



## FEP Medical Policy Manual

### FEP 2.01.04 Hyperbaric Oxygen Therapy

**Effective Policy Date: April 1, 2023**

**Original Policy Date: September 2012**

**Related Policies:**

None

## Hyperbaric Oxygen Therapy

### Description

#### Description

Hyperbaric oxygen therapy (HBOT) involves breathing 100% oxygen at pressures between 1.5 and 3.0 atmospheres. It is generally applied systemically with the patient inside a hyperbaric chamber. HBOT can also be applied topically; ie, the body part to be treated is isolated (eg, in an inflatable bag and exposed to pure oxygen). HBOT has been investigated for various conditions that have potential to respond to increased oxygen delivery to tissue.

#### OBJECTIVE

The objective of this evidence review is to determine whether the use of systemic hyperbaric oxygen pressurization improves net health outcomes for a variety of indications.

## POLICY STATEMENT

Systemic hyperbaric oxygen pressurization may be considered **medically necessary** in the treatment of the following conditions:

- nonhealing diabetic wounds of the lower extremities in patients who meet the following 3 criteria:
  - Patient has type 1 or type 2 diabetes and has a lower-extremity wound due to diabetes;
  - Patient has a wound classified as Wagner grade 3 or higher (see Policy Guidelines section); and
  - Patient has no measurable signs of healing after 30 days of an adequate course of standard wound therapy;
- acute traumatic ischemia (eg, crush injuries, reperfusion injury, compartment syndrome);
- decompression sickness;
- gas embolism, acute;
- cyanide poisoning, acute;
- acute carbon monoxide poisoning;
- soft-tissue radiation necrosis (eg, radiation enteritis, cystitis, proctitis) and osteoradionecrosis;
- pre- and posttreatment for patients undergoing dental surgery (non-implant-related) of an irradiated jaw;
- gas gangrene (ie, clostridial myonecrosis);
- profound anemia with exceptional blood loss: only when blood transfusion is impossible or must be delayed; and
- chronic refractory osteomyelitis.

Hyperbaric oxygen pressurization is considered **investigational** in all other situations, including but not limited to, the treatment of the following conditions:

- compromised skin grafts or flaps;
- acute osteomyelitis;
- bisphosphonate-related osteonecrosis of the jaw;
- necrotizing soft tissue infections;
- acute thermal burns;
- acute surgical and traumatic wounds;
- chronic wounds, other than those in patients with diabetes who meet the criteria specified in the medically necessary statement;
- spinal cord injury;
- traumatic brain injury;
- inflammatory bowel disease (Crohn disease or ulcerative colitis);
- brown recluse spider bites;
- bone grafts;

- carbon tetrachloride poisoning, acute;
- cerebrovascular disease, acute (thrombotic or embolic) or chronic;
- fracture healing;
- hydrogen sulfide poisoning;
- intra-abdominal and intracranial abscesses;
- lepromatous leprosy;
- meningitis;
- pseudomembranous colitis (antimicrobial agent-induced colitis);
- radiation myelitis;
- sickle cell crisis and/or hematuria;
- demyelinating diseases (eg, multiple sclerosis, amyotrophic lateral sclerosis);
- retinal artery insufficiency, acute;
- retinopathy, adjunct to scleral buckling procedures in patients with sickle cell peripheral retinopathy and retinal detachment;
- pyoderma gangrenosum;
- acute arterial peripheral insufficiency;
- acute coronary syndromes and as an adjunct to coronary interventions, including but not limited to, percutaneous coronary interventions and cardiopulmonary bypass;
- idiopathic sudden sensorineural hearing loss;
- refractory mycoses: mucormycosis, actinomycosis, conidiobolus coronato;
- cerebral edema, acute;
- migraine;
- in vitro fertilization;
- cerebral palsy;
- tumor sensitization for cancer treatments, including but not limited to, radiotherapy or chemotherapy;
- delayed-onset muscle soreness;
- idiopathic femoral neck necrosis;
- chronic arm lymphedema following radiotherapy for cancer;
- radiation-induced injury in the head and neck, except as noted earlier in the medically necessary statement;
- early treatment (beginning at completion of radiotherapy) to reduce adverse events of radiotherapy;
- autism spectrum disorder;
- Bell palsy;
- acute ischemic stroke;
- motor dysfunction associated with stroke;

- herpes zoster;
- vascular dementia;
- fibromyalgia; and
- mental illness (ie, posttraumatic stress disorder, generalized anxiety disorder or depression).

## POLICY GUIDELINES

### Systemic Hyperbaric Oxygen

The Wagner classification system categorizes wounds as follows: grade 0, no open lesion; grade 1, superficial ulcer without penetration to deeper layers; grade 2, ulcer penetrates to tendon, bone, or joint; grade 3, lesion has penetrated deeper than grade 2, and there is abscess, osteomyelitis, pyarthrosis, plantar space abscess, or infection of the tendon and tendon sheaths; grade 4, wet or dry gangrene in the toes or forefoot; grade 5, gangrene involves the whole foot or such a percentage that no local procedures are possible and amputation (at least at the below the knee level) is indicated.

Following are recommended indications from the Undersea and Hyperbaric Medical Society's (UHMS) 2019 Hyperbaric Oxygen Therapy Committee report on utilization of HBOT (14th edition):

- Air or gas embolism
- Carbon monoxide poisoning and carbon monoxide complicated by cyanide poisoning
- Clostridial myositis and myonecrosis (gas gangrene)
- Crush injury, compartment syndrome, and other acute traumatic ischemias
- Decompression sickness
- Central retinal artery occlusion
- Diabetic foot ulcer
- Healing of other problem wounds
- Severe anemia
- Intracranial abscess
- Necrotizing soft tissue infections
- Refractory osteomyelitis
- Delayed radiation injury (soft tissue and bony necrosis)
- Compromised grafts and flaps
- Acute thermal burn injury
- Sudden sensorineural hearing loss.

## BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

## FDA REGULATORY STATUS

Since 1979, the U.S. Food and Drug Administration (FDA) has cleared multiple topical and systemic hyperbaric oxygen administration devices through the 510(k) pathway. In 2013, the FDA published a statement warning that non-FDA approved uses of HBOT may endanger the health of patients.<sup>2</sup> If patients mistakenly believe that HBOT devices have been proven safe for uses not cleared by the FDA, they may delay or forgo proven medical therapies.

## RATIONALE

### Summary of Evidence

For individuals with chronic diabetic ulcers who receive systemic hyperbaric oxygen therapy (HBOT), the evidence includes randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are symptoms and change in disease status. Meta-analyses of RCTs found significantly higher diabetic ulcer healing rates with HBOT than with control conditions. One of the 2 meta-analyses found that HBOT was associated with a significantly lower rate of major amputation. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with carbon monoxide poisoning who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are overall survival (OS) and symptoms. A meta-analysis in a Cochrane review of low-quality RCT data did not find HBOT to be associated with a significantly lower risk of neurologic deficits after carbon monoxide poisoning. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with radionecrosis, osteoradionecrosis, or treatment of irradiated jaw who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms and change in disease status. A meta-analysis in a Cochrane review of RCTs found evidence that HBOT improved radionecrosis and osteoradionecrosis outcomes and resulted in better outcomes before tooth extraction in an irradiated jaw. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with chronic refractory osteomyelitis who receive systemic HBOT, the evidence includes case series. Relevant outcomes are symptoms and change in disease status. The case series reported high rates of successful outcomes (no drainage, pain, tenderness, or cellulitis) in patients with chronic refractory osteomyelitis treated with HBOT. However, controlled studies are needed to determine conclusively the impact of HBOT on health outcomes compared with other interventions. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with acute thermal burns who receive systemic HBOT, the evidence includes a systematic review of 2 RCTs. Relevant outcomes are overall survival (OS), symptoms, and change in disease status. Both RCTs were judged to have poor methodologic quality. Evidence from well-conducted controlled trials is needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with acute surgical and traumatic wounds who receive systemic HBOT, the evidence includes RCTs, controlled nonrandomized studies, and systematic reviews. Relevant outcomes are OS, symptoms, change in disease status, and functional outcomes. There was considerable heterogeneity across the 4 RCTs identified (eg, patient population, comparison group, treatment regimen, outcomes). This heterogeneity prevented pooling of trial findings and limits the ability to conclude the impact of HBOT on health outcomes for patients with acute surgical and traumatic wounds. Additional evidence from high-quality RCTs is needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with bisphosphonate-related osteonecrosis of the jaw who receive systemic HBOT, the evidence includes an randomized controlled trial (RCT). Relevant outcomes are symptoms and change in disease status. The RCT was unblinded and reported initial benefits at 3-month follow-up; however, there were no significant benefits of HBOT for most health outcomes compared with standard care in the long-term (6 months to 2 years). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with necrotizing soft tissue infections who receive systemic HBOT, the evidence includes systematic reviews and a retrospective cohort study. Relevant outcomes are OS, symptoms, and change in disease status. A Cochrane review did not identify any RCTs. Another systematic review identified a retrospective cohort study, which did not find better outcomes after HBOT than after standard care without HBOT in patients with necrotizing soft tissue infections. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with acute coronary syndrome who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are OS, symptoms, change in disease status, and functional outcomes. A Cochrane review identified 6 RCTs. There were 2 pooled analyses, 1 found significantly lower rates of death with HBOT and the other reported inconsistent results in left ventricular function. Additional RCT data are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with acute ischemic stroke who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are OS, symptoms, change in disease status, and functional outcomes. Cochrane reviewers could only pool data for a single outcome (mortality at 3-6 months), and for that outcome, there was no significant difference between active and sham HBOT treatments. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with motor dysfunction associated with stroke who receive systemic HBOT, the evidence includes an RCT. Relevant outcomes are symptoms and functional outcomes. The RCT, which used a crossover design, found better outcomes with HBOT at 2 months than with delayed treatment. However, the trial had a number of methodologic limitations (eg, lack of patient blinding, heterogeneous population, high dropout rate) that make it difficult to evaluate the efficacy of HBOT. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with Bell palsy who receive systemic HBOT, the evidence includes a systematic review. Relevant outcomes are symptoms, change in disease status, and functional outcomes. A Cochrane review did not identify any RCTs meeting selection criteria; the single RCT found did not have a blinded outcome assessment. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with traumatic brain injury who receive systemic HBOT, the evidence includes RCTs and systematic reviews. Relevant outcomes are OS, symptoms, change in disease status, and functional outcomes. RCTs were heterogeneous regarding intervention protocols, patient populations, and outcomes reported. Multiple RCTs of US military service members showed no statistical difference in outcomes between HBOT groups and those that received sham treatment. Systematic reviews conducted pooled analyses only on a minority of the published RCTs, and these findings were inconsistent. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with inflammatory bowel disease who receive systemic HBOT, the evidence includes an RCT, observational studies, and a systematic review. Relevant outcomes are symptoms, change in disease status and functional outcomes. One small RCT has been published, and this trial did not find a significant improvement in health outcomes when HBOT was added to standard medical therapy. A systematic review including the RCT and observational studies found a high rate of bias in the literature due to attrition and reporting bias. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

A Cochrane review of RCTs had mixed findings from studies that included individuals with tinnitus. Some outcomes (ie, improvement in hearing of all frequencies, >25% return of hearing) were better with HBOT than with a control intervention, but more than 50% return of hearing did not differ significantly between groups. There was important variability in the patients enrolled in the studies. A subsequent systematic review had similarly limited conclusions due to the inclusion of non-randomized studies. One RCT included in this review included patients with idiopathic sudden sensorineural hearing loss and found no differences in HBOT treatment compared with steroid injections in mean hearing thresholds at 0.25, 0.5, 1, and 4 kHz; however, a significant difference was detected at the 2-kHz level. Nonrandomized studies of HBOT used as adjunctive therapy did not support incremental value, although 1 systematic review evaluated HBOT along with steroid therapy and found benefit specifically for those with severe-to-profound idiopathic sudden sensorineural hearing loss. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with delayed-onset muscle soreness who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms and functional outcomes. A Cochrane review of RCTs found worse short-term pain outcomes with HBOT than with control and no difference in longer-term pain or other outcomes (eg, swelling). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with autism spectrum disorder who receive systemic HBOT, the evidence includes an RCT and a systematic review. Relevant outcomes are symptoms and functional outcomes. A Cochrane review identified a single RCT on HBOT for autism spectrum disorder and this trial did not find significantly better parental-assessed or clinician-assessed outcomes with HBOT compared with sham. A subsequent controlled trial reached the same conclusion. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with cerebral palsy who receive systemic HBOT, the evidence includes 2 RCTs and an observational study. Relevant outcomes are symptoms and functional outcomes. One RCT was stopped early due to futility, and the other did not find significantly better outcomes with HBOT than with a sham intervention. The observational study focused on sleep disorders in children with cerebral palsy and reported improvements with the HBOT treatment. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with vascular dementia who receive systemic HBOT, the evidence includes an RCT and a systematic review. Relevant outcomes are symptoms and functional outcomes. The Cochrane review identified only a single RCT with methodologic limitations. Well-conducted controlled trials are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with radiotherapy adverse events who receive systemic HBOT, the evidence includes RCTs, nonrandomized comparator trials, case series, and systematic reviews. Relevant outcomes are symptoms and functional outcomes. Three systematic reviews included few RCTs and provide limited evidence on the effect of HBOT. Two RCTs identified had inconsistent findings. One reported no short-term benefit with HBOT, but some benefits 12 months after radiotherapy; the other did not find a significant benefit of HBOT at 12-month follow-up. Another RCT assessed HBOT for radiation-induced cystitis and found significant benefit by some measures but not others. An observational study for dry mouth (xerostomia) caused by radiotherapy found some benefit with HBOT. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with idiopathic femoral neck necrosis who receive systemic HBOT, the evidence includes an RCT. Relevant outcomes are symptoms, change in disease status, and functional outcomes. The RCT, which had a small sample, only reported short-term (ie, 6-week) outcomes. Larger well-conducted RCTs reporting longer-term outcomes are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with a migraine who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms, change in disease status, and functional outcomes. The Cochrane review conducted a pooled analysis including 3 of the 11 trials. Meta-analysis of these 3 RCTs found significantly greater relief of migraine symptoms with HBOT than with a comparator intervention within 45 minutes of treatment. Longer-term data are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with herpes zoster who receive systemic HBOT, the evidence includes an RCT. Relevant outcomes are symptoms and change in disease status. The RCT was unblinded and only reported short-term (ie, 6-week) outcomes. Additional well-conducted RCTs with longer follow-up are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with fibromyalgia who receive systemic HBOT, the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, and functional outcomes. Only 2 RCTs were identified, and both reported positive effects of HBOT on tender points and pain. However, the trials had relatively small samples and methodologic limitations (eg, quasi-randomization, no or uncertain sham control for a condition with subjective outcomes susceptible to a placebo effect). Moreover, the HBOT protocols varied. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with multiple sclerosis who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms and functional outcomes. A Cochrane review of RCTs did not find a significant difference in Expanded Disability Status Scale scores when patients with multiple sclerosis were treated with HBOT versus a comparator intervention. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with cancer and are undergoing chemotherapy who receive systemic HBOT, the evidence includes an RCT and a systematic review. Relevant outcomes are OS and change in disease status. While the systematic review reported improvements in tumor control in patients with head and neck cancer who received HBOT, the adverse events accompanying the treatment (eg, radiation tissue injury, seizures) were significant. The single RCT did not find a significant difference in survival for cancer patients who received HBOT before chemotherapy compared with usual care. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

## SUPPLEMENTAL INFORMATION

### Practice Guidelines and Position Statements

#### Infectious Disease Society of America

In 2012, the Infectious Disease Society of America published guidelines on the diagnosis and treatment of diabetic foot infections.<sup>74</sup> The guidelines stated that "for selected diabetic foot wounds that are slow to heal, clinicians might consider using hyperbaric oxygen therapy (strength of evidence: strong; quality of evidence: moderate)."

#### Society of Vascular Surgery et al

In 2016, the Society of Vascular Surgery in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine published guidelines on the management of the diabetic foot.<sup>75</sup> According to the guidelines, for diabetic foot ulcers that fail to demonstrate improvement (>50% wound area reduction) after a minimum of 4 weeks of standard wound therapy, adjunctive therapy such as HBOT is recommended (grade 1B). Also, for diabetic foot ulcers with adequate perfusion that fail to respond to 4 to 6 weeks of conservative management, HBOT is suggested (grade 2B).



## Undersea and Hyperbaric Medical Society

In 2015, the Undersea and Hyperbaric Medical Society (UHMS) published guidelines on the use of hyperbaric oxygen therapy (HBOT) for treating diabetic foot ulcers.<sup>76</sup> Recommendations in the current version include:

- Suggest against using HBOT in patients with "Wagner Grade 2 or lower diabetic foot ulcers..."
- Suggest adding HBOT in patients with "Wagner Grade 3 or higher diabetic foot ulcers that have not shown significant improvement after 30 days of [standard of care] therapy..."
- Suggest "adding acute post-operative hyperbaric oxygen therapy to the standard of care" in patients with "Wagner Grade 3 or higher diabetic foot ulcers" who have just had foot surgery related to their diabetic ulcers.

The 2019 UHMS hyperbaric oxygen therapy indications committee report included the following indications as recommended:<sup>77</sup>

1. Air or Gas Embolism
2. Carbon Monoxide Poisoning and carbon monoxide complicated by cyanide poisoning
3. Clostridial Myositis and Myonecrosis (Gas Gangrene)
4. Crush Injury, Compartment Syndrome and Other Acute Traumatic Ischemias
5. Decompression Sickness
6. Central retinal artery occlusion
7. Diabetic foot ulcer
8. Healing of other problem wounds
9. Severe anemia
10. Intracranial abscess
11. Necrotizing soft tissue infections
12. Refractory osteomyelitis
13. Delayed radiation injury ( soft tissue and bony necrosis)
14. Compromised grafts and flaps
15. Acute thermal burn injury
16. Sudden Sensorineural hearing loss.

The UHMS has also published position statements that concluded there was insufficient evidence to recommend topical HBOT for chronic wounds (2005),<sup>78</sup> multiple sclerosis,<sup>79</sup> and autism spectrum disorder (2009).<sup>80</sup>

## American Academy of Otolaryngology-Head and Neck Surgery

In 2018, the American Academy of Otolaryngology-Head and Neck Surgery updated clinical guidelines on treatment of sudden hearing loss.<sup>81</sup> They give the following options regarding HBOT:

"Clinicians may offer, or refer to a physician who can offer, hyperbaric oxygen therapy (HBOT) combined with steroid therapy within 2 weeks of onset of SSNH."

"Clinicians may offer, or refer to a physician who can offer, hyperbaric oxygen therapy (HBOT) combined with steroid therapy as salvage within 1 months of onset of SSNHL."

## U.S. Preventive Services Task Force Recommendations

Not applicable.

## Medicare National Coverage

In 2003, the Centers for Medicare & Medicaid added Medicare coverage of HBOT for diabetic wounds of the lower extremities meeting certain criteria. As of the current coverage statement, Medicare coverage is provided for HBOT administered in a chamber for the following conditions<sup>82</sup>:

1. "Acute carbon monoxide intoxication,
2. Decompression illness,
3. Gas embolism,
4. Gas gangrene,
5. Acute traumatic peripheral ischemia. HBO therapy is a valuable adjunctive treatment to be used in combination with accepted standard therapeutic measures when loss of function, limb, or life is threatened.
6. Crush injuries and suturing of severed limbs. As in the previous conditions, HBO therapy would be an adjunctive treatment when loss of function, limb, or life is threatened.
7. Progressive necrotizing infections (necrotizing fasciitis),
8. Acute peripheral arterial insufficiency,
9. Preparation and preservation of compromised skin grafts (not for primary management of wounds),
10. Chronic refractory osteomyelitis, unresponsive to conventional medical and surgical management,
11. Osteoradionecrosis as an adjunct to conventional treatment,
12. Soft tissue radionecrosis as an adjunct to conventional treatment,
13. Cyanide poisoning,
14. Actinomycosis, only as an adjunct to conventional therapy when the disease process is refractory to antibiotics and surgical treatment,
15. Diabetic wounds of the lower extremities in patients who meet the following 3 criteria:
  1. Patient has type I or type II diabetes and has a lower extremity wound that is due to diabetes;
  2. Patient has a wound classified as Wagner grade III or higher; and
  3. Patient has failed an adequate course of standard wound therapy."

The use of HBO therapy is covered as adjunctive therapy only after there are no measurable signs of healing for at least 30-days of treatment with standard wound therapy and must be used in addition to standard wound care. Standard wound care in patients with diabetic wounds includes: assessment of a patient's vascular status and correction of any vascular problems in the affected limb if possible, optimization of nutritional status, optimization of glucose control, debridement by any means to remove devitalized tissue, maintenance of a clean, moist bed of granulation tissue with appropriate moist dressings, appropriate off-loading, and necessary treatment to resolve any infection that might be present. Failure to respond to standard wound care occurs when there are no measurable signs of healing for at least 30 consecutive days. Wounds must be evaluated at least every 30 days during administration of HBO therapy. Continued treatment with HBO therapy is not covered if measurable signs of healing have not been demonstrated within any 30-day period of treatment."

Systemic HBOT for other indications is not covered, nor is topical HBOT for any indication.

## REFERENCES

1. Sadri RA, Cooper JS. Hyperbaric, complications. NCBI Bookshelf 2017; <https://www.ncbi.nlm.nih.gov/books/NBK459191/>. Accessed November 14, 2020.
2. U.S. Food and Drug Administration. Hyperbaric Oxygen Therapy: Don't Be Misled. 2013; <https://www.talkingaboutthescience.com/studies/FDA2013.pdf>. Accessed December 15, 2020.
3. de Smet GHJ, Kroese LF, Menon AG, et al. Oxygen therapies and their effects on wound healing. *Wound Repair Regen*. Aug 2017; 25(4): 591-608. PMID 28783878
4. Leslie CA, Sapico FL, Ginunas VJ, et al. Randomized controlled trial of topical hyperbaric oxygen for treatment of diabetic foot ulcers. *Diabetes Care*. Feb 1988; 11(2): 111-5. PMID 3289861
5. Kranke P, Bennett MH, Martyn-St James M, et al. Hyperbaric oxygen therapy for chronic wounds. *Cochrane Database Syst Rev*. Jun 24 2015; (6): CD004123. PMID 26106870
6. Elraiyah T, Tsapas A, Prutsky G, et al. A systematic review and meta-analysis of adjunctive therapies in diabetic foot ulcers. *J Vasc Surg*. Feb 2016; 63(2 Suppl): 46S-58S.e1-2. PMID 26804368
7. Buckley NA, Juurlink DN, Isbister G, et al. Hyperbaric oxygen for carbon monoxide poisoning. *Cochrane Database Syst Rev*. Apr 13 2011; (4): CD002041. PMID 21491385
8. Nakajima M, Aso S, Matsui H, et al. Hyperbaric oxygen therapy and mortality from carbon monoxide poisoning: A nationwide observational study. *Am J Emerg Med*. Feb 2020; 38(2): 225-230. PMID 30797609
9. Bennett MH, Feldmeier J, Hampson NB, et al. Hyperbaric oxygen therapy for late radiation tissue injury. *Cochrane Database Syst Rev*. Apr 28 2016; 4: CD005005. PMID 27123955
10. Borab Z, Mirmanesh MD, Gantz M, et al. Systematic review of hyperbaric oxygen therapy for the treatment of radiation-induced skin necrosis. *J Plast Reconstr Aesthet Surg*. Apr 2017; 70(4): 529-538. PMID 28081957
11. Ravi P, Vaishnavi D, Gnanam A, et al. The role of hyperbaric oxygen therapy in the prevention and management of radiation-induced complications of the head and neck - a systematic review of literature. *J Stomatol Oral Maxillofac Surg*. Dec 2017; 118(6): 359-362. PMID 28838774
12. Maynor ML, Moon RE, Camporesi EM, et al. Chronic osteomyelitis of the tibia: treatment with hyperbaric oxygen and autogenous microsurgical muscle transplantation. *J South Orthop Assoc*. 1998; 7(1): 43-57. PMID 9570731
13. Davis JC, Heckman JD, DeLee JC, et al. Chronic non-hematogenous osteomyelitis treated with adjuvant hyperbaric oxygen. *J Bone Joint Surg Am*. Oct 1986; 68(8): 1210-7. PMID 3771602
14. Chen CE, Ko JY, Fu TH, et al. Results of chronic osteomyelitis of the femur treated with hyperbaric oxygen: a preliminary report. *Chang Gung Med J*. Feb 2004; 27(2): 91-7. PMID 15095953
15. Chen CE, Shih ST, Fu TH, et al. Hyperbaric oxygen therapy in the treatment of chronic refractory osteomyelitis: a preliminary report. *Chang Gung Med J*. Feb 2003; 26(2): 114-21. PMID 12718388
16. Chen CY, Lee SS, Chan YS, et al. Chronic refractory tibia osteomyelitis treated with adjuvant hyperbaric oxygen: a preliminary report. *Changgeng Yi Xue Za Zhi*. Jun 1998; 21(2): 165-71. PMID 9729650
17. Villanueva E, Bennett MH, Wasiak J, et al. Hyperbaric oxygen therapy for thermal burns. *Cochrane Database Syst Rev*. 2004; (3): CD004727. PMID 15266540
18. Eskes A, Vermeulen H, Lucas C, et al. Hyperbaric oxygen therapy for treating acute surgical and traumatic wounds. *Cochrane Database Syst Rev*. Dec 16 2013; (12): CD008059. PMID 24343585
19. Dauwe PB, Pulikkottil BJ, Lavery L, et al. Does hyperbaric oxygen therapy work in facilitating acute wound healing: a systematic review. *Plast Reconstr Surg*. Feb 2014; 133(2): 208e-215e. PMID 24469192
20. Freiburger JJ, Padilla-Burgos R, McGraw T, et al. What is the role of hyperbaric oxygen in the management of bisphosphonate-related osteonecrosis of the jaw: a randomized controlled trial of hyperbaric oxygen as an adjunct to surgery and antibiotics. *J Oral Maxillofac Surg*. Jul 2012; 70(7): 1573-83. PMID 22698292

21. Levett D, Bennett MH, Millar I. Adjunctive hyperbaric oxygen for necrotizing fasciitis. *Cochrane Database Syst Rev*. Jan 15 2015; 1: CD007937. PMID 25879088
22. Jallali N, Withey S, Butler PE. Hyperbaric oxygen as adjuvant therapy in the management of necrotizing fasciitis. *Am J Surg*. Apr 2005; 189(4): 462-6. PMID 15820462
23. George ME, Rueth NM, Skarda DE, et al. Hyperbaric oxygen does not improve outcome in patients with necrotizing soft tissue infection. *Surg Infect (Larchmt)*. Feb 2009; 10(1): 21-8. PMID 18991520
24. Bennett MH, Lehm JP, Jepson N. Hyperbaric oxygen therapy for acute coronary syndrome. *Cochrane Database Syst Rev*. Jul 23 2015; (7): CD004818. PMID 26202854
25. Bennett MH, Weibel S, Wasiak J, et al. Hyperbaric oxygen therapy for acute ischaemic stroke. *Cochrane Database Syst Rev*. Nov 12 2014; (11): CD004954. PMID 25387992
26. Efrati S, Fishlev G, Bechor Y, et al. Hyperbaric oxygen induces late neuroplasticity in post stroke patients--randomized, prospective trial. *PLoS One*. 2013; 8(1): e53716. PMID 23335971
27. Holland NJ, Bernstein JM, Hamilton JW. Hyperbaric oxygen therapy for Bell's palsy. *Cochrane Database Syst Rev*. Feb 15 2012; (2): CD007288. PMID 22336830
28. Wang F, Wang Y, Sun T, et al. Hyperbaric oxygen therapy for the treatment of traumatic brain injury: a meta-analysis. *Neurol Sci*. May 2016; 37(5): 693-701. PMID 26746238
29. Crawford C, Teo L, Yang E, et al. Is Hyperbaric Oxygen Therapy Effective for Traumatic Brain Injury? A Rapid Evidence Assessment of the Literature and Recommendations for the Field. *J Head Trauma Rehabil*. May/June 2017; 32(3): E27-E37. PMID 27603765
30. Bennett MH, Trytko B, Jonker B. Hyperbaric oxygen therapy for the adjunctive treatment of traumatic brain injury. *Cochrane Database Syst Rev*. Dec 12 2012; 12: CD004609. PMID 23235612
31. Hart BB, Weaver LK, Gupta A, et al. Hyperbaric oxygen for mTBI-associated PCS and PTSD: Pooled analysis of results from Department of Defense and other published studies. *Undersea Hyperb Med. BIMA* 2019; 46(3): 353-383. PMID 31394604
32. Wolf G, Cifu D, Baugh L, et al. The effect of hyperbaric oxygen on symptoms after mild traumatic brain injury. *J Neurotrauma*. Nov 20 2012; 29(17): 2606-12. PMID 23031217
33. Cifu DX, Walker WC, West SL, et al. Hyperbaric oxygen for blast-related postconcussion syndrome: three-month outcomes. *Ann Neurol*. Feb 2014; 75(2): 277-86. PMID 24255008
34. Miller RS, Weaver LK, Bahraini N, et al. Effects of hyperbaric oxygen on symptoms and quality of life among service members with persistent postconcussion symptoms: a randomized clinical trial. *JAMA Intern Med*. Jan 2015; 175(1): 43-52. PMID 25401463
35. Marois P, Mukherjee A, Ballaz L. Hyperbaric Oxygen Treatment for Persistent Postconcussion Symptoms--A Placebo Effect?. *JAMA Intern Med*. Jul 2015; 175(7): 1239-40. PMID 26146912
36. mTBI mechanisms of action of HBO2 for persistent post-concussive symptoms. U.S. National Library of Medicine. *ClinicalTrials.gov*. <https://clinicaltrials.gov/ct2/show/NCT01611194>. Updated November 21, 2018. Accessed November 14, 2020.
37. Hart BB, Wilson SH, Churchill S, et al. Extended follow-up in a randomized trial of hyperbaric oxygen for persistent post-concussive symptoms. *Undersea Hyperb Med. BIMA* 2019; 46(3): 313-327. PMID 31394601
38. Weaver LK, Churchill S, Wilson SH, et al. A composite outcome for mild traumatic brain injury in trials of hyperbaric oxygen. *Undersea Hyperb Med. BIMA* 2019; 46(3): 341-352. PMID 31394603
39. Hyperbaric oxygen therapy (HBO2) for persistent post-concussive symptoms after mild traumatic brain injury (mTBI) (HOPPS). U.S. National Library of Medicine. *ClinicalTrials.gov*. <https://clinicaltrials.gov/ct2/show/NCT01306968>. Updated September 5, 2014. Accessed November 14, 2020.
40. Churchill S, Deru K, Weaver LK, et al. Adverse events and blinding in two randomized trials of hyperbaric oxygen for persistent post-concussive symptoms. *Undersea Hyperb Med. BIMA* 2019; 46(3): 331-340. PMID 31394602
41. Dulai PS, Gleeson MW, Taylor D, et al. Systematic review: The safety and efficacy of hyperbaric oxygen therapy for inflammatory bowel disease. *Aliment Pharmacol Ther*. Jun 2014; 39(11): 1266-75. PMID 24738651
42. Pagoldh M, Hultgren E, Arnell P, et al. Hyperbaric oxygen therapy does not improve the effects of standardized treatment in a severe attack of ulcerative colitis: a prospective randomized study. *Scand J Gastroenterol*. Sep 2013; 48(9): 1033-40. PMID 23879825
43. Lewis JD, Chuai S, Nessel L, et al. Use of the noninvasive components of the Mayo score to assess clinical response in ulcerative colitis. *Inflamm Bowel Dis*. Dec 2008; 14(12): 1660-6. PMID 18623174
44. Bennett MH, Kertesz T, Perleth M, et al. Hyperbaric oxygen for idiopathic sudden sensorineural hearing loss and tinnitus. *Cochrane Database Syst Rev*. Oct 17 2012; 10: CD004739. PMID 23076907
45. Rhee TM, Hwang D, Lee JS, et al. Addition of Hyperbaric Oxygen Therapy vs Medical Therapy Alone for Idiopathic Sudden Sensorineural Hearing Loss: A Systematic Review and Meta-analysis. *JAMA Otolaryngol Head Neck Surg*. Dec 01 2018; 144(12): 1153-1161. PMID 30267033
46. Eryigit B, Ziyilan F, Yaz F, et al. The effectiveness of hyperbaric oxygen in patients with idiopathic sudden sensorineural hearing loss: a systematic review. *Eur Arch Otorhinolaryngol*. Dec 2018; 275(12): 2893-2904. PMID 30324404
47. Cvorovic L, Jovanovic MB, Milutinovic Z, et al. Randomized prospective trial of hyperbaric oxygen therapy and intratympanic steroid injection as salvage treatment of sudden sensorineural hearing loss. *Otol Neurotol*. Aug 2013; 34(6): 1021-6. PMID 23820795
48. Sun H, Qiu X, Hu J, et al. Comparison of intratympanic dexamethasone therapy and hyperbaric oxygen therapy for the salvage treatment of refractory high-frequency sudden sensorineural hearing loss. *Am J Otolaryngol*. Sep 2018; 39(5): 531-535. PMID 29891394
49. Almosnino G, Holm JR, Schwartz SR, et al. The Role of Hyperbaric Oxygen as Salvage Therapy for Sudden Sensorineural Hearing Loss. *Ann Otol Rhinol Laryngol*. Oct 2018; 127(10): 672-676. PMID 30009614
50. Xie S, Qiang Q, Mei L, et al. Multivariate analysis of prognostic factors for idiopathic sudden sensorineural hearing loss treated with adjuvant hyperbaric oxygen therapy. *Eur Arch Otorhinolaryngol*. Jan 2018; 275(1): 47-51. PMID 29071444

51. Bennett M, Best TM, Babul S, et al. Hyperbaric oxygen therapy for delayed onset muscle soreness and closed soft tissue injury. *Cochrane Database Syst Rev.* Oct 19 2005; (4): CD004713. PMID 16235376
52. Xiong T, Chen H, Luo R, et al. Hyperbaric oxygen therapy for people with autism spectrum disorder (ASD). *Cochrane Database Syst Rev.* Oct 13 2016; 10: CD010922. PMID 27737490
53. Sampanthavivat M, Singkhwa W, Chaiyakul T, et al. Hyperbaric oxygen in the treatment of childhood autism: a randomised controlled trial. *Diving Hyperb Med.* Sep 2012; 42(3): 128-33. PMID 22987458
54. Rizzato A, D'Alessandro N, Berenci E, et al. Effect of mild hyperbaric oxygen therapy on children diagnosed with autism. *Undersea Hyperb Med.* Nov-Dec 2018; 45(6): 639-645. PMID 31158930
55. Lacey DJ, Stolfi A, Pilati LE. Effects of hyperbaric oxygen on motor function in children with cerebral palsy. *Ann Neurol.* Nov 2012; 72(5): 695-703. PMID 23071074
56. Collet JP, Vanasse M, Marois P, et al. Hyperbaric oxygen for children with cerebral palsy: a randomised multicentre trial. *HBO-CP Research Group. Lancet.* Feb 24 2001; 357(9256): 582-6. PMID 11558483
57. Long Y, Tan J, Nie Y, et al. Hyperbaric oxygen therapy is safe and effective for the treatment of sleep disorders in children with cerebral palsy. *Neurol Res.* Mar 2017; 39(3): 239-247. PMID 28079475
58. Xiao Y, Wang J, Jiang S, et al. Hyperbaric oxygen therapy for vascular dementia. *Cochrane Database Syst Rev.* Jul 11 2012; (7): CD009425. PMID 22786527
59. Spiegelberg L, Djasim UM, van Neck HW, et al. Hyperbaric oxygen therapy in the management of radiation-induced injury in the head and neck region: a review of the literature. *J Oral Maxillofac Surg.* Aug 2010; 68(8): 1732-9. PMID 20493616
60. Villeirs L, Taily T, Ost P, et al. Hyperbaric oxygen therapy for radiation cystitis after pelvic radiotherapy: Systematic review of the recent literature. *Int J Urol.* Feb 2020; 27(2): 98-107. PMID 31617263
61. Teguh DN, Levendag PC, Noever I, et al. Early hyperbaric oxygen therapy for reducing radiotherapy side effects: early results of a randomized trial in oropharyngeal and nasopharyngeal cancer. *Int J Radiat Oncol Biol Phys.* Nov 01 2009; 75(3): 711-6. PMID 19386439
62. Sherlock S, Way M, Tabah A. Hyperbaric oxygen treatment for the management of radiation-induced xerostomia. *J Med Imaging Radiat Oncol.* Dec 2018; 62(6): 841-846. PMID 30113763
63. Gothard L, Haviland J, Bryson P, et al. Randomised phase II trial of hyperbaric oxygen therapy in patients with chronic arm lymphoedema after radiotherapy for cancer. *Radiother Oncol.* Oct 2010; 97(1): 101-7. PMID 20605648
64. Oscarsson N, Muller B, Rosen A, et al. Radiation-induced cystitis treated with hyperbaric oxygen therapy (RICH-ART): a randomised, controlled, phase 2-3 trial. *Lancet Oncol.* Nov 2019; 20(11): 1602-1614. PMID 31537473
65. Camporesi EM, Vezzani G, Bosco G, et al. Hyperbaric oxygen therapy in femoral head necrosis. *J Arthroplasty.* Sep 2010; 25(6 Suppl): 118-23. PMID 20637561
66. Bennett MH, French C, Schnabel A, et al. Normobaric and hyperbaric oxygen therapy for the treatment and prevention of migraine and cluster headache. *Cochrane Database Syst Rev.* Dec 28 2015; (12): CD005219. PMID 26709672
67. Peng Z, Wang S, Huang X, et al. Effect of hyperbaric oxygen therapy on patients with herpes zoster. *Undersea Hyperb Med.* Nov-Dec 2012; 39(6): 1083-7. PMID 23342765
68. Efrati S, Golan H, Bechor Y, et al. Hyperbaric oxygen therapy can diminish fibromyalgia syndrome—prospective clinical trial. *PLoS One.* 2015; 10(5): e0127012. PMID 26010952
69. Yildiz S, Kiralp MZ, Akin A, et al. A new treatment modality for fibromyalgia syndrome: hyperbaric oxygen therapy. *J Int Med Res.* May-Jun 2004; 32(3): 263-7. PMID 15174219
70. Bennett M, Heard R. Hyperbaric oxygen therapy for multiple sclerosis. *CNS Neurosci Ther.* Apr 2010; 16(2): 115-24. PMID 20415839
71. Bennett M, Feldmeier J, Smee R, et al. Hyperbaric oxygenation for tumour sensitisation to radiotherapy. *Cochrane Database Syst Rev.* Oct 19 2005; (4): CD005007. PMID 16235387
72. Bennett MH, Feldmeier J, Smee R, et al. Hyperbaric oxygenation for tumour sensitisation to radiotherapy. *Cochrane Database Syst Rev.* Apr 18 2012; (4): CD005007. PMID 22513926
73. Heys SD, Smith IC, Ross JA, et al. A pilot study with long term follow up of hyperbaric oxygen pretreatment in patients with locally advanced breast cancer undergoing neo-adjuvant chemotherapy. *Undersea Hyperb Med.* Jan-Feb 2006; 33(1): 33-43. PMID 16602255
74. Lipsky BA, Berendt AR, Cornia PB, et al. 2012 infectious diseases society of america clinical practice guideline for the diagnosis and treatment of diabetic foot infections. *J Am Podiatr Med Assoc.* Jan-Feb 2013; 103(1): 2-7. PMID 23328846
75. Hingorani A, LaMuraglia GM, Henke P, et al. The management of diabetic foot: A clinical practice guideline by the Society for Vascular Surgery in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine. *J Vasc Surg.* Feb 2016; 63(2 Suppl): 3S-21S. PMID 26804367
76. Huang ET, Mansouri J, Murad MH, et al. A clinical practice guideline for the use of hyperbaric oxygen therapy in the treatment of diabetic foot ulcers. *Undersea Hyperb Med.* May-Jun 2015; 42(3): 205-47. PMID 26152105
77. Moon RE, editor. *Hyperbaric Oxygen Therapy Indications.* 14th ed. North Palm Beach, FL: Undersea and Hyperbaric Medical Society; 2019.
78. Feldmeier JJ, Hopf HW, Warriner RA, et al. UHMS position statement: topical oxygen for chronic wounds. *Undersea Hyperb Med.* May-Jun 2005; 32(3): 157-68. PMID 16119307
79. Bennett M., Heard R. UHMS Position Paper: the treatment of multiple sclerosis with hyperbaric oxygen therapy. North Palm Beach, FL: Undersea & Hyperbaric Medical Society (UHMS); n.d.
80. Bennett M., B. H. UHMS Position Paper: the treatment of autism spectrum disorder with hyperbaric oxygen therapy. North Palm Beach, FL: Undersea & Hyperbaric Medical Society (UHMS); 2009.
81. Chandrasekhar SS, Tsai Do BS, Schwartz SR, et al. Clinical Practice Guideline: Sudden Hearing Loss (Update). *Otolaryngol Head Neck Surg.* Aug 2019; 161(1\_suppl): S1-S45. PMID 31369359
82. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for Hyperbaric Oxygen Therapy (20.29). 2006; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?ncid=12&ver=3>. Accessed November 14, 2020.

**POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:**

<b>Date</b>	<b>Action</b>	<b>Description</b>
September 2012	New Policy	
March 2014	Replace Policy	Policy updated with literature review. Bisphosphonate-related osteonecrosis of the jaw, motor dysfunction associated with stroke, herpes zoster and vascular dementia added as not medically necessary. References added; other references renumbered or removed. Additional conditions added as medically necessary.
December 2014	Replace Policy	Policy updated with literature review through July 17, 2014. In not medically necessary statement, severe or refractory Crohn's disease changed to inflammatory bowel disease (Crohn disease or ulcerative colitis). Clarification added to bullet point in not medically necessary statement on radiation-induced injury in the head and neck. Title changed from "Hyperbaric Oxygen Pressurization (HBO)" to "Hyperbaric Oxygen Therapy". References 2, 10, 12-13, 43-46, and 49 added.
December 2015	Replace Policy	Policy updated with literature review through June 17, 2015; references 27, 34, 40, 41 and 66-70 added. Bullet points on (1) fibromyalgia and (2) mental illness (i.e., posttraumatic stress disorder, generalized anxiety disorder or depression) added to the not medically necessary statement
March 2017	Replace Policy	Policy updated with literature review through November 8, 2016; references 6-7, 15, 22, 26-27, 39, 48, and 56 added. Policy statements unchanged.
March 2018	Replace Policy	Policy updated with literature review through November 6, 2017; references 1, 3, 12-13, 47, 58-59, 62-63, 65-67, and 69-70 added. Policy statements unchanged.
March 2019	Replace Policy	Policy updated with literature review through November 6, 2017; references 1, 3, 12-13, 47, 58-59, 62-63, 65-67, and 69-70 added. Policy statements unchanged.
March 2020	Replace Policy	Policy updated with literature review through November 11, 2019; references added. Policy statements unchanged except "not medically necessary" changed to "investigational" due to HBOT FDA 501(k) clearance.
March 2022	Replace policy	Policy updated with literature review through November 14, 2020; references added. Policy statements unchanged.
March 2023	Replace Policy	Policy updated with literature review through November 30, 2021; references added. Minor editorial change to policy statements; intent unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.