

5.21.022

Section:	Prescription Drugs	Effective Date:	April 1, 2024
Subsection:	Antineoplastic Agents	Original Policy Date:	October 17, 2012
Subject:	Bosulif	Page:	1 of 5

Last Review Date: March 8, 2024

Bosulif

Description

Bosulif (bosutinib)

Background

Bosulif (bosutinib) is a tyrosine kinase inhibitor indicated for the treatment of chronic myelogenous leukemia (CML). Bosulif is intended for patients with Philadelphia chromosome positive CML (Ph+ CML) who are newly-diagnosed or resistant to or who cannot tolerate other therapies, including imatinib. Bosulif inhibits the BCR-ABL kinase, an enzyme that promotes chronic myelogenous leukemia (CML) (1-2).

Regulatory Status

FDA-approved indications: Bosulif is a kinase inhibitor indicated for the treatment of: (1)

1. Adult and pediatric patients 1 year of age and older with chronic phase Ph+ chronic myelogenous leukemia (CML), newly-diagnosed or resistant or intolerant to prior therapy.
2. Adult patients with accelerated, or blast phase Ph+ chronic myelogenous leukemia (CML) with resistance or intolerance to prior therapy.

Off-Label Uses: (2-4)

1. Follow-up therapy for CML patients after hematopoietic stem cell transplant (HSCT)
2. Relapsed or refractory Ph+ acute lymphoblastic leukemia (ALL)

Bosulif has warnings and precautions regarding gastrointestinal toxicity, myelosuppression, hepatic toxicity, fluid retention and embryo-fetal toxicity. Patients should be monitored and managed using standards of care. Therapy should be interrupted, the dose reduced or discontinued as necessary (1).

Section:	Prescription Drugs	Effective Date:	April 1, 2024
Subsection:	Antineoplastic Agents	Original Policy Date:	October 17, 2012
Subject:	Bosulif	Page:	2 of 5

Liver enzymes should be monitored at least monthly for the first 3 months and as needed. Thrombocytopenia, anemia and neutropenia can occur; therefore, a complete blood count should be performed weekly for the first month and then monthly or as clinically indicated (1).

The safety and efficacy of Bosulif in patients less than 1 year of age have not been established (1).

Related policies

Gleevec, Iclusig, Scemblix, Sprycel, Tassigna

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Bosulif may be considered **medically necessary** if the conditions indicated below are met.

Bosulif may be considered **investigational** for all other indications.

Prior-Approval Requirements

Diagnoses

Patient must have **ONE** of the following:

1. Newly-diagnosed chronic phase Ph+ chronic myelogenous leukemia (CML)
 - a. 1 year of age or older
2. Chronic phase Ph+ chronic myelogenous leukemia (CML)
 - a. 1 year of age or older
 - b. Resistant or intolerant to prior therapy
3. Accelerated phase Ph+ chronic myelogenous leukemia (CML)
 - a. 18 years of age or older
 - b. Resistant or intolerant to prior therapy
4. Blast phase Ph+ chronic myelogenous leukemia (CML)
 - a. 18 years of age or older
 - b. Resistant or intolerant to prior therapy
5. Chronic myeloid leukemia (CML) post-hematopoietic stem cell transplant (HSCT)
 - a. 18 years of age or older

Section:	Prescription Drugs	Effective Date:	April 1, 2024
Subsection:	Antineoplastic Agents	Original Policy Date:	October 17, 2012
Subject:	Bosulif	Page:	3 of 5

6. Relapsed or refractory Ph+ acute lymphoblastic leukemia (ALL)
 - a. 18 years of age or older

AND ALL of the following for **ALL** indications:

1. Confirmed by detection of the Ph chromosome or BCR-ABL gene by cytogenetic and/or molecular testing prior to initiation of therapy
2. If the patient has had prior therapy with a TKI then **ONE** of the following requirements must be met:
 - a. Member experienced resistance to prior therapy with TKI
 - i. Results from mutational testing are negative for the T315I mutation
 - b. Member experienced toxicity or intolerance to prior therapy with a TKI

Prior – Approval *Renewal* Requirements

Diagnoses

Patient must have **ONE** of the following:

1. Chronic phase Ph+ chronic myelogenous leukemia (CML)
 - a. 1 year of age or older
2. Accelerated phase Ph+ chronic myelogenous leukemia (CML)
 - a. 18 years of age or older
3. Blast phase Ph+ chronic myelogenous leukemia (CML)
 - a. 18 years of age or older
4. Chronic myeloid leukemia (CML) post-hematopoietic stem cell transplant (HSCT)
 - a. 18 years of age or older
5. Relapsed or refractory Ph+ acute lymphoblastic leukemia (ALL)
 - a. 18 years of age or older

Policy Guidelines

Pre - PA Allowance

None

Prior - Approval Limits

Quantity 600 mg per day

Duration 12 months

Section:	Prescription Drugs	Effective Date:	April 1, 2024
Subsection:	Antineoplastic Agents	Original Policy Date:	October 17, 2012
Subject:	Bosulif	Page:	4 of 5

Prior – Approval *Renewal* Limits

Same as above

Rationale

Summary

Bosulif (bosutinib) is a kinase inhibitor that inhibits the BCR-ABL kinase, an enzyme that promotes chronic myelogenous leukemia (CML). In studies, treatment with bosutinib reduced the size of CML tumors relative to controls and inhibited growth of murine myeloid tumors expressing several imatinib-resistant forms of BCR-ABL. Bosulif has warnings and precautions regarding gastrointestinal toxicity, myelosuppression, hepatic toxicity, fluid retention and embryo-fetal toxicity (1).

Prior approval is required to ensure the safe, clinically appropriate, and cost-effective use of Bosulif while maintaining optimal therapeutic outcomes.

References

1. Bosulif [package insert]. New York, NY: Pfizer Labs; September 2023.
2. NCCN Drugs & Biologics Compendium® Bosutinib 2024. National Comprehensive Cancer Network, Inc. Accessed on January 22, 2024.
3. NCCN Clinical Practice Guidelines in Oncology® Chronic Myeloid Leukemia (CML) (Version 2.2024). National Comprehensive Cancer Network, Inc. December 2023. Accessed on January 22, 2024.
4. NCCN Clinical Practice Guidelines in Oncology® Acute Lymphoblastic Leukemia (Version 3.2023). National Comprehensive Cancer Network, Inc. October 2023. Accessed on January 22, 2024.

Policy History

Date	Action
October 2012	New addition
March 2013	Annual review and update.
September 2014	Annual editorial review and reference update
December 2015	Annual editorial review and reference update Removed tyrosine kinase inhibitors examples
June 2016	Annual editorial review and reference update Policy number change from 5.04.22 to 5.21.22

5.21.022

Section:	Prescription Drugs	Effective Date:	April 1, 2024
Subsection:	Antineoplastic Agents	Original Policy Date:	October 17, 2012
Subject:	Bosulif	Page:	5 of 5

March 2017	Annual review and reference update Addition of no dual therapy with another tyrosine kinase inhibitor and addition of the age requirement in the renewal section
January 2018	Addition of new indication of newly-diagnosed chronic phase Ph+ chronic myelogenous leukemia (CML), chronic myeloid leukemia (CML) post hematopoietic stem cell transplant (HSCT), Ph+ acute lymphoblastic leukemia (ALL); and accelerated, or blast phase Ph+ chronic myelogenous leukemia (CML) with no prior therapy Addition of quantity limits
March 2018	Annual editorial review Addition of mutational testing requirement to “If the patient has had prior therapy with a TKI then ONE of the following requirements must be met: member experienced resistance to prior therapy with TKI and results from mutational testing are negative for the T315I mutation or member experienced toxicity or intolerance to prior therapy with a TKI
June 2019	Annual review and reference update
June 2020	Annual editorial review and reference update. Removed no dual therapy with another TKI requirement
March 2021	Annual review and reference update
March 2022	Annual editorial review and reference update
December 2022	Annual review and reference update. Changed policy number to 5.21.022
March 2023	Annual review and reference update
October 2023	Per PI update, changed age requirement to 1 year or older for chronic phase CML. Added “resistant or intolerant to prior therapy” to accelerated and blast phase CML. Revised quantity limit to 600 mg per day
December 2023	Annual review
March 2024	Annual review and reference update

Keywords

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on March 8, 2024 and is effective on April 1, 2024.