



FEP Medical Policy Manual

FEP 2.04.153 Tumor-Informed Circulating Tumor DNA Testing for Cancer Management

Effective Policy Date: July 1, 2022

Original Policy Date: July 2022

Related Policies:

None

Tumor-Informed Circulating Tumor DNA Testing for Cancer Management

Description

Description

This evidence review addresses the use of tumor-informed circulating tumor DNA (ctDNA) testing for cancer management. The purpose of tumor-informed ctDNA testing in individuals with cancer is to predict disease course to inform treatment decisions and to monitor for recurrence following treatment.

OBJECTIVE

The objective of this evidence review is to determine whether using tumor-informed ctDNA testing improves the net health outcome in individuals with bladder, breast, colorectal, esophageal, or non-small cell lung cancer, or in individuals with solid tumors who are receiving immunotherapy.

POLICY STATEMENT

Tumor-informed circulating tumor DNA testing (e.g., Signatera) is considered **not medically necessary** for all indications.

POLICY GUIDELINES

None

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

FDA REGULATORY STATUS

Signatera is a laboratory developed test regulated under CLIA. Signatera has been developed and its performance characteristics determined by Natera, the CLIA-certified laboratory performing the test. The test has not been cleared or approved by the US Food and Drug Administration (FDA), but has received 3 Breakthrough Device Designations from FDA:

- In May 2019, Signatera was granted a BDD for the detection of ctDNA in localized or advanced colorectal cancer patients to optimize the use of chemotherapy alone or in combination with durvalumab.
- A March 2021 press release announced that FDA granted 2 additional Breakthrough Device Designations covering new intended uses.¹

RATIONALE

Summary of Evidence

For individuals with colorectal cancer (CRC) who receive tumor-informed circulating tumor DNA (ctDNA) testing with Signatera to guide treatment decisions and monitor for recurrence, the evidence includes 3 noncomparative studies (N = 410) and 1 retrospective comparative study (N = 48). Relevant outcomes are overall survival, disease-specific survival, test validity, other test performance measures, change in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related mortality. Nonrandomized studies have reported an association between ctDNA results measured at diagnosis, following surgery, during adjuvant therapy, and during surveillance after curative treatment and prognosis, but these studies are limited by a lack of comparison to tests used for the same purpose, imprecise estimates due to small sample sizes, and clinical heterogeneity of study populations. No study reported management changes made in response to ctDNA test results. A retrospective observational study found no advantage to surveillance with Signatera compared to standard surveillance conducted according to National Comprehensive Cancer Network (NCCN) guidelines ($p > .99$ for sensitivity and specificity compared to imaging). There is no direct evidence that the use of the test improves health outcomes, and indirect evidence is not sufficient to draw conclusions about clinical validity. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with breast cancer who receive tumor-informed ctDNA testing with Signatera to guide treatment decisions and monitor for recurrence, the evidence includes 2 noncomparative studies (N = 133). Relevant outcomes are overall survival, disease-specific survival, test validity, other test performance measures, change in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related mortality. One study evaluated Signatera testing for disease surveillance following primary treatment, and 1 reported the association of test results at different timepoints with response to neoadjuvant chemotherapy. Although the studies found an association of test results with prognosis, the studies are limited by a lack of comparison to tests used for the same purpose, imprecise estimates due to small sample sizes, and clinical heterogeneity of study populations. No study reported management changes made in response to ctDNA test results. There is no direct evidence that the use of the test improves health outcomes, and indirect evidence is not sufficient to draw conclusions about clinical validity. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with bladder cancer who receive tumor-informed ctDNA testing with Signatera to guide treatment decisions and monitor for recurrence, the evidence includes 1 uncontrolled prospective cohort study (N = 68) and 1 retrospective subgroup analysis from a randomized controlled trial (RCT) (N = 581). Relevant outcomes are overall survival, disease-specific survival, test validity, other test performance measure, change in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related mortality. The prospective study reported an association between Signatera test results at diagnosis, during chemotherapy treatment, and during surveillance following cystectomy to prognosis. The retrospective analysis reported an association between test results and response to atezolizumab treatment. Study limitations, including a lack of comparison to tests used for the same purpose preclude drawing conclusions about clinical validity and usefulness. No study reported management changes made in response to ctDNA test results. There is no direct evidence that the use of the test improves health outcomes, and indirect evidence is not sufficient to draw conclusions about clinical validity. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with non-small cell lung cancer (NSCLC) who receive tumor-informed ctDNA testing with Signatera to guide treatment decisions and monitor for recurrence, the evidence includes 1 subgroup analysis of participants enrolled in a prospective observational study (N = 24). Relevant outcomes are overall survival, disease-specific survival, test validity, other test performance measures, change in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related mortality. Of 14 individuals with confirmed relapse, 13 (93%) had a positive ctDNA test (defined as at least 2 single-nucleotide variants detected). Of 10 individuals with no relapse after a median follow up of 775 days, (range 688 to 945 days), 1 had a positive ctDNA test (10%). This study's small sample size and lack of a comparator preclude drawing conclusions about clinical validity. There is no direct evidence that the use of the test improves health outcomes, and indirect evidence is not sufficient to draw conclusions about clinical validity. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with esophageal cancer who receive tumor-informed ctDNA testing with Signatera to guide treatment decisions and monitor for recurrence, the evidence includes 1 noncomparative, retrospective study (N = 17). Relevant outcomes are overall survival, disease-specific survival, test validity, other test performance measure, change in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related mortality. Patients who were ctDNA-positive before surgery had significantly poorer disease-free survival (DFS) ($p < .042$), with a median DFS of 32.0 months versus 63.0 months in ctDNA-negative preoperative patients. This study was limited by its small number sample size and retrospective design. There is no direct evidence that the use of the test improves health outcomes. Due to the study's limitations and lack of additional supporting studies, the evidence is not sufficient to draw conclusions on clinical validity. Additionally, the management pathway for Signatera testing in esophageal cancer has not been clearly defined. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with solid tumors who receive tumor-informed ctDNA testing with Signatera to monitor response to immunotherapy, the evidence includes a subgroup analysis of individuals enrolled in a nonrandomized trial of pembrolizumab (N = 106). Relevant outcomes are overall survival, disease-specific survival, test validity, other test performance measures, change in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related mortality. The subgroup analysis evaluated Signatera testing to monitor response to immunotherapy in individuals with advanced solid tumors who were enrolled in a Phase II clinical trial of pembrolizumab. Lower-than-median ctDNA levels at baseline were associated with improved overall survival (adjusted hazard ratio [HR] 0.49, 95% CI 0.29 to 0.83) and progression free survival (adjusted HR 0.54, 95% CI 0.34 to 0.85). The study was limited by a small sample size, variability in results across different tumor types, and lack of a comparison to standard methods of monitoring response to treatment. There is no direct evidence that the use of the test improves health outcomes, and indirect evidence is not sufficient to draw conclusions about clinical validity. Additionally, the management pathway for Signatera testing for monitoring response to immunotherapy has not been clearly defined. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

National Comprehensive Cancer Network

National Comprehensive Cancer Network guidelines do not specifically address tumor-informed ctDNA testing for any of the cancer types included in this review. The guidelines on colon cancer state: "The panel believes that there are insufficient data to recommend the use of multigene assays, Immunoscore or post-surgical ctDNA to estimate risk of recurrence or determine adjuvant therapy."¹³

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

REFERENCES

1. Natera. FDA Grants Two New Breakthrough Device Designations for Nateras Signatera MRD Test. March 24, 2021. <https://www.natera.com/company/news/fda-grants-two-new-breakthrough-device-designations-for-nateras-signatera-mrd-test/>. Accessed April 5, 2022
2. Fakih M, Sandhu J, Wang C, et al. Evaluation of Comparative Surveillance Strategies of Circulating Tumor DNA, Imaging, and Carcinoembryonic Antigen Levels in Patients With Resected Colorectal Cancer. *JAMA Netw Open*. Mar 01 2022; 5(3): e221093. PMID 35258578
3. Reinert T, Henriksen TV, Christensen E, et al. Analysis of Plasma Cell-Free DNA by Ultradeep Sequencing in Patients With Stages I to III Colorectal Cancer. *JAMA Oncol*. Aug 01 2019; 5(8): 1124-1131. PMID 31070691
4. Henriksen TV, Tarazona N, Frydendahl A, et al. Circulating Tumor DNA in Stage III Colorectal Cancer, beyond Minimal Residual Disease Detection, toward Assessment of Adjuvant Therapy Efficacy and Clinical Behavior of Recurrences. *Clin Cancer Res*. Feb 01 2022; 28(3): 507-517. PMID 34625408
5. Loupakis F, Sharma S, Derouazi M, et al. Detection of Molecular Residual Disease Using Personalized Circulating Tumor DNA Assay in Patients With Colorectal Cancer Undergoing Resection of Metastases. *JCO Precis Oncol*. Jul 2021; 5. PMID 34327297
6. Coombes RC, Page K, Salari R, et al. Personalized Detection of Circulating Tumor DNA Antedates Breast Cancer Metastatic Recurrence. *Clin Cancer Res*. Jul 15 2019; 25(14): 4255-4263. PMID 30992300
7. Magbanua MJM, Swigart LB, Wu HT, et al. Circulating tumor DNA in neoadjuvant-treated breast cancer reflects response and survival. *Ann Oncol*. Feb 2021; 32(2): 229-239. PMID 33232761
8. Christensen E, Birkenkamp-Demtroder K, Sethi H, et al. Early Detection of Metastatic Relapse and Monitoring of Therapeutic Efficacy by Ultra-Deep Sequencing of Plasma Cell-Free DNA in Patients With Urothelial Bladder Carcinoma. *J Clin Oncol*. Jun 20 2019; 37(18): 1547-1557. PMID 31059311
9. Powles T, Assaf ZJ, Davarpanah N, et al. ctDNA guiding adjuvant immunotherapy in urothelial carcinoma. *Nature*. Jul 2021; 595(7867): 432-437. PMID 34135506
10. Abbosh C, Birkbak NJ, Wilson GA, et al. Phylogenetic ctDNA analysis depicts early-stage lung cancer evolution. *Nature*. Apr 26 2017; 545(7655): 446-451. PMID 28445469
11. Ococks E, Sharma S, Ng AWT, et al. Serial Circulating Tumor DNA Detection Using a Personalized, Tumor-Informed Assay in Esophageal Adenocarcinoma Patients Following Resection. *Gastroenterology*. Nov 2021; 161(5): 1705-1708.e2. PMID 34284036
12. Bratman SV, Yang SYC, lafolla MAJ, et al. Personalized circulating tumor DNA analysis as a predictive biomarker in solid tumor patients treated with pembrolizumab. *Nat Cancer*. Sep 2020; 1(9): 873-881. PMID 35121950
13. National Comprehensive Cancer Network. Guidelines on Colon Cancer. Version 1.2022. https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf. Accessed April 5, 2022.

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
June 2022	New policy - Add to Pathology/Laboratory section	Policy created with literature review through March 11, 2022. Tumor-informed circulating tumor DNA testing is considered investigational for all indications.