



FEP Medical Policy Manual

FEP 4.01.19 Laparoscopic, Percutaneous, and Transcervical Techniques for Uterine Fibroid Myolysis

Effective Policy Date: July 1, 2022

Original Policy Date: September 2013

Related Policies:

7.01.109 - Magnetic Resonance Imaging-Guided Focused Ultrasound

Laparoscopic, Percutaneous, and Transcervical Techniques for Uterine Fibroid Myolysis

Description

Description

Various minimally invasive treatments for uterine fibroids have been proposed as alternatives to surgery. Among these approaches are laparoscopic, percutaneous, and transcervical techniques to induce myolysis, which includes radiofrequency ablation (RFA), laser and bipolar needles, cryomyolysis, and magnetic resonance imaging-guided laser ablation.

OBJECTIVE

The objective of this evidence review is to determine whether laparoscopic, percutaneous, and/or transcervical techniques to induce myolysis improve the net health outcome in individuals with uterine fibroids.

POLICY STATEMENT

Laparoscopic or transcervical radiofrequency ablation (RFA) as a treatment of symptomatic uterine fibroids is considered **medically necessary** in women 18 years and older when ALL of the following conditions are met:

- Evidence of uterine fibroids via ultrasound that are less than 10 cm in diameter for laparoscopic RFA with Acessa or 7 cm for transcervical RFA with Sonata; AND
- Patient desires a uterine-sparing treatment approach or is ineligible for hysterectomy or other uterine-sparing alternatives to RFA (e.g., laparoscopic myomectomy, uterine artery embolization [UAE]) (see Policy Guidelines); AND
- Patient has experienced at least 1 of the following symptoms that are a direct result of the fibroid(s):
 - Menorrhagia or other abnormal uterine bleeding that interferes with daily activities or causes anemia (see Policy Guidelines);
 - Pelvic pain or pressure;
 - Urinary symptoms (e.g., urinary frequency, urgency) related to bulk compression of the bladder;
 - Gastrointestinal symptoms related to bulk compression of the bowel (e.g., constipation, bloating);
 - Dyspareunia (painful or difficult sexual relations).

Other laparoscopic, transcervical, or percutaneous techniques for myolysis of uterine fibroids, including use of laser or bipolar needles, cryomyolysis, and magnetic resonance imaging-guided laser ablation, are considered **investigational**.

POLICY GUIDELINES

Eligibility Considerations

Abnormal uterine bleeding refers to uterine bleeding of abnormal frequency, duration, and volume that interferes with a patient's quality of life. Patients with abnormal uterine bleeding with an inadequate response to appropriately selected medical therapy may be considered for alternate uterine-sparing interventions. In patients >45 years of age with menorrhagia or other abnormal bleeding, endometrial biopsy is recommended prior to treatment to rule out endometrial malignancy and/or additional assessment to rule out a risk for uterine leiomyosarcoma.

Clinical trial experience with radiofrequency ablation (RFA) has been limited to patients with overall uterine size ≤ 16 gestational weeks size based on pelvic examination. In patients where fibroids cannot be distinguished from adenomyosis on ultrasound, advanced imaging (e.g., magnetic resonance imaging [MRI]) may be required. For patients with pelvic pain, alternative causes such as endometritis and active pelvic inflammatory disease should be excluded prior to treatment with RFA.

Treatment Approach Considerations for Radiofrequency Ablation

Uterine fibroids are categorized according to the International Federation of Gynaecology and Obstetrics (FIGO) leiomyoma subclassification system (see Table PG1). Choice of laparoscopic versus transcervical RFA treatment is dependent on fibroid number, size, type and location, and patient preferences. For example, predominantly lower uterine segment or cervical leiomyomata, or those with a predominant submucosal location or intramural FIGO type 2 or 3 fibroids, may suggest a transcervical approach, whereas fibroids with largely fundal or extramural components may suggest a laparoscopic approach. Patients aiming to avoid future deliveries via obligate cesarean section may prefer a transcervical approach. Select women with numerous fibroids may benefit from combined laparoscopic RFA and laparoscopic myomectomy. Patients with intramural fibroids, intra-abdominal adhesions, or medical contraindications may not be candidates for alternative uterine-sparing interventions.

Table PG1. FIGO Leiomyoma Subclassification System

Group	Type	Description
Submucosal	0	Pedunculated intracavitary
	1	<50% intramural ($\geq 50\%$ submucosal)

	2	≥50% intramural (<50% submucosal)
Other	3	100% intramural, contacting endometrium
	4	100% intramural, no endometrial or subserosal contact
	5	Subserosal, ≥50% intramural
	6	Subserosal, <50% intramural
	7	Pedunculated subserosal
	8	Non-myometrial location (eg, cervical, broad ligament, parasitic)
Hybrid	X-X	Both submucosal and subserosal components. Submucosal component designated by first number and subserosal component designated by second number.

FIGO: International Federation of Gynaecology and Obstetrics.

Table adapted from Gomez et al (2021). MRI-based pictorial review of the FIGO classification system for uterine fibroids. *Abdom Radiol.* 46(5): 2146-2155. PMID: 33385249.

Reinterventions

Reintervention with RFA may be considered for patients meeting policy criteria with documentation of new or recurrent fibroid development following a partial response with the initial procedure. However, data on reinterventions for new or recurrent fibroids is limited and documentation procedures for repeat anatomic mapping of fibroids are not standardized.

In November 2014, the U.S. Food and Drug Administration published a safety communication on laparoscopic power morcellators used for myomectomy and hysterectomy in most women. (Morcellators are not otherwise addressed herein). The Administration recommended that manufacturers of these devices include in their product labels a boxed safety warning and wording on contraindications (see <https://www.fda.gov/downloads/MedicalDevices/DeviceRegulationandGuidance/GuidanceDocuments/UCM424123.pdf>).

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

Some of the lysis procedures are specialized and not widely disseminated; therefore, requests for out-of-network referral may occur.

FDA REGULATORY STATUS

In 2012, the Acessa™ System (Acessa Health, formerly Halt Medical) was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process for percutaneous laparoscopic coagulation and ablation of soft tissue and treatment of symptomatic uterine fibroids under laparoscopic ultrasound guidance (K121858). The technology was previously approved in 2010, at which time it was called the Halt 2000GI™ Electrosurgical Radiofrequency Ablation System. In 2014, the ultrasound guidance system received marketing clearance from the FDA (K132744). FDA product code: GEI. In 2018, the third-generation Acessa™ ProVu System was cleared for marketing by the FDA through the 510(k) process for use in percutaneous, laparoscopic coagulation and ablation of soft tissue, including treatment of symptomatic uterine fibroids under laparoscopic ultrasound guidance. (K181124). Hologic acquired Accessa Health in 2020. FDA product code: HFG.

In 2018, the Sonata Sonography-Guided Transcervical Fibroid Ablation System (Gynesonics) was cleared for marketing by the FDA through the 510(k) process for diagnostic intrauterine imaging and transcervical radiofrequency ablation as treatment of symptomatic uterine fibroids (K173703). The Sonata System 2.1 received marketing clearance in 2020 (K193516) and the Sonata System 2.2 received marketing clearance in 2021 (K211535). The Sonata system was previously known as Vizablate. FDA product codes: KNF, ITX, and IYO.

Cryoablation is a surgical procedure that uses previously approved and available cryoablation systems; and as a surgical procedure, it is not subject to regulation by the FDA. Other products addressed in this review (eg, Nd:YAG lasers, bipolar electrodes) have long-standing FDA approval, and there

are no products specifically approved for the treatment of uterine fibroids.

RATIONALE

Summary of Evidence

For individuals who have symptomatic uterine fibroids who receive radiofrequency ablation (RFA), the evidence includes prospective cohorts, randomized controlled trials (RCTs), and systematic reviews. Relevant outcomes are symptoms, quality of life, and treatment-related morbidity. The meta-analysis found low rates of reintervention with RFA and quality of life outcomes that were similar to uterine artery embolization and myomectomy at 12 months. Data on reintervention rates at 36 months were limited to 1 RCT and 1 cohort study with high loss to follow-up. No studies reported reintervention rates at 60 months. Two RCTs found that RFA was noninferior to laparoscopic myomectomy on the primary outcome: length of hospitalization. A number of secondary outcomes were reported at 12 and 24 months in 1 RCT, including symptoms and quality of life. None of the secondary outcomes demonstrated significant between-group differences in a subgroup analysis of 43 patients. The procedure has faster recovery than myomectomy, and provides a reduction in symptoms and improvement in quality of life in the short term. Recurrence and reintervention rates at longer follow-up are unknown. Well-designed comparative trials with longer follow-up are needed to determine the effect of RFA on health outcomes compared with other treatment options such as myomectomy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have symptomatic uterine fibroids who receive laser or bipolar needles, the evidence includes case series. Relevant outcomes are symptoms, quality of life, and treatment-related morbidity. The case series were published in the 1990s, and the procedures used then may not reflect current practice. RCTs comparing laser or bipolar needles with alternative treatments for uterine fibroids are needed to evaluate the safety and efficacy of this technology adequately. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have symptomatic uterine fibroids who receive cryomyolysis, the evidence includes case series. Relevant outcomes are symptoms, quality of life, and treatment-related morbidity. Among the few case series, sample sizes were small (≤ 20 patients). RCTs comparing cryomyolysis with alternative treatments for uterine fibroids are needed to evaluate the safety and efficacy of this technology adequately. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have symptomatic uterine fibroids who receive magnetic resonance imaging-guided laser ablation, the evidence includes a study with historical controls. Relevant outcomes are symptoms, quality of life, and treatment-related morbidity. A single study with historical controls is not sufficiently robust to evaluate this technology. RCTs comparing magnetic resonance imaging-guided laser ablation with alternative treatments for uterine fibroids are needed to evaluate safety and efficacy adequately. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Obstetricians and Gynecologists

In 2021, the American College of Obstetricians and Gynecologists updated its practice bulletin on the management of symptomatic leiomyomas.¹ Recommendations based on a review of evidence included the following:

- Radiofrequency ablation can be considered as a minimally invasive treatment option in patients who desire to retain their uterus, provided they are counseled about the limited data on reproductive outcomes. Laparoscopic, transvaginal, or transcervical approaches using ultrasound guidance are considered similarly effective.
- Focused ultrasound is associated with a reduction in leiomyoma and uterine size, but is associated with less improvement in symptoms and quality of life and a higher risk of reintervention compared with uterine artery embolization.

- Myomectomy was recommended as an option in patients who desire uterine preservation or future pregnancy and are counseled about the risk of recurrence. The laparoscopic approach is associated with shorter hospitalization, less postoperative pain, faster return to work, and earlier return to normal activities.
- Hysterectomy is recommended as a definitive surgical management option in patients who do not desire future childbearing or do not wish to retain their uterus.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

REFERENCES

1. Stewart MR, Adelman VL, Jacoby MR. Management of Symptomatic Uterine Leiomyomas: ACOG Practice Bulletin, Number 228. *Obstet Gynecol.* Jun 01 2021; 137(6): e100-e115. PMID 34011888
2. Laughlin-Tommaso SK, Jacoby VL, Myers ER. Disparities in Fibroid Incidence, Prognosis, and Management. *Obstet Gynecol Clin North Am.* Mar 2017; 44(1): 81-94. PMID 28160895
3. Stewart EA, Nicholson WK, Bradley L, et al. The burden of uterine fibroids for African-American women: results of a national survey. *J Womens Health (Larchmt).* Oct 2013; 22(10): 807-16. PMID 24033092
4. Jones S, O'Donovan P, Toub D. Radiofrequency ablation for treatment of symptomatic uterine fibroids. *Obstet Gynecol Int.* 2012; 2012: 194839. PMID 21961009
5. Davis MR, Soliman AM, Castelli-Haley J, et al. Reintervention Rates After Myomectomy, Endometrial Ablation, and Uterine Artery Embolization for Patients with Uterine Fibroids. *J Womens Health (Larchmt).* Oct 2018; 27(10): 1204-1214. PMID 30085898
6. Sandberg EM, Tummers FHMP, Cohen SL, et al. Reintervention risk and quality of life outcomes after uterine-sparing interventions for fibroids: a systematic review and meta-analysis. *Fertil Steril.* Apr 2018; 109(4): 698-707.e1. PMID 29653718
7. Havryliuk Y, Setton R, Carlow JJ, et al. Symptomatic Fibroid Management: Systematic Review of the Literature. *JLS.* Jul-Sep 2017; 21(3). PMID 28951653
8. Lin L, Ma H, Wang J, et al. Quality of Life, Adverse Events, and Reintervention Outcomes after Laparoscopic Radiofrequency Ablation for Symptomatic Uterine Fibroids: A Meta-Analysis. *J Minim Invasive Gynecol.* Mar 2019; 26(3): 409-416. PMID 30253997
9. Bradley LD, Pasic RP, Miller LE. Clinical Performance of Radiofrequency Ablation for Treatment of Uterine Fibroids: Systematic Review and Meta-Analysis of Prospective Studies. *J Laparoendosc Adv Surg Tech A.* Dec 2019; 29(12): 1507-1517. PMID 31702440
10. Arnreiter C, Oppelt P. A Systematic Review of the Treatment of Uterine Myomas Using Transcervical Ultrasound-Guided Radiofrequency Ablation with the Sonata System. *J Minim Invasive Gynecol.* Aug 2021; 28(8): 1462-1469. PMID 33892184
11. Brucker SY, Hahn M, Kraemer D, et al. Laparoscopic radiofrequency volumetric thermal ablation of fibroids versus laparoscopic myomectomy. *Int J Gynaecol Obstet.* Jun 2014; 125(3): 261-5. PMID 24698202
12. Rattray DD, Weins L, Regush LC, et al. Clinical outcomes and health care utilization pre- and post-laparoscopic radiofrequency ablation of symptomatic fibroids and laparoscopic myomectomy: a randomized trial of uterine-sparing techniques (TRUST) in Canada. *Clinicoecon Outcomes Res.* 2018; 10: 201-212. PMID 29670382
13. Yu S, Silverberg K, Bhagavath B, et al. Post-Market Safety of Laparoscopic Ultrasound-Guided Radiofrequency Ablation. *JLS.* Oct-Dec 2020; 24(4). PMID 33510567
14. Hahn M, Brucker S, Kraemer D, et al. Radiofrequency Volumetric Thermal Ablation of Fibroids and Laparoscopic Myomectomy: Long-Term Follow-up From a Randomized Trial. *Geburtshilfe Frauenheilkd.* May 2015; 75(5): 442-449. PMID 26097247
15. Kramer B, Hahn M, Taran FA, et al. Interim analysis of a randomized controlled trial comparing laparoscopic radiofrequency volumetric thermal ablation of uterine fibroids with laparoscopic myomectomy. *Int J Gynaecol Obstet.* May 2016; 133(2): 206-11. PMID 26892690
16. Jacoby VL, Parvataneni R, Oberman E, et al. Laparoscopic Radiofrequency Ablation of Uterine Leiomyomas: Clinical Outcomes during Early Adoption into Surgical Practice. *J Minim Invasive Gynecol.* May 2020; 27(4): 915-925. PMID 31376584
17. Miller CE, Osman KM. Transcervical Radiofrequency Ablation of Symptomatic Uterine Fibroids: 2-Year Results of the SONATA Pivotal Trial. *J Gynecol Surg.* Dec 01 2019; 35(6): 345-349. PMID 32226268
18. Lukes A, Green MA. Three-Year Results of the SONATA Pivotal Trial of Transcervical Fibroid Ablation for Symptomatic Uterine Myomata. *J Gynecol Surg.* Oct 01 2020; 36(5): 228-233. PMID 33061253
19. Brolmann H, Bongers M, Garza-Leal JG, et al. The FAST-EU trial: 12-month clinical outcomes of women after intrauterine sonography-guided transcervical radiofrequency ablation of uterine fibroids. *Gynecol Surg.* NA 2016; 13: 27-35. PMID 26918001
20. Shifrin G, Engelhardt M, Gee P, et al. Transcervical fibroid ablation with the Sonata system for treatment of submucous and large uterine fibroids. *Int J Gynaecol Obstet.* Oct 2021; 155(1): 79-85. PMID 33544889

21. Christoffel L, Romer T, Schiermeier S. Transcervical Radiofrequency Ablation of Uterine Fibroids Global Registry (SAGE): Study Protocol and Preliminary Results. *Med Devices (Auckl)*. 2021; 14: 77-84. PMID 33688276
22. Keltz J, Levie M, Chudnoff S. Pregnancy Outcomes After Direct Uterine Myoma Thermal Ablation: Review of the Literature. *J Minim Invasive Gynecol*. May 2017; 24(4): 538-545. PMID 28109894
23. Berman JM, Shashoua A, Olson C, et al. Case Series of Reproductive Outcomes after Laparoscopic Radiofrequency Ablation of Symptomatic Myomas. *J Minim Invasive Gynecol*. Mar 2020; 27(3): 639-645. PMID 31238151
24. Goldfarb HA. Bipolar laparoscopic needles for myoma coagulation. *J Am Assoc Gynecol Laparosc*. Feb 1995; 2(2): 175-9. PMID 9050553
25. Goldfarb HA. Nd:YAG laser laparoscopic coagulation of symptomatic myomas. *J Reprod Med*. Jul 1992; 37(7): 636-8. PMID 1387912
26. Nisolle M, Smets M, Malvaux V, et al. Laparoscopic myolysis with the Nd:YAG laser. *J Gynecol Surg*. 1993; 9(2): 95-9. PMID 10171973
27. Donnez J, Squifflet J, Polet R, et al. Laparoscopic myolysis. *Hum Reprod Update*. Nov-Dec 2000; 6(6): 609-13. PMID 11129695
28. Phillips DR, Nathanson HG, Milim SJ, et al. Laparoscopic Leiomyoma Coagulation. *J Am Assoc Gynecol Laparosc*. Aug 1996; 3(4, Supplement): S39. PMID 9074213
29. Zreik TG, Rutherford TJ, Palter SF, et al. Cryomyolysis, a new procedure for the conservative treatment of uterine fibroids. *J Am Assoc Gynecol Laparosc*. Feb 1998; 5(1): 33-8. PMID 9454874
30. Zupi E, Piredda A, Marconi D, et al. Directed laparoscopic cryomyolysis: a possible alternative to myomectomy and/or hysterectomy for symptomatic leiomyomas. *Am J Obstet Gynecol*. Mar 2004; 190(3): 639-43. PMID 15041993
31. Zupi E, Marconi D, Sbracia M, et al. Directed laparoscopic cryomyolysis for symptomatic leiomyomata: one-year follow up. *J Minim Invasive Gynecol*. Jul-Aug 2005; 12(4): 343-6. PMID 16036195
32. Hindley JT, Law PA, Hickey M, et al. Clinical outcomes following percutaneous magnetic resonance image guided laser ablation of symptomatic uterine fibroids. *Hum Reprod*. Oct 2002; 17(10): 2737-41. PMID 12351555
33. National Institute for Health and Care Excellence (NICE). Interventional procedures guidance: Transcervical ultrasound-guided radiofrequency ablation for symptomatic uterine fibroids [IPG689]. March 31, 2021; <https://www.nice.org.uk/guidance/ipg689>. Accessed on January 3, 2022.

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
September 2013	New policy	
September 2014	Replace policy	Policy updated with literature review. References 2, 4, and 15 added. Policy statement unchanged.
September 2015	Replace policy	Policy updated with literature review; references 5 and 15 added. Policy statement unchanged.
September 2016	Replace policy	Policy updated with literature review; references 3-4 added. Policy statement unchanged.
December 2017	Replace policy	Policy updated with literature review through 2017; references 7 and 18 added. Policy statement unchanged.
December 2018	Replace policy	Policy updated with literature review through June 4, 2018; reference 2 added. Policy statement unchanged.
December 2019	Replace policy	Policy updated with literature review through June 16, 2019; references added. Policy statement unchanged.
December 2020	Replace policy	Policy updated with literature review through June 26, 2020; references added. Policy statement unchanged.
March 2022	Replace policy	Policy updated with clinical input and literature review through January 3, 2022; references added. Title changed to "Laparoscopic, percutaneous, and transcervical techniques for uterine fibroid myolysis." Policy statements revised. Use of laparoscopic or transcervical radiofrequency ablation for the treatment of symptomatic uterine fibroids may be considered medically necessary in patients meeting criteria. Use of laser or bipolar needles, cryomyolysis, or MRI-guided laser ablation maintained as investigational. Policy previously tabled in 2021 to obtain clinical input.
June 2022	Administrative Review	No change in policy statements or references.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.