



FEP Medical Policy Manual

FEP 7.01.48 Autologous Chondrocyte Implantation for Focal Articular Cartilage Lesions

Effective Policy Date: July 1, 2022

Original Policy Date: December 2011

Related Policies:

7.01.15 - Meniscal Allografts and Other Meniscal Implants

8.01.52 - Orthopedic Applications of Stem Cell Therapy (Including Allografts and Bone Substitutes Used With Autologous Bone Marrow)

Autologous Chondrocyte Implantation for Focal Articular Cartilage Lesions

Description

Description

A variety of procedures are being developed to resurface articular cartilage defects. Autologous chondrocyte implantation involves harvesting chondrocytes from healthy tissue, expanding the cells in vitro, and implanting the expanded cells into the chondral defect. Second- and third-generation techniques include combinations of autologous chondrocytes, scaffolds, and growth factors.

OBJECTIVE

The objective of this evidence review is to determine whether autologous chondrocyte implantation improves net health outcomes in patients with focal articular cartilage lesions of the knee and other joints.

POLICY STATEMENT

Autologous chondrocyte implantation may be considered **medically necessary** for the treatment of disabling full-thickness articular cartilage defects of the knee caused by acute or repetitive trauma when all of the following criteria are met:

- Adolescent patients should be skeletally mature with documented closure of growth plates (eg, ≥ 15 years). Adult patients should be too young to be considered an appropriate candidate for total knee arthroplasty or other reconstructive knee surgery (eg, < 55 years)
- Focal, full-thickness (grade III or IV) unipolar lesions of the weight-bearing surface of the femoral condyles, trochlea, or patella at least 1.5 cm^2 in size
- Documented minimal to absent degenerative changes in the surrounding articular cartilage (Outerbridge grade II or less), and normal-appearing hyaline cartilage surrounding the border of the defect
- Normal knee biomechanics or alignment and stability achieved concurrently with autologous chondrocyte implantation.

Autologous chondrocyte implantation for all other joints, including the talar, and any indications other than those listed above is considered **investigational**.

POLICY GUIDELINES

For smaller lesions (eg, $< 4 \text{ cm}^2$), if debridement is the only prior surgical treatment, then consideration should be given to marrow-stimulating techniques before autologous chondrocyte implantation is performed.

The average defect size reported in the literature is about 5 cm^2 ; many studies treated lesions as large as 15 cm^2 .

Severe obesity (eg, body mass index $> 35 \text{ kg/m}^2$) may affect outcomes due to the increased stress on weight-bearing surfaces of the joint.

Misalignment and instability of the joint are contraindications. Therefore, additional procedures, such as repair of ligaments or tendons or creation of an osteotomy for realignment of the joint, may be performed at the same time. In addition, meniscal allograft transplantation may be performed in combination, either concurrently or sequentially, with autologous chondrocyte implantation. The charges for the culturing component of the procedure are submitted as part of the hospital bill.

The entire matrix-induced autologous chondrocyte implantation procedure consists of 4 steps: (1) initial arthroscopy and biopsy of normal cartilage, (2) culturing of chondrocytes on an absorbable collagen matrix, (3) a separate arthrotomy to place the implant, and (4) postsurgical rehabilitation. The initial arthroscopy may be scheduled as a diagnostic procedure; as part of this procedure, a cartilage defect may be identified, prompting biopsy of normal cartilage in anticipation of a possible chondrocyte transplant. The biopsied material is then sent for culturing and returned to the hospital when the implantation procedure (ie, arthrotomy) is scheduled.

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

FDA REGULATORY STATUS

The culturing of chondrocytes is considered by the FDA to fall into the category of manipulated autologous structural cells, which are subject to a biologic licensing requirement. In 1997, Carticel (Genzyme; now Vericel) received the FDA approval for the repair of clinically significant, "...symptomatic cartilaginous defects of the femoral condyle (medial-lateral or trochlear) caused by acute or repetitive trauma..."

In December 2016, MACI (Vericel) received FDA approval for "the repair of symptomatic, single or multiple full-thickness cartilage defects of the knee with or without bone involvement in adults." MACI consists of autologous chondrocytes that are cultured onto a bioresorbable porcine-derived collagen membrane. In 2017, production of Carticel was phased out, and MACI is the only autologous chondrocyte implantation product available in the United States.

A number of other second-generation methods for implanting autologous chondrocytes in a biodegradable matrix are currently in development or testing or are available outside of the United States. They include Atelocollagen (Koken), a collagen gel; Bioseed C (BioTissue Technologies), a polymer scaffold; CaReS (Ars Arthro), collagen gel; Cartilix (Biomet), a polymer hydrogel; Chondron (Sewon Cellontech), a fibrin gel; Hyalograft C (Fidia Advanced Polymers), a hyaluronic acid-based scaffold; NeoCart (Histogenics), an autologous chondrocyte implantation with a 3-dimensional chondromatrix in a phase 3 trial; and Novocart3D (Aesculap Biologics), a collagen-chondroitin sulfate scaffold in a phase 3 trial. ChondroCelect (TiGenix), characterized as a chondrocyte implantation with a completed phase 3 trial, uses a gene marker profile to determine in vivo cartilage-forming potential and thereby optimizes the phenotype (eg, hyaline cartilage vs. fibrocartilage) of the tissue produced with each autologous chondrocyte implantation cell batch. Each batch of chondrocytes is graded based on the quantitative gene expression of a selection of positive and negative markers for hyaline cartilage formation. Both Hyalograft C and ChondroCelect have been withdrawn from the market in Europe. In 2020, the FDA granted breakthrough status to Agili-C (CartiHeal, Ltd.), a proprietary biocompatible and biodegradable tapered-shape implant for the treatment of cartilage lesions in arthritic and non-arthritic joints that, when implanted into a pre-prepared osteochondral hole, acts as a 3-dimensional scaffold that potentially supports and promotes the regeneration of the articular cartilage and its underlying subchondral bone.

RATIONALE

Summary of Evidence

For individuals who have focal articular cartilage lesion(s) of the weight-bearing surface of the femoral condyles, trochlea, or patella who receive autologous chondrocyte implantation, the evidence includes systematic reviews, randomized controlled trials (RCTs), and prospective observational studies. Relevant outcomes are symptoms, change in disease status, morbid events, functional outcomes, and quality of life. There is a large body of evidence on autologous chondrocyte implantation for the treatment of focal articular cartilage lesions of the knee. For large lesions, autologous chondrocyte implantation results in better outcomes than microfracture, particularly in the long term. In addition, there is a limit to the size of lesions that can be treated with osteochondral autograft transfer, due to a limit on the number of osteochondral cores that can be safely harvested. As a result, autologous chondrocyte implantation has become the established treatment for large articular cartilage lesions in the knee. In 2017, first-generation autologous chondrocyte implantation with a collagen cover was phased out and replaced with an autologous chondrocyte implantation preparation that seeds the chondrocytes onto a bioresorbable collagen sponge. Although the implantation procedure for this second-generation autologous chondrocyte implantation is less technically demanding, studies to date have not shown improved outcomes compared with first-generation autologous chondrocyte implantation. Some evidence has suggested an increase in hypertrophy (overgrowth) of the new implant that may exceed that of the collagen membrane-covered implant. Long-term studies with a larger number of patients will be needed to determine whether this hypertrophy impacts graft survival. Based on mid-term outcomes that approximate those of first-generation autologous chondrocyte implantation and the lack of alternatives, second-generation autologous chondrocyte implantation may be considered an option for large disabling full-thickness cartilage lesions of the knee. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have focal articular cartilage lesions of joints other than the knee who receive autologous chondrocyte implantation, the evidence includes systematic reviews of case series. Relevant outcomes are symptoms, change in disease status, morbid events, functional outcomes, and quality of life. The greatest amount of literature is for autologous chondrocyte implantation of the talus. Comparative trials are needed to determine whether autologous chondrocyte implantation improves outcomes for lesions in joints other than the knee. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in "Supplemental Information" if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Academy of Orthopaedic Surgeons

In its 2010 guidelines on the diagnosis and treatment of osteochondritis dissecans, the American Academy of Orthopaedic Surgeons did not recommend for or against a specific cartilage repair technique in symptomatic skeletally immature or mature patients with an unsalvageable

osteochoondritis dissecans lesion.⁴³ This finding of insufficient evidence was based on a systematic review that found 4 level IV studies addressing cartilage repair techniques for an unsalvageable osteochoondritis dissecans lesion. Because each level IV article used different techniques, different outcome measures, and differing lengths of follow-up, the Academy deemed the evidence for any specific technique inconclusive.

National Institute for Health and Care Excellence

In 2018, NICE updated its 2005 guidance on the use of autologous chondrocyte implantation.⁴⁴ The NICE recommendations are stated below:

"...as an option for treating symptomatic articular cartilage defects of the femoral condyle and patella of the knee (International Cartilage Repair Society grade III or IV) in adults, only if:

- the person has not had previous surgery to repair articular cartilage defects;
- there is minimal osteoarthritic damage to the knee (as assessed by clinicians experienced in investigating knee cartilage damage using a validated measure for knee osteoarthritis); and
- the defect is over 2 cm²."

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

REFERENCES

1. Niemeyer P, Pestka JM, Kreuz PC, et al. Characteristic complications after autologous chondrocyte implantation for cartilage defects of the knee joint. *Am J Sports Med.* Nov 2008; 36(11): 2091-9. PMID 18801942
2. Free online Modified Cincinnati Knee Rating System calculator. OrthoToolKit. <https://www.orthotoolkit.com/cincinnati/>. Accessed February 16, 2022.
3. Greco NJ, Anderson AF, Mann BJ, et al. Responsiveness of the International Knee Documentation Committee Subjective Knee Form in comparison to the Western Ontario and McMaster Universities Osteoarthritis Index, modified Cincinnati Knee Rating System, and Short Form 36 in patients with focal articular cartilage defects. *Am J Sports Med.* May 2010; 38(5): 891-902. PMID 20044494
4. Gusi N, Olivares PR, Rajendram R. The EQ-5D Health-Related Quality of Life Questionnaire [Abstract]. In: Preedy VR, Watson RR, eds. *Handbook of Disease Burdens and Quality of Life Measures.* New York: Springer; 2010:87-89.
5. Roos EM, Engelhart L, Ranstam J, et al. ICRS Recommendation Document: Patient-Reported Outcome Instruments for Use in Patients with Articular Cartilage Defects. *Cartilage.* Apr 2011; 2(2): 122-36. PMID 26069575
6. Roos EM, Lohmander LS. The Knee injury and Osteoarthritis Outcome Score (KOOS): from joint injury to osteoarthritis. *Health Qual Life Outcomes.* Nov 03 2003; 1: 64. PMID 14613558
7. Collins NJ, Misra D, Felson DT, et al. Measures of knee function: International Knee Documentation Committee (IKDC) Subjective Knee Evaluation Form, Knee Injury and Osteoarthritis Outcome Score (KOOS), Knee Injury and Osteoarthritis Outcome Score Physical Function Short Form (KOOS-PS), Knee Outcome Survey Activities of Daily Living Scale (KOS-ADL), Lysholm Knee Scoring Scale, Oxford Knee Score (OKS), Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), Activity Rating Scale (ARS), and Tegner Activity Score (TAS). *Arthritis Care Res (Hoboken).* Nov 2011; 63 Suppl 11: S208-28. PMID 22588746
8. Lee WC, Kwan YH, Chong HC, et al. The minimal clinically important difference for Knee Society Clinical Rating System after total knee arthroplasty for primary osteoarthritis. *Knee Surg Sports Traumatol Arthrosc.* Nov 2017; 25(11): 3354-3359. PMID 27324635
9. Clement ND, MacDonald D, Simpson AH. The minimal clinically important difference in the Oxford knee score and Short Form 12 score after total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc.* Aug 2014; 22(8): 1933-9. PMID 24253376
10. Copay AG, Eyberg B, Chung AS, et al. Minimum Clinically Important Difference: Current Trends in the Orthopaedic Literature, Part II: Lower Extremity: A Systematic Review. *JBJS Rev.* Sep 2018; 6(9): e2. PMID 30179898
11. Bin Abd Razak HR, Acharyya S, Tan SM, et al. Predictors of Midterm Outcomes after Medial Unicompartmental Knee Arthroplasty in Asians. *Clin Orthop Surg.* Dec 2017; 9(4): 432-438. PMID 29201296
12. Lee WC, Bin Abd Razak HR, Allen JC, et al. Achieving Minimum Clinically Important Difference in Oxford Knee Score and Short Form-36 Physical Component Summary Is Less Likely with Single-Radius Compared with Multiradius Total Knee Arthroplasty in Asians. *J Knee Surg.*

- Mar 2019; 32(3): 227-232. PMID 29635649
13. Abraamyan T, Johnson AJ, Wiedrick J, et al. Marrow Stimulation Has Relatively Inferior Patient-Reported Outcomes in Cartilage Restoration Surgery of the Knee: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Am J Sports Med.* Apr 23 2021: 3635465211003595. PMID 33890799
 14. Zamborsky R, Danisovic L. Surgical Techniques for Knee Cartilage Repair: An Updated Large-Scale Systematic Review and Network Meta-analysis of Randomized Controlled Trials. *Arthroscopy.* Mar 2020; 36(3): 845-858. PMID 32139062
 15. Riboh JC, Cvetanovich GL, Cole BJ, et al. Comparative efficacy of cartilage repair procedures in the knee: a network meta-analysis. *Knee Surg Sports Traumatol Arthrosc.* Dec 2017; 25(12): 3786-3799. PMID 27605128
 16. Devitt BM, Bell SW, Webster KE, et al. Surgical treatments of cartilage defects of the knee: Systematic review of randomised controlled trials. *Knee.* Jun 2017; 24(3): 508-517. PMID 28189406
 17. Mundi R, Bedi A, Chow L, et al. Cartilage Restoration of the Knee: A Systematic Review and Meta-analysis of Level 1 Studies. *Am J Sports Med.* Jul 2016; 44(7): 1888-95. PMID 26138733
 18. Gou GH, Tseng FJ, Wang SH, et al. Autologous Chondrocyte Implantation Versus Microfracture in the Knee: A Meta-analysis and Systematic Review. *Arthroscopy.* Jan 2020; 36(1): 289-303. PMID 31708355
 19. Mistry H, Connock M, Pink J, et al. Autologous chondrocyte implantation in the knee: systematic review and economic evaluation. *Health Technol Assess.* Feb 2017; 21(6): 1-294. PMID 28244303
 20. Harris JD, Siston RA, Pan X, et al. Autologous chondrocyte implantation: a systematic review. *J Bone Joint Surg Am.* Sep 15 2010; 92(12): 2220-33. PMID 20844166
 21. Sacolick DA, Kirven JC, Abouljoud MM, et al. The Treatment of Adult Osteochondritis Dissecans with Autologous Cartilage Implantation: A Systematic Review. *J Knee Surg.* Nov 2019; 32(11): 1102-1110. PMID 30396204
 22. Bartlett W, Skinner JA, Gooding CR, et al. Autologous chondrocyte implantation versus matrix-induced autologous chondrocyte implantation for osteochondral defects of the knee: a prospective, randomised study. *J Bone Joint Surg Br.* May 2005; 87(5): 640-5. PMID 15855365
 23. Saris D, Price A, Widuchowski W, et al. Matrix-Applied Characterized Autologous Cultured Chondrocytes Versus Microfracture: Two-Year Follow-up of a Prospective Randomized Trial. *Am J Sports Med.* Jun 2014; 42(6): 1384-94. PMID 24714783
 24. Brittberg M, Recker D, Ilgenfritz J, et al. Matrix-Applied Characterized Autologous Cultured Chondrocytes Versus Microfracture: Five-Year Follow-up of a Prospective Randomized Trial. *Am J Sports Med.* May 2018; 46(6): 1343-1351. PMID 29565642
 25. Basad E, Ishaque B, Bachmann G, et al. Matrix-induced autologous chondrocyte implantation versus microfracture in the treatment of cartilage defects of the knee: a 2-year randomised study. *Knee Surg Sports Traumatol Arthrosc.* Apr 2010; 18(4): 519-27. PMID 20062969
 26. Basad E, Wissing FR, Fehrenbach P, et al. Matrix-induced autologous chondrocyte implantation (MACI) in the knee: clinical outcomes and challenges. *Knee Surg Sports Traumatol Arthrosc.* Dec 2015; 23(12): 3729-35. PMID 25218576
 27. Schuette HB, Kraeutler MJ, McCarty EC. Matrix-Assisted Autologous Chondrocyte Transplantation in the Knee: A Systematic Review of Mid- to Long-Term Clinical Outcomes. *Orthop J Sports Med.* Jun 2017; 5(6): 2325967117709250. PMID 28620621
 28. Meyerkort D, Ebert JR, Ackland TR, et al. Matrix-induced autologous chondrocyte implantation (MACI) for chondral defects in the patellofemoral joint. *Knee Surg Sports Traumatol Arthrosc.* Oct 2014; 22(10): 2522-30. PMID 24817164
 29. Zak L, Aldrian S, Wondrasch B, et al. Ability to return to sports 5 years after matrix-associated autologous chondrocyte transplantation in an average population of active patients. *Am J Sports Med.* Dec 2012; 40(12): 2815-21. PMID 23108635
 30. Ebert JR, Fallon M, Wood DJ, et al. A Prospective Clinical and Radiological Evaluation at 5 Years After Arthroscopic Matrix-Induced Autologous Chondrocyte Implantation. *Am J Sports Med.* Jan 2017; 45(1): 59-69. PMID 27587741
 31. Ebert JR, Fallon M, Zheng MH, et al. A randomized trial comparing accelerated and traditional approaches to postoperative weightbearing rehabilitation after matrix-induced autologous chondrocyte implantation: findings at 5 years. *Am J Sports Med.* Jul 2012; 40(7): 1527-37. PMID 22539536
 32. Ebert JR, Smith A, Edwards PK, et al. Factors predictive of outcome 5 years after matrix-induced autologous chondrocyte implantation in the tibiofemoral joint. *Am J Sports Med.* Jun 2013; 41(6): 1245-54. PMID 23618699
 33. Ebert JR, Schneider A, Fallon M, et al. A Comparison of 2-Year Outcomes in Patients Undergoing Tibiofemoral or Patellofemoral Matrix-Induced Autologous Chondrocyte Implantation. *Am J Sports Med.* Dec 2017; 45(14): 3243-3253. PMID 28910133
 34. Harris JD, Cavo M, Brophy R, et al. Biological knee reconstruction: a systematic review of combined meniscal allograft transplantation and cartilage repair or restoration. *Arthroscopy.* Mar 2011; 27(3): 409-18. PMID 21030203
 35. Andriolo L, Merli G, Filardo G, et al. Failure of Autologous Chondrocyte Implantation. *Sports Med Arthrosc Rev.* Mar 2017; 25(1): 10-18. PMID 28045868
 36. Nawaz SZ, Bentley G, Briggs TW, et al. Autologous chondrocyte implantation in the knee: mid-term to long-term results. *J Bone Joint Surg Am.* May 21 2014; 96(10): 824-30. PMID 24875023
 37. Minas T, Von Keudell A, Bryant T, et al. The John Insall Award: A minimum 10-year outcome study of autologous chondrocyte implantation. *Clin Orthop Relat Res.* Jan 2014; 472(1): 41-51. PMID 23979923
 38. Minas T, Gomoll AH, Rosenberger R, et al. Increased failure rate of autologous chondrocyte implantation after previous treatment with marrow stimulation techniques. *Am J Sports Med.* May 2009; 37(5): 902-8. PMID 19261905
 39. Ebert JR, Smith A, Fallon M, et al. Incidence, degree, and development of graft hypertrophy 24 months after matrix-induced autologous chondrocyte implantation: association with clinical outcomes. *Am J Sports Med.* Sep 2015; 43(9): 2208-15. PMID 26163536
 40. Niemeyer P, Salzmann G, Schmal H, et al. Autologous chondrocyte implantation for the treatment of chondral and osteochondral defects of the talus: a meta-analysis of available evidence. *Knee Surg Sports Traumatol Arthrosc.* Sep 2012; 20(9): 1696-703. PMID 22037894
 41. Hu M, Li X, Xu X. Efficacy and safety of autologous chondrocyte implantation for osteochondral defects of the talus: a systematic review and meta-analysis. *Arch Orthop Trauma Surg.* Jun 14 2021. PMID 34128117

42. Shimozono Y, Yasui Y, Ross AW, et al. Scaffolds based therapy for osteochondral lesions of the talus: A systematic review. *World J Orthop.* Oct 18 2017; 8(10): 798-808. PMID 29094011
43. American Academy of Orthopaedic Surgeons. *Clinical Practice Guideline on the Diagnosis and Treatment of Osteochondritis Dissecans.* Rosemont, IL: AAOS; 2010.
44. National Institute for Health and Care Excellence (NICE). *Autologous chondrocyte implantation for treating symptomatic articular cartilage defects of the knee [TA508]*. 2018; <https://www.nice.org.uk/guidance/TA508/chapter/1-Recommendations>. Accessed February 16, 2022.

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
December 2011	New policy	
September 2013	Replace policy	Policy updated with literature review, references 12 and 39-42 added; sections and statements on minced cartilage moved to policy No.7.01.78. Policy title change (Osteochondral Autografts and Allografts and Other Cell-based Treatments removed from title.
September 2015	Replace policy	Policy updated with literature review; references 5 and 7 added; policy statements unchanged.
June 2017		Clinical input reviewed; references 8 and 32-33 added. Autologous chondrocyte implantation of the patella considered medically necessary; need for a prior surgical procedure removed from policy statement. Policy updated with literature review through March 2, 2017; references 5, 7, 10, 12, and 19 added. Rationale extensively revised to focus on available products. Investigational statement on matrix-induced autologous chondrocyte implantation removed.
March 2018	Replace policy	Policy updated with literature review through November 13, 2017, focusing on matrix-induced autologous chondrocyte implantation of the patella; references 12-18 added. Matrix-induced autologous chondrocyte implantation of the patella is considered medically necessary. In the investigational statement the wording: "and any indications other than those listed above" changed to "and any non-FDA approved indications" for clarification.
June 2018	Replace policy	Policy updated with literature review through February 5, 2018; references 6, 8, 22, 27, and 30 added. Policy statements unchanged.
April 2019	Replace policy	Policy updated with literature review through February 5, 2019, no references added; reference 30 updated. Policy statements unchanged.
June 2020	Replace policy	Policy updated with literature review through February 11, 2020; references added. Policy statements unchanged.
June 2021	Replace policy	Policy updated with literature review through February 23, 2021; references added. Policy statements unchanged.
June 2022	Replace policy	Policy updated with literature review through February 16, 2022; references added. Policy statements unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.