



FEP Medical Policy Manual

FEP 8.01.02 Chelation Therapy for Off-Label Uses

Effective Policy Date: July 1, 2022

Original Policy Date: December 2011

Related Policies:

None

Chelation Therapy for Off-Label Uses

Description

Description

Chelation therapy, an established treatment for heavy metal toxicities and transfusional hemosiderosis, has been investigated for a variety of off-label applications, such as treatment of atherosclerosis, Alzheimer disease, and autism. This evidence review does not address indications for chelation therapy approved by the U.S. Food and Drug Administration. Instead, it addresses off-label indications, including Alzheimer disease, cardiovascular disease, autism spectrum disorder, diabetes, multiple sclerosis, and arthritis.

OBJECTIVE

The objective of this evidence review is to determine whether chelation therapy, when used as a treatment for various off-label applications such as Alzheimer disease, cardiovascular disease, autism spectrum disorder, diabetes, multiple sclerosis, and arthritis, improves the net health outcome.

POLICY STATEMENT

Off-label applications of chelation therapy (see Policy Guidelines section for uses approved by the U.S. Food and Drug Administration) are considered **investigational**, including, but not limited to:

- Alzheimer disease
- atherosclerosis (eg, coronary artery disease, secondary prevention in patients with myocardial infarction, or peripheral vascular disease)
- autism
- diabetes
- multiple sclerosis
- arthritis (includes rheumatoid arthritis).

POLICY GUIDELINES

A number of indications for chelation therapy have received U.S. Food and Drug Administration (FDA) approval and for which chelation therapy is considered standard of care. These indications include:

- extreme conditions of metal toxicity
- treatment of chronic iron overload due to blood transfusions (transfusional hemosiderosis) or due to non-transfusion-dependent thalassemia
- Wilson disease (hepatolenticular degeneration)
- lead poisoning
- control of ventricular arrhythmias or heart block associated with digitalis toxicity
- emergency treatment of hypercalcemia.

For the last 2 bullet points, most patients should be treated with other modalities. Digitalis toxicity is currently treated in most patients with Fab monoclonal antibodies. The FDA removed the approval for NaEDTA as chelation therapy due to safety concerns and recommended that other chelators be used. NaEDTA was the most common chelation agent used to treat digitalis toxicity and hypercalcemia.

Suggested toxic or normal levels of select heavy metals are listed in Table 1.

Table 1. Toxic or Normal Concentrations of Heavy Metals

Metal	Toxic Levels (Normal Levels Where Indicated)
Arsenic	24-h urine: ≥ 50 $\mu\text{g/L}$ urine or 100 $\mu\text{g/g}$ creatinine
Bismuth	No clear reference standard
Cadmium	Proteinuria and/or ≥ 15 $\mu\text{g/g}$ creatinine
Chromium	No clear reference standard
Cobalt	Normative excretion: 0.1-1.2 $\mu\text{g/L}$ (serum), 0.1-2.2 $\mu\text{g/L}$ (urine)
Copper	Normative excretion: 25 $\mu\text{g}/24$ h (urine)

Metal	Toxic Levels (Normal Levels Where Indicated)
Iron	<ul style="list-style-type: none"> • Nontoxic: <300 µg/dL • Severe: >500 µg/dL
Lead	<p>Pediatric</p> <ul style="list-style-type: none"> • Symptoms or blood lead level ≥45 µg/dL (blood) • CDC level of concern: 5 µg/dL³³, <p>Adult</p> <ul style="list-style-type: none"> • Symptoms or blood lead level ≥70 µg/dL • CDC level of concern: 10 µg/dL³⁴,
Manganese	No clear reference standard
Mercury	Background exposure normative limits: 1-8 µg/L (whole blood); 4-5 µg/L (urine) ^{35,,a}
Nickel	<ul style="list-style-type: none"> • Excessive exposure: ≥8 µg/L (blood) • Severe poisoning: ≥500 µg/L (8-h urine)
Selenium	<ul style="list-style-type: none"> • Mild toxicity: >1 mg/L (serum) • Serious toxicity: >2 mg/L
Silver	Asymptomatic workers have mean levels of 11 µg/L (serum) and 2.6 µg/L (spot urine)
Thallium	24-hour urine thallium >5 µg/L ³⁶ ,
Zinc	Normative range: 0.6-1.1 mg/L (plasma), 10-14 mg/L (red cells)

Adapted from Adal (2018).³⁷.

CDC: Centers for Disease Control and Prevention.

^a Hair analysis is useful to assess mercury exposure in epidemiologic studies. However, hair analysis in individual patients must be interpreted with consideration of the patient's history, signs, and symptoms, and possible alternative explanations.

Measurement of blood and urine mercury levels can exclude exogenous contamination; therefore, blood or urine mercury levels may be more robust measures of exposure in individual patients. [Kempson IM, Lombi E. Hair analysis as a biomonitor.... 40\(7\): 3915-40. PMID 21468435\]](#)

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

FDA REGULATORY STATUS

In 1953, EDTA (Versenate) was approved by the FDA for lowering blood lead levels among both pediatric and adult patients with lead poisoning. In 1991, succimer (Chemet) was approved by the FDA for the treatment of lead poisoning in pediatric patients only. The FDA approved disodium-EDTA for use in selected patients with hypercalcemia and use in patients with heart rhythm problems due to intoxication with digitalis. In 2008, the FDA withdrew approval of disodium-EDTA due to safety concerns and recommended that other forms of chelation therapy be used.²

Several iron-chelating agents are FDA approved:

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

- In 1968, deferoxamine (Desferal; Novartis) was approved by the FDA for subcutaneous, intramuscular, or intravenous injections to treat acute iron intoxication and chronic iron overload due to transfusion-dependent anemia. Several generic forms of deferoxamine have been approved by the FDA.
- In 2005, deferasirox (Exjade; Novartis) was approved by the FDA, is available as a tablet for oral suspension, and is indicated for the treatment of chronic iron overload due to blood transfusions in patients ages 2 years and older. Under the accelerated approval program, the FDA expanded the indications for deferasirox in 2013 to include treatment of patients aged 10 years and older with chronic iron overload due to non-transfusion-dependent thalassemia syndromes and specific liver iron concentration and serum ferritin levels. A generic version of deferasirox tablet for oral suspension has also been approved by the FDA. In 2015, an oral tablet formulation for deferasirox (Jadenu) was approved by the FDA. All formulations of deferasirox carry a black box warning because it may cause serious and fatal renal toxicity and failure, hepatic toxicity and failure, and gastrointestinal hemorrhage. As a result, treatment with deferasirox requires close patient monitoring, including laboratory tests of renal and hepatic function.
- In 2011, the iron chelator deferiprone (Ferriprox) was approved by the FDA for treatment of patients with transfusional overload due to thalassemia syndromes when another chelation therapy is inadequate. Deferiprone is available in tablet and oral solution. Ferriprox carries a black box warning because it can cause agranulocytosis, which can lead to serious infections and death. As a result, absolute neutrophil count should be monitored before and during treatment.

In a June 2014 warning to consumers, the FDA advised that FDA-approved chelating agents would be available by prescription only. There are no FDA approved over-the-counter chelation products.

RATIONALE

Summary of Evidence

For individuals who have Alzheimer disease, or cardiovascular disease, or autism spectrum disorder, or diabetes, or multiple sclerosis, or arthritis who receive chelation therapy, the evidence includes a small number of randomized controlled trials (RCTs) and case series. Relevant outcomes are symptoms, change in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related morbidity. One RCT (the Trial to Assess Chelation Therapy) reported that chelation therapy reduced cardiovascular events in patients with previous myocardial infarction and that the benefit was greater in diabetic patients compared with nondiabetic patients. However, this trial had significant limitations (eg, high dropout rates) and, therefore, conclusions are not definitive. For other conditions, the available RCTs did not report improvements in health outcomes with chelation therapy and, as evidence, the case series are inadequate to determine efficacy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

The purpose of the remaining sections in Supplemental Information is to provide reference material regarding existing practice guidelines and position statements, U.S. Preventive Services Task Force Recommendations and Medicare National Coverage Decisions and registered, ongoing clinical trials. Inclusion in the Supplemental Information does not imply endorsement and information may not necessarily be used in formulating the evidence review conclusions.

Guidelines or position statements will be considered for inclusion in Supplemental Information if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Heart Association and American College of Cardiology

In 2016, the American College of Cardiology (ACC) and the American Heart Association (AHA) published a joint guideline on the management of patients with lower extremity peripheral artery disease, which recommended that chelation therapy (eg, ethylenediaminetetraacetic acid) is not

beneficial for the treatment of claudication.²⁷

In 2014, the ACC and AHA published a focused update of the guideline for the management of stable ischemic heart disease, in conjunction with the American Association for Thoracic Surgery, Preventative Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons. This update included a revised recommendation on chelation therapy stating that the "usefulness of chelation therapy is uncertain for reducing cardiovascular events in patients with stable IHD."²⁸ Compared to the original publication of this guideline in 2012, the recommendation was upgraded from a class III (no benefit) to class IIb (benefit \geq risk), and the level of evidence from C (only consensus expert opinion, case studies, or standard of care) to B (data from a single randomized trial or nonrandomized studies).²⁹

American Academy of Pediatrics

In 2019, the American Academy of Pediatrics published guidance for the management of children with autism spectrum disorder. The guidance cautioned against the use of chelation therapy due to safety concerns and lack of supporting efficacy data.³⁰

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

The Centers for Medicare & Medicaid have issued 2 national coverage determinations on chelation therapy relevant to this evidence review. Section 20.21 states³¹:

"The application of chelation therapy using ethylenediamine-tetra-acetic acid (EDTA) for the treatment and prevention of atherosclerosis is controversial. There is no widely accepted rationale to explain the beneficial effects attributed to this therapy. Its safety is questioned and its clinical effectiveness has never been established by well designed, controlled clinical trials. It is not widely accepted and practiced by American physicians. EDTA chelation therapy for atherosclerosis is considered experimental. For these reasons, EDTA chelation therapy for the treatment or prevention of atherosclerosis is not covered.

Some practitioners refer to this therapy as chemoendarterectomy and may also show a diagnosis other than atherosclerosis, such as arteriosclerosis or calcinosis. Claims employing such variant terms should also be denied under this section."

Section 20.22 states³²:

"The use of EDTA as a chelating agent to treat atherosclerosis, arteriosclerosis, calcinosis, or similar generalized condition not listed by the FDA [U.S. Food and Drug Administration] as an approved use is not covered. Any such use of EDTA is considered experimental."

These national coverage determinations are long-standing; effective dates of these versions have not been posted.

REFERENCES

- Centers for Disease Control and Prevention (CDC). Deaths associated with hypocalcemia from chelation therapy--Texas, Pennsylvania, and Oregon, 2003-2005. *MMWR Morb Mortal Wkly Rep.* Mar 03 2006; 55(8): 204-7. PMID 16511441
- Food and Drug Administration. Hospira, Inc., et al.; Withdrawal of Approval of One New Drug Application and Two Abbreviated New Drug Application. *Federal Register.* 2008;73(113):33440-33441.
- Sampson E, Jenagaratnam L, McShane R. Metal protein attenuating compounds for the treatment of Alzheimer's disease. *Cochrane Database Syst Rev.* Jan 23 2008; (1): CD005380. PMID 18254079
- Ritchie CW, Bush AI, Mackinnon A, et al. Metal-protein attenuation with iodochlorhydroxyquin (clioquinol) targeting Abeta amyloid deposition and toxicity in Alzheimer disease: a pilot phase 2 clinical trial. *Arch Neurol.* Dec 2003; 60(12): 1685-91. PMID 14676042
- Sampson EL, Jenagaratnam L, McShane R. Metal protein attenuating compounds for the treatment of Alzheimer's dementia. *Cochrane Database Syst Rev.* May 16 2012; (5): CD005380. PMID 22592705
- Lannfelt L, Blennow K, Zetterberg H, et al. Safety, efficacy, and biomarker findings of PBT2 in targeting Abeta as a modifying therapy for Alzheimer's disease: a phase IIa, double-blind, randomised, placebo-controlled trial. *Lancet Neurol.* Sep 2008; 7(9): 779-86. PMID 18672400
- Villarruz-Sulit MV, Forster R, Dans AL, et al. Chelation therapy for atherosclerotic cardiovascular disease. *Cochrane Database Syst Rev.* May 05 2020; 5: CD002785. PMID 32367513
- Lamas GA, Goertz C, Boineau R, et al. Effect of disodium EDTA chelation regimen on cardiovascular events in patients with previous myocardial infarction: the TACT randomized trial. *JAMA.* Mar 27 2013; 309(12): 1241-50. PMID 23532240

9. Mark DB, Anstrom KJ, Clapp-Channing NE, et al. Quality-of-life outcomes with a disodium EDTA chelation regimen for coronary disease: results from the trial to assess chelation therapy randomized trial. *Circ Cardiovasc Qual Outcomes*. Jul 2014; 7(4): 508-16. PMID 24987051
10. Lamas GA, Boineau R, Goertz C, et al. EDTA chelation therapy alone and in combination with oral high-dose multivitamins and minerals for coronary disease: The factorial group results of the Trial to Assess Chelation Therapy. *Am Heart J*. Jul 2014; 168(1): 37-44.e5. PMID 24952858
11. Lewis EF, Ujueta F, Lamas GA, et al. Differential Outcomes With Edetate Disodium-Based Treatment Among Stable Post Anterior vs. Non-Anterior Myocardial Infarction Patients. *Cardiovasc Revasc Med*. Nov 2020; 21(11): 1389-1395. PMID 32303436
12. Nissen SE. Concerns about reliability in the Trial to Assess Chelation Therapy (TACT). *JAMA*. Mar 27 2013; 309(12): 1293-4. PMID 23532246
13. Maron DJ, Hlatky MA. Trial to Assess Chelation Therapy (TACT) and equipoise: When evidence conflicts with beliefs. *Am Heart J*. Jul 2014; 168(1): 4-5. PMID 24952853
14. Bernard S, Enayati A, Redwood L, et al. Autism: a novel form of mercury poisoning. *Med Hypotheses*. Apr 2001; 56(4): 462-71. PMID 11339848
15. Nelson KB, Bauman ML. Thimerosal and autism?. *Pediatrics*. Mar 2003; 111(3): 674-9. PMID 12612255
16. Ng DK, Chan CH, Soo MT, et al. Low-level chronic mercury exposure in children and adolescents: meta-analysis. *Pediatr Int*. Feb 2007; 49(1): 80-7. PMID 17250511
17. Rossignol DA. Novel and emerging treatments for autism spectrum disorders: a systematic review. *Ann Clin Psychiatry*. Oct-Dec 2009; 21(4): 213-36. PMID 19917212
18. Cooper GJ, Young AA, Gamble GD, et al. A copper(II)-selective chelator ameliorates left-ventricular hypertrophy in type 2 diabetic patients: a randomised placebo-controlled study. *Diabetologia*. Apr 2009; 52(4): 715-22. PMID 19172243
19. Escolar E, Lamas GA, Mark DB, et al. The effect of an EDTA-based chelation regimen on patients with diabetes mellitus and prior myocardial infarction in the Trial to Assess Chelation Therapy (TACT). *Circ Cardiovasc Qual Outcomes*. Jan 2014; 7(1): 15-24. PMID 24254885
20. Ujueta F, Arenas IA, Escolar E, et al. The effect of EDTA-based chelation on patients with diabetes and peripheral artery disease in the Trial to Assess Chelation Therapy (TACT). *J Diabetes Complications*. Jul 2019; 33(7): 490-494. PMID 31101487
21. Escolar E, Ujueta F, Kim H, et al. Possible differential benefits of edetate disodium in post-myocardial infarction patients with diabetes treated with different hypoglycemic strategies in the Trial to Assess Chelation Therapy (TACT). *J Diabetes Complications*. Aug 2020; 34(8): 107616. PMID 32446881
22. Chen KH, Lin JL, Lin-Tan DT, et al. Effect of chelation therapy on progressive diabetic nephropathy in patients with type 2 diabetes and high-normal body lead burdens. *Am J Kidney Dis*. Oct 2012; 60(4): 530-8. PMID 22721929
23. U.S. Department of Labor, Occupational Health and Safety Administration. Safety and Health Regulations for Construction: Substance Data Sheet for Occupational Exposure to Lead. 1993; http://www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=STANDARDS&p_id=10642. Accessed January 31, 2022.
24. Weinreb O, Mandel S, Youdim MBH, et al. Targeting dysregulation of brain iron homeostasis in Parkinson's disease by iron chelators. *Free Radic Biol Med*. Sep 2013; 62: 52-64. PMID 23376471
25. Grolez G, Moreau C, Sablonniere B, et al. Ceruloplasmin activity and iron chelation treatment of patients with Parkinson's disease. *BMC Neurol*. May 06 2015; 15: 74. PMID 25943368
26. van Eijk LT, Heemskerk S, van der Pluijm RW, et al. The effect of iron loading and iron chelation on the innate immune response and subclinical organ injury during human endotoxemia: a randomized trial. *Haematologica*. Mar 2014; 99(3): 579-87. PMID 24241495
27. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the Management of Patients With Lower Extremity Peripheral Artery Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. Mar 21 2017; 135(12): e726-e779. PMID 27840333
28. Fihn SD, Blankenship JC, Alexander KP, et al. 2014 ACC/AHA/AATS/PCNA/SCAI/STS focused update of the guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines, and the American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. Nov 04 2014; 64(18): 1929-49. PMID 25077860
29. Qaseem A, Fihn SD, Dallas P, et al. Management of stable ischemic heart disease: summary of a clinical practice guideline from the American College of Physicians/American College of Cardiology Foundation/American Heart Association/American Association for Thoracic Surgery/Preventive Cardiovascular Nurses Association/Society of Thoracic Surgeons. *Ann Intern Med*. Nov 20 2012; 157(10): 735-43. PMID 23165665
30. Hyman SL, Levy SE, Myers SM, et al. Identification, Evaluation, and Management of Children With Autism Spectrum Disorder. *Pediatrics*. Jan 2020; 145(1). PMID 31843864
31. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination (NCD) for CHELATION THERAPY for Treatment of Atherosclerosis (20.21). n.d.; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=86>. Accessed January 29, 2022.
32. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination (NCD) for Ethylenediamine- Tetra-Acetic (EDTA) CHELATION THERAPY for Treatment of Atherosclerosis (20.22). n.d.; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=146&ncdver=1&bc=AAAAQAAAAAA&>. Accessed January 28, 2022.
33. Centers for Disease Control and Prevention (CDC). What Do Parents Need to Know to Protect Their Children? 2017, May 17; http://www.cdc.gov/nceh/lead/ACCLPP/blood_lead_levels.htm. Accessed January 30, 2022.
34. Centers for Disease Control and Prevention (CDC). Very high blood lead levels among adults - United States, 2002-2011. *MMWR Morb Mortal Wkly Rep*. Nov 29 2013; 62(47): 967-71. PMID 24280917
35. Agency for Toxic Substances and Disease Registry. Toxicological profile for mercury. 1999 March; <https://www.atsdr.cdc.gov/ToxProfiles/tp46.pdf>. Accessed January 31, 2022.

36. Centers for Disease Control and Prevention (CDC). Emergency preparedness and response. Case definition: thallium. 2015 November 18; <https://emergency.cdc.gov/agent/thallium/casedef.asp>. Accessed January 31, 2022.
37. Adal A. Medscape. Heavy metal toxicity. 2018; <http://emedicine.medscape.com/article/814960-overview>. Accessed January 31, 2022.
38. Kempson IM, Lombi E. Hair analysis as a biomonitor for toxicology, disease and health status. *Chem Soc Rev*. Jul 2011; 40(7): 3915-40. PMID 21468435

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
December 2011	New policy	
September 2013	Replace policy	Policy updated with literature review. References 16-21 added, others removed or renumbered. Chronic iron overload due to nontransfusion-dependent thalassemia (NDTD) added to medically necessary statement based on new FDA approval. Secondary prevention in patients with myocardial infarction added to bullet point in investigational statement on atherosclerosis; in that bullet point, "i.e.€" changed to "e.g.€"
September 2014	Replace policy	Policy updated with literature review through May 21, 2014; references 14, 22-24, and 28-29 added; references 2, 19, and 25 updated. Title changed to "Chelation Therapy for Off-Label Uses.€" Medically necessary policy statement for on-label uses deleted from policy statement and moved to policy guidelines. Investigational policy statement unchanged.
September 2015	Replace policy	Policy updated with literature review through May 21, 2015; references 3, 4, 23-25, 27, 33, 35, 36, 38 and 41 added. Hypoglycemia deleted from policy statement; this indication is not reviewed in the policy. Policy statements otherwise unchanged.
June 2018	Replace policy	Policy updated with literature review through December 11, 2017; reference 8 removed; reference 38 and 44 updated; reference 39 added. Policy statement unchanged.
June 2019	Replace policy	Policy updated with literature review through January 3, 2019; no reference added. Policy statement unchanged.
June 2020	Replace policy	Policy updated with literature review through December 9, 2019; no references added. Policy statement unchanged.
June 2021	Replace policy	Policy updated with literature review through December 8, 2020; references added. Policy statement unchanged.
June 2022	Replace policy	Policy updated with literature review through November 15, 2021; no references added. Policy statements unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.