

## 5.40.33

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Cardiovascular Agent	<b>Original Policy Date:</b>	May 27, 2022
<b>Subject:</b>	Camzyos	<b>Page:</b>	1 of 5

---

**Last Review Date:** June 16, 2022

## Camzyos

### Description

#### Camzyos (mavacamten)

#### Background

Camzyos (mavacamten) is an allosteric and reversible inhibitor selective for cardiac myosin. Camzyos modulates the number of myosin heads that can enter “on actin” (power-generating) states, thus reducing the probability of force-producing (systolic) and residual (diastolic) cross-bridge formation. Excess myosin actin cross-bridge formation and dysregulation of the super-relaxed state are mechanistic hallmarks of hypertrophic cardiomyopathy (HCM). Camzyos shifts the overall myosin population towards an energy-sparing, recruitable, super-relaxed state. In HCM patients, myosin inhibition with Camzyos reduces dynamic left ventricular outflow tract (LVOT) obstruction and improves cardiac filling pressures (1).

#### Regulatory Status

FDA-approved indication: Camzyos is a cardiac myosin inhibitor indicated for the treatment of adults with symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) to improve functional capacity and symptoms (1).

Camzyos has a boxed warning regarding risk of heart failure. Camzyos reduces left ventricular ejection fraction (LVEF) and can cause heart failure due to systolic dysfunction.

Echocardiogram assessments of LVEF are required prior to and during treatment with Camzyos. Initiation of Camzyos in patients with LVEF <55% is not recommended. Concomitant use of Camzyos with certain cytochrome P450 inhibitors or discontinuation of certain cytochrome P450 inducers may increase the risk of heart failure due to systolic dysfunction; therefore, the use of Camzyos is contraindicated with the following:

- Moderate to strong CYP2C19 inhibitors or strong CYP3A4 inhibitors

# 5.40.33

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Cardiovascular Agent	<b>Original Policy Date:</b>	May 27, 2022
<b>Subject:</b>	Camzyos	<b>Page:</b>	2 of 5

---

- Moderate to strong CYP2C19 inducers or moderate to strong CYP3A4 inducers

Because of the risk of heart failure due to systolic dysfunction, Camzyos is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called Camzyos REMS Program (1).

Regular LVEF and Valsalva left ventricular outflow tract (LVOT) gradient assessment is required for careful titration to achieve an appropriate target Valsalva LVOT gradient, while maintaining LVEF  $\geq 50\%$  and avoiding heart failure symptoms (1).

Camzyos also contains a warning for embryo-fetal toxicity. Camzyos may cause fetal toxicity when administered to a pregnant female. Confirm absence of pregnancy in females of reproductive potential prior to treatment and advise patients to use effective contraception during treatment with Camzyos and for 4 months after the last dose. Camzyos may reduce the effectiveness of combine hormonal contraceptives (CHCs). Advise patients using CHCs to use an alternative method that is not affected by CYP450 enzyme induction or to add nonhormonal contraception (1).

The safety and effectiveness of Camzyos in pediatric patients less than 18 year of age have not been established (1).

## Related policies

### Policy

*This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.*

Camzyos may be considered **medically necessary** in patients 18 years of age or older with obstructive hypertrophic cardiomyopathy (HCM) and if the conditions indicated below are met.

Camzyos may be considered **investigational** in patients less than 18 years of age and for all other indications.

## Prior-Approval Requirements

**Age** 18 years of age or older

### Diagnosis

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Cardiovascular Agent	<b>Original Policy Date:</b>	May 27, 2022
<b>Subject:</b>	Camzyos	<b>Page:</b>	3 of 5

---

Patient must have the following:

Obstructive hypertrophic cardiomyopathy (HCM)

**AND ALL** of the following:

1. NYHA activity class II – III
2. Prescribed by or recommended by a cardiologist
3. Left ventricular ejection fraction (LVEF)  $\geq$  55%
4. Prescriber agrees to monitor echocardiogram, LVEF, and Valsalva left ventricular outflow tract (LVOT) gradient during treatment with Camzyos
5. Prescriber agrees to monitor for and counsel patient regarding CYP450 drug interactions with Camzyos
6. Patient and prescriber are enrolled in the Camzyos REMS Program
7. Females of reproductive potential **only**: absence of pregnancy has been confirmed and patient will be advised to use effective contraception during treatment with Camzyos and for 4 months after the last dose

## **Prior – Approval *Renewal* Requirements**

**Age** 18 years of age or older

### **Diagnosis**

Patient must have the following:

Obstructive hypertrophic cardiomyopathy (HCM)

**AND ALL** of the following:

1. Symptoms have improved or stabilized
2. Prescriber agrees to monitor echocardiogram, LVEF, and Valsalva left ventricular outflow tract (LVOT) gradient during treatment with Camzyos
3. Prescriber agrees to interrupt treatment with Camzyos if LVEF  $<$ 50%
4. Prescriber agrees to monitor for and counsel patient regarding CYP450 drug interactions with Camzyos
5. Patient and prescriber are enrolled in the Camzyos REMS Program
6. Females of reproductive potential **only**: patient will be advised to use effective contraception during treatment with Camzyos and for 4 months after the last dose

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Cardiovascular Agent	<b>Original Policy Date:</b>	May 27, 2022
<b>Subject:</b>	Camzyos	<b>Page:</b>	4 of 5

---

### Policy Guidelines

#### Pre - PA Allowance

None

#### Prior - Approval Limits

**Quantity** 90 capsules per 90 days

**Duration** 12 months

#### Prior – Approval *Renewal* Limits

Same as above

### Rationale

#### Summary

Camzyos (mavacamten) is an allosteric and reversible inhibitor selective for cardiac myosin and is indicated for the treatment of adults with symptomatic NYHA class II-III obstructive hypertrophic cardiomyopathy (HCM). Camzyos contains a boxed warning regarding the risk of heart failure. Camzyos is only available through the Camzyos REMS Program. The safety and effectiveness of Camzyos in pediatric patients less than 18 year of age have not been established (1).

Prior approval is required to ensure the safe, clinically appropriate and cost-effective use of Camzyos while maintaining optimal therapeutic outcomes.

#### References

1. Camzyos [package insert]. Brisbane, CA: Bristol Myers Squibb; May 2022.

### Policy History

Date	Action
May 2022	Addition to PA
June 2022	Annual review and reference update

### Keywords

---

# 5.40.33

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Cardiovascular Agent	<b>Original Policy Date:</b>	May 27, 2022
<b>Subject:</b>	Camzyos	<b>Page:</b>	5 of 5

---

**This policy was approved by the FEP® Pharmacy and Medical Policy Committee on June 16, 2022 and is effective on July 1, 2022.**