

# 5.21.155

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	August 27, 2020
<b>Subject:</b>	Tecartus	<b>Page:</b>	1 of 6

---

**Last Review Date:** June 16, 2022

---

## Tecartus

### Description

#### Tecartus (brexucabtagene autoleucel)

#### Background

Tecartus (brexucabtagene autoleucel) is a genetically modified autologous T cell immunotherapy indicated for the treatment of relapsed or refractory mantle cell lymphoma (MCL) and relapsed or refractory B-cell precursor acute lymphoblastic leukemia (ALL). The patient's own T-cells are collected and sent to a manufacturing center where they are genetically modified to include a new gene that contains a specific protein (a chimeric antigen receptor or CAR) that directs the T-cells to target and kill cancer cells that have a specific antigen (CD19) on the surface. Once the cells are modified, they are infused back into the patient to kill the cancer cells (1).

#### Regulatory Status

FDA-approved indications: Tecartus is a CD19-directed genetically modified autologous T cell immunotherapy indicated for the treatment of: (1)

- Adult patients with relapsed or refractory mantle cell lymphoma (MCL)
- Adult patients with relapsed or refractory B-cell precursor acute lymphoblastic leukemia (ALL)

Tecartus has a boxed warning for cytokine release syndrome (CRS) and neurological toxicities. Patients with an active infection or inflammatory disorders should not receive Tecartus and monitoring for neurological events should be done after treatment of Tecartus (1).

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	August 27, 2020
<b>Subject:</b>	Tecartus	<b>Page:</b>	2 of 6

---

Tecartus is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Healthcare facilities that dispense and administer Tecartus must be enrolled and comply with the REMS requirements. Certified healthcare facilities must have on-site, immediate access to tocilizumab, and ensure that a minimum of two doses of tocilizumab are available for each patient for administration within 2 hours after Tecartus infusion, if needed for treatment of CRS (1).

Serious infections, including life-threatening or fatal infections, occurred in patients after Tecartus infusion. Hepatitis B virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, and hepatic failure, can occur in patients treated with drugs directed against B cells. Perform screening for HBV, HCV, and HIV in accordance with clinical guidelines before collection of cells for manufacturing (1).

CD19-directed CAR-T cell therapy is supported by the National Comprehensive Cancer Network (NCCN) Guidelines for the treatment of B-cell lymphomas only after two or more chemoimmunotherapy regimens and if not previously given (2).

The safety and effectiveness of Tecartus have not been established in pediatric patients (1).

## Related policies

Breyanzi, Kymriah, Yescarta

## Policy

*This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.*

Tecartus may be considered **medically necessary** for use in patients 18 years of age and older with relapsed or refractory mantle cell lymphoma or relapsed or refractory acute lymphoblastic leukemia, and if the conditions indicated below are met.

Tecartus may be considered **investigational** in patients below 18 years of age and for all other indications.

## Prior-Approval Requirements

**Age** 18 years of age or older

## Diagnosis

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	August 27, 2020
<b>Subject:</b>	Tecartus	<b>Page:</b>	3 of 6

---

Patient must have the following:

1. Relapsed or refractory mantle cell lymphoma (MCL)
  - a. Patient must have previously received **ALL** of the following therapies:
    - i. Anthracycline **OR** bendamustine-containing chemotherapy
    - ii. Anti-CD20 monoclonal antibody
    - iii. Bruton tyrosine kinase inhibitor
  - b. Patient experienced disease progression after their last regimen **OR** refractory disease to their most recent therapy
  - c. Patient has adequate organ and bone marrow function as determined by the prescriber
  - d. **NO** history of primary central nervous system lymphoma or CNS disorders (such as seizures or cerebrovascular ischemia)
  - e. **NO** detectable cerebrospinal fluid malignant cells or brain metastases
2. B-cell precursor acute lymphoblastic leukemia (ALL)
  - a. Patient has **ONE** of the following:
    - i. Primary refractory ALL, first relapse following a remission lasting  $\leq 12$  months
    - ii. Relapsed or refractory ALL after second-line or higher therapy
    - iii. Relapsed or refractory ALL at least 100 days after allogeneic stem cell transplantation (HSCT)
  - b. Documentation of CD19 tumor expression in bone marrow or peripheral blood
  - c. Lymphoblasts  $\geq 5\%$
  - d. Patient has adequate organ function with no significant deterioration in organ function expected within 4 weeks after apheresis

**AND NONE** of the following:

- a. Burkitt lymphoma
- b. Active graft-vs-host disease
- c. Concomitant genetic syndrome associated with bone marrow failure with the exception of Down syndrome
- d. Received immunosuppressive medications within 4 weeks prior to treatment with Tecartus
- e. Active central nervous system acute lymphoblastic leukemia (ALL) (i.e., white blood cell count  $\geq 5$  cells/ $\mu$ L in cerebrospinal fluid with presence of lymphoblasts)

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	August 27, 2020
<b>Subject:</b>	Tecartus	<b>Page:</b>	4 of 6

---

- f. Received allogeneic cellular therapy, such as donor lymphocyte infusion, within six weeks prior to Tecartus infusion
- g. History of any CNS disorders, including CNS-2 disease with neurologic changes and CNS-3 disease irrespective of neurological changes

**AND ALL** of the following for **ALL** diagnoses:

- a. **NO** active infections (including TB, HBV, HCV, and HIV)
- b. Patient is not at risk for HBV infection **OR** patient is at risk for HBV infection and HBV infection has been ruled out or treatment for HBV infection has been initiated
- c. Prescriber agrees to monitor the patient for signs and symptoms of cytokine release syndrome (CRS) and administer tocilizumab if needed
- d. Prescriber agrees to monitor the patient for signs and symptoms of neurological toxicities
- e. Prescriber agrees to monitor for Hemophagocytic Lymphohistiocytosis/Macrophage activation Syndrome and treat per institutional standards
- f. Administered in a healthcare facility enrolled in the Tecartus REMS Program
- g. **NO** prior therapy with any other gene therapy (e.g., Abecma, Breyanzi, Carvykti, Kymriah, Yescarta)
- h. **NO** dual therapy with any other gene therapy (e.g., Abecma, Breyanzi, Carvykti, Kymriah, Yescarta)

## **Prior – Approval *Renewal* Requirements**

None

## [Policy Guidelines](#)

## **Pre – PA Allowance**

None

## **Prior - Approval Limits**

**Quantity** One infusion  
**Duration** 3 months (only one PA approval for one infusion per lifetime)

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	August 27, 2020
<b>Subject:</b>	Tecartus	<b>Page:</b>	5 of 6

## Rationale

### Summary

Tecartus is a CD19-directed genetically modified autologous T cell immunotherapy indicated for the treatment of adult patients with relapsed or refractory mantle cell lymphoma (MCL) and relapsed or refractory B-cell precursor acute lymphoblastic leukemia (ALL). Tecartus may cause cytokine release syndrome (CRS) and neurological toxicities. Tecartus should not be administered in patients with an active infection, central nervous system lymphoma or CNS disorders. Safety and efficacy have not been established in pediatric patients (1).

Prior approval is required to ensure the safe, clinically appropriate and cost-effective use of Tecartus while maintaining optimal therapeutic outcomes.

### Reference

1. Tecartus [package insert]. Santa Monica, CA: Kite Pharma, Inc.; October 2021.
2. NCCN Clinical Practice Guidelines in Oncology<sup>®</sup> B-Cell Lymphomas (Version 3.2022). National Comprehensive Cancer Network, Inc. April 2022. Accessed on May 10, 2022.
3. NCCN Clinical Practice Guidelines in Oncology<sup>®</sup> Acute Lymphoblastic Leukemia (Version 1.2022). National Comprehensive Cancer Network, Inc. April 2022. Accessed on May 10, 2022.

## Policy History

Date	Action
August 2020	Addition to PA
September 2020	Annual review
December 2020	Annual editorial review
March 2021	Added the requirement: No prior therapy with another CD19-directed CAR-T cell therapy, and No dual therapy with another CD19-directed CAR-T cell therapy per NCCN Guidelines. Revised PA quantity limit from 1 infusion per lifetime to 1 infusion, 3 months duration. Added clarifying statement indicating that only 1 infusion/one PA approval allowed per member's lifetime
April 2021	Revised no prior therapy and no dual therapy statements to include any other gene therapy
June 2021	Annual review and reference update
October 2021	Addition of indication: Relapsed or refractory B-cell precursor acute lymphoblastic leukemia (ALL). Added requirement "Prescriber agrees to

# 5.21.155

---

<b>Section:</b>	Prescription Drugs	<b>Effective Date:</b>	July 1, 2022
<b>Subsection:</b>	Antineoplastic Agents	<b>Original Policy Date:</b>	August 27, 2020
<b>Subject:</b>	Tecartus	<b>Page:</b>	6 of 6

---

	monitor for Hemophagocytic Lymphohistiocytosis/Macrophage activation Syndrome and treat per institutional standards”
December 2021	Annual review and reference update
March 2022	Per FEP: Added requirement for MCL to include "patient has adequate organ and bone marrow function as determined by the prescriber". Removed MCL requirement of <b>NO</b> prior allogenic hematopoietic stem cell transplant (HSCT). Added ALL requirement "patient has adequate organ function with no significant deterioration in organ function expected within 4 weeks after apheresis" and added/revised "NONE of the following: Burkitt lymphoma; Concomitant genetic syndrome associated with bone marrow failure with the exception of Down syndrome; Active central nervous system ALL; Received allogeneic cellular therapy, such as donor lymphocyte infusion, within six weeks prior to Tecartus infusion"
June 2022	Annual review and reference update. Addition of Carvykti to gene therapy requirement

## Keywords

**This policy was approved by the FEP® Pharmacy and Medical Policy Committee on June 16, 2022 and is effective on July 1, 2022.**