
5.21.163

Section:	Prescription Drugs	Effective Date:	July 1, 2022
Subsection:	Antineoplastic Agents	Original Policy Date:	October 2, 2020
Subject:	Lysodren	Page:	1 of 4

Last Review Date: June 16, 2022

Lysodren

Description

Lysodren (mitotane)

Background

Lysodren (mitotane) is an adrenal cytotoxic agent with an unknown mechanism of action. Lysodren modifies the peripheral metabolism of steroids and directly suppresses the adrenal cortex. This leads to a reduction in 17-hydroxycorticosteroids in the absence of decreased corticosteroid concentrations and increased formation of 6- β -hydroxycortisol (1).

Regulatory Status

FDA-approved indication: Lysodren is an adrenal cytotoxic agent indicated for the treatment of inoperable, functional or nonfunctional, adrenal cortical carcinoma (1).

Lysodren dose should be increased incrementally to achieve a blood concentration of 14 to 20 mg/L, or as tolerated (1).

Lysodren has a boxed warning regarding adrenal crisis in the setting of shock or severe trauma. In patients taking Lysodren, adrenal crisis occurs in the setting of shock or severe trauma and response to shock is impaired. Hydrocortisone should be administered and patient should be monitored for escalating signs of shock. Lysodren should be discontinued until recovery (1).

Lysodren also has warnings regarding CNS toxicity and adrenal insufficiency. Mitotane plasma concentrations exceeding 20 mcg/mL are associated with a greater incidence of CNS toxicity. Free cortisol and corticotropin (ACTH) levels should be measured to achieve optimal steroid replacement (1).

Section:	Prescription Drugs	Effective Date:	July 1, 2022
Subsection:	Antineoplastic Agents	Original Policy Date:	October 2, 2020
Subject:	Lysodren	Page:	2 of 4

Lysodren can cause fetal harm when administered to a pregnant woman. Pregnant women should be advised of the potential risk to a fetus. Females of reproductive potential should be advised to use effective contraception during treatment with Lysodren and after discontinuation of treatment for as long as mitotane plasma levels are detectable (1).

The safety and effectiveness of Lysodren in pediatric patients have not been established (1).

Related policies

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Lysodren may be considered **medically necessary** for use in patients 18 years of age and older with adrenal cortical carcinoma and if the conditions indicated below are met.

Lysodren may be considered **investigational** in patients below 18 years of age and for all other indications.

Prior-Approval Requirements

Age 18 years of age or older

Diagnosis

Patient must have the following:

Adrenal cortical carcinoma

AND ALL of the following:

- The tumor is inoperable
- Prescriber agrees to monitor the patient for signs and symptoms of adrenal crisis in the setting of shock or severe trauma
- Prescriber agrees to monitor for CNS toxicity
- Prescriber agrees to monitor free cortisol and corticotropin (ACTH) levels
- Female patients of reproductive potential **only**: patient will be advised to use effective contraception during treatment with Lysodren and after

Section:	Prescription Drugs	Effective Date:	July 1, 2022
Subsection:	Antineoplastic Agents	Original Policy Date:	October 2, 2020
Subject:	Lysodren	Page:	3 of 4

discontinuation of treatment for as long as mitotane plasma levels are detectable

Prior – Approval *Renewal* Requirements

Age 18 years of age or older

Diagnosis

Patient must have the following:

Adrenal cortical carcinoma

AND ALL of the following:

- NO** disease progression or unacceptable toxicity
- Prescriber agrees to monitor the patient for signs and symptoms of adrenal crisis in the setting of shock or severe trauma
- Prescriber agrees to monitor for CNS toxicity
- Prescriber agrees to monitor free cortisol and corticotropin (ACTH) levels
- Female patients of reproductive potential **only**: patient will be advised to use effective contraception during treatment with Lysodren and after discontinuation of treatment for as long as mitotane plasma levels are detectable

[Policy Guidelines](#)

Pre – PA Allowance

None

Prior - Approval Limits

Duration 12 months

Prior – Approval *Renewal* Limits

Same as above

[Rationale](#)

Summary

Section:	Prescription Drugs	Effective Date:	July 1, 2022
Subsection:	Antineoplastic Agents	Original Policy Date:	October 2, 2020
Subject:	Lysodren	Page:	4 of 4

Lysodren (mitotane) is an adrenal cytotoxic agent with an unknown mechanism of action indicated for the treatment of inoperable, functional or nonfunctional, adrenal cortical carcinoma. Lysodren modifies the peripheral metabolism of steroids and directly suppresses the adrenal cortex. This leads to a reduction in 17-hydroxycorticosteroids in the absence of decreased corticosteroid concentrations and increased formation of 6- β -hydroxycortisol (1).

Prior approval is required to ensure the safe, clinically appropriate and cost-effective use of Lysodren while maintaining optimal therapeutic outcomes.

References

1. Lysodren [package insert]. Farmingdale, NJ: Direct Success Inc.; June 2021.
2. NCCN Drugs & Biologics Compendium[®] Mitotane 2022. National Comprehensive Cancer Network, Inc. Accessed on April 25, 2022.

Policy History

Date	Action
October 2020	Addition to PA
December 2020	Annual review
June 2021	Annual review and reference update
June 2022	Annual review and reference update

Keywords

This policy was approved by the FEP[®] Pharmacy and Medical Policy Committee on June 16, 2022 and is effective on July 1, 2022.