



## FEP Medical Policy Manual

### FEP 7.01.106 Percutaneous Tibial Nerve Stimulation

**Effective Policy Date: January 1, 2022**

**Original Policy Date: December 2012**

**Related Policies:**

- 1.01.17 - Pelvic Floor Stimulation as a Treatment of Urinary and Fecal Incontinence
- 5.75.01 - Botox (onabotulinum toxin A)
- 7.01.19 - Injectable Bulking Agents for the Treatment of Urinary and Fecal Incontinence
- 7.01.29 - Percutaneous Electrical Nerve Stimulation and Percutaneous Neuromodulation Therapy

## Percutaneous Tibial Nerve Stimulation

### Description

Percutaneous tibial nerve stimulation (PTNS; also known as posterior tibial nerve stimulation) is an electrical neuromodulation technique used primarily for treating voiding dysfunction.

The current indication cleared by the U.S. Food and Drug Administration (FDA) for PTNS is overactive bladder and associated symptoms of urinary frequency, urinary urgency, and urge incontinence.

Altering the function of the posterior tibial nerve with PTNS is believed to improve voiding function and control. The mechanism of action is believed to be retrograde stimulation of the lumbosacral nerves (L4-S3) via the posterior tibial nerve located near the ankle. The lumbosacral nerves control the bladder detrusor and perineal floor.

Administration of PTNS consists of inserting a needle above the medial malleolus into the posterior tibial nerve followed by the application of low-voltage (10 mA, 1-10 Hz frequency) electrical stimulation that produces sensory and motor responses as evidenced by a tickling sensation and plantarflexion or fanning of all toes. Noninvasive PTNS has also been delivered with transcutaneous or surface electrodes. The recommended course of treatment is an initial series of 12 weekly office-based treatments followed by an individualized maintenance treatment schedule.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

Percutaneous tibial nerve stimulation is less invasive than traditional sacral nerve neuromodulation (see evidence review 7.01.69), which has been successfully used to treat urinary dysfunction but requires implantation of a permanent device. In sacral root neuromodulation, an implantable pulse generator that delivers controlled electrical impulses is attached to wire leads that connect to the sacral nerves, most commonly the S3 nerve root that modulates the neural pathways controlling bladder function.

Percutaneous tibial nerve stimulation has also been proposed as a treatment for non-neurogenic and neurogenic bladder syndromes and fecal incontinence.

## OBJECTIVE

The objective of this evidence review is to determine whether the use of percutaneous tibial nerve stimulation improves the net health outcome in individuals who have urinary dysfunction associated with overactive bladder syndrome, neurogenic bladder, or fecal incontinence.

## POLICY STATEMENT

Percutaneous tibial nerve stimulation for an initial 12-week course is considered **medically necessary** for individuals with non-neurogenic urinary dysfunction including overactive bladder who have both:

- failed behavioral therapy following an appropriate duration of 8 to 12 weeks without meeting treatment goals; and
- failed pharmacologic therapy following 4 to 8 weeks of treatment without meeting treatment goals.

Maintenance therapy using monthly percutaneous tibial nerve stimulation is considered **medically necessary** for individuals following a 12-week initial course of percutaneous tibial nerve stimulation that resulted in improved urinary dysfunction meeting treatment goals.

Percutaneous tibial nerve stimulation is considered **investigational** for all other indications, including but not limited to the following:

- Neurogenic bladder dysfunction
- Fecal incontinence.

## POLICY GUIDELINES

Patients may be considered to have failed behavioral therapies following an appropriate duration of 8 to 12 weeks without meeting treatment goals.

Patients may be considered to have failed pharmacologic therapies following 4 to 8 weeks of treatment without meeting treatment goals.

Annual evaluation by a physician may be performed to ensure efficacy is continuing for maintenance percutaneous tibial nerve stimulation treatments.

## BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

## FDA REGULATORY STATUS

In 2005, the Urgent PC Neuromodulation System was the initial PTNS device cleared for marketing by FDA through the 510(k) process to treat patients suffering from urinary urgency, urinary frequency, and urge incontinence. Additional PTNS devices have been cleared for marketing through the 510(k) process. They are listed in Table 1.

The Urgent PC Neuromodulation System, NURO™ Neuromodulation System, and ZIDA Wearable Neuromodulation System are not FDA cleared for other indications, such as the treatment of fecal incontinence.

Wireless technology is evolving for the treatment of overactive bladder; it is approved in Europe. BlueWind (BlueWind Medical) is a wireless, battery-less, miniature implantable neurostimulator activated by an external device worn at the ankle.

**Table 1. FDA-Cleared Percutaneous Tibial Nerve Stimulators (FDA Product Code: NAM)**

Device Name	Manufacturer	Cleared	510(k)	Indications
Urgent PC Neuromodulation System	Uroplasty, now Cogentix Medical	Oct 2005	K052025	Treatment of urinary urgency, urinary frequency, and urge incontinence
Urgent PC Neuromodulation System	Uroplasty, now Cogentix Medical	Jul 2006	K061333	FDA determined the 70% isopropyl alcohol prep pad contained in the kit is subject to regulation as a drug
Urgent PC Neuromodulation System	Uroplasty, now Cogentix Medical	Aug 2007	K071822	Labeling update, intended use is unchanged
Urgent PC Neuromodulation System	Uroplasty, now Cogentix Medical	Oct 2010	K101847	Intended use statement adds the diagnosis of overactive bladder
NURO™ Neuromodulation System	Advanced Uro-Solutions, now Medtronic	Nov 2013	K132561	Treatment of patients with overactive bladder and associated symptoms of urinary urgency, urinary frequency, and urge incontinence
ZIDA Wearable Neuromodulation System	Exodus Innovations	Mar 2021	K192731	Treatment of patients with an overactive bladder and associated symptoms of urinary urgency, urinary frequency, and urge incontinence

FDA: U.S. Food and Drug Administration.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

## RATIONALE

### Summary of Evidence

For individuals who have non-neurogenic urinary dysfunction including overactive bladder and have failed behavioral and pharmacologic therapy who receive an initial course of Percutaneous tibial nerve stimulation (PTNS), the evidence includes randomized sham-controlled trials, randomized controlled trials (RCTs) with an active comparator, and systematic reviews. Relevant outcomes are symptoms, change in disease status, functional outcomes, quality of life, and treatment-related morbidity. The SUmIT and OrBIT trials are 2 key industry-sponsored RCTs. Systematic reviews that included these and other published trials have found short-term reductions in voiding dysfunction with PTNS. The largest, highest quality study was the double-blind, sham-controlled SUmIT trial, which reported a statistically significant benefit of PTNS versus sham at 12 weeks. In an additional, small sham-controlled trial, a 50% reduction in urge incontinent episodes was attained in 71% of the PTNS group compared with 0% in the sham group. The nonblinded OrBIT trial found that PTNS was noninferior to medication therapy at 12 weeks. Adverse events were limited to local irritation effects. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have overactive bladder syndrome that has failed behavioral and pharmacologic therapy who respond to an initial course of PTNS who receive maintenance PTNS, the evidence includes observational studies and systematic reviews. Relevant outcomes are symptoms, change in disease status, functional outcomes, quality of life, and treatment-related morbidity. The SUmIT and OrBIT trials each included extension studies that followed individuals who responded to the initial course of PTNS and continued to receive periodic maintenance therapy. There is variability in the interval between and frequency of maintenance treatments, and an optimal maintenance regimen remains unclear. There are up to 36 months of observational data available, reporting that there is a durable effect for some of these patients. While comparative data are not available after the initial 12-week treatment period, the observational data support a clinically meaningful benefit for use in individuals who have already failed behavioral and pharmacologic therapy and who respond to the initial course of PTNS. Percutaneous tibial nerve stimulation may allow such individuals to avoid more invasive interventions. Adverse events appear to be limited to local irritation for both short- and long-term PTNS use. Typical regimens schedule maintenance treatments every 4-6 weeks. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have neurogenic bladder dysfunction who receive PTNS, the evidence includes several RCTs and a systematic review of RCTs and observational data. Relevant outcomes are symptoms, change in disease status, functional outcomes, quality of life, and treatment-related morbidity. Only a few RCTs evaluating tibial nerve stimulation for treating neurogenic bladder have been published to date, and all but 1 performed transcutaneous stimulation rather than PTNS. Studies varied widely in factors such as study populations and comparator interventions. Study findings have not reported that tibial nerve stimulation significantly reduced incontinence symptoms and improved other outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have fecal incontinence who receive PTNS, the evidence includes several RCTs and systematic reviews. Relevant outcomes are symptoms, change in disease status, functional outcomes, quality of life, and treatment-related morbidity. The available RCTs have not found a clear benefit of PTNS. Neither of the sham-controlled trials found that active stimulation was superior to sham for achieving the primary outcome, at least a 50% reduction in mean weekly fecal incontinence episodes. The larger sham-controlled randomized trial did find a significantly greater decrease in the absolute number of weekly incontinence episodes in the active treatment group, but the overall trial findings did not suggest the superiority of PTNS over sham treatment. A meta-analysis of a single RCT and several observational studies reported that patients receiving sacral nerve stimulation experienced significant benefits compared with patients receiving PTNS. A post hoc analysis of the larger trial suggested a subset of patients with fecal incontinence (those without concomitant obstructive defecation) may benefit from PTNS. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

## SUPPLEMENTAL INFORMATION

### Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in "Supplemental Information" if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

## American Urological Association et al

In 2019, the American Urological Association and the Society of Urodynamics, Female Pelvic Medicine & Urogenital Reconstruction published updated guidelines on the diagnosis and treatment of non-neurogenic overactive bladder in adults.<sup>44</sup> The guidelines included a statement that clinicians may offer PTNS as a third-line treatment option in carefully selected patients. The statement carried a grade C rating, indicating that the balance of benefits and risks/burdens are uncertain.

## American College of Obstetricians and Gynecologists

In 2015, the American College of Obstetricians and Gynecologists practice bulletin on the treatment of urinary incontinence in women did not address PTNS or other types of nerve stimulation.<sup>45</sup>

## American Gastroenterological Association

In 2017, the American Gastroenterological Association issued an expert review and clinical practice update on surgical interventions and device-aided therapy for the treatment of fecal incontinence.<sup>46</sup> The update stated that "until further evidence is available, percutaneous tibial nerve stimulation should not be used for managing FI [fecal incontinence] in clinical practice."

## U.S. Preventive Services Task Force Recommendations

Not applicable.

## Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

## REFERENCES

1. Wang M, Jian Z, Ma Y, et al. Percutaneous tibial nerve stimulation for overactive bladder syndrome: a systematic review and meta-analysis. *Int Urogynecol J*. Dec 2020; 31(12): 2457-2471. PMID 32681345
2. Xiong SC, Peng L, Hu X, et al. Effectiveness and safety of tibial nerve stimulation versus anticholinergic drugs for the treatment of overactive bladder syndrome: a meta-analysis. *Ann Palliat Med*. Jun 09 2021. PMID 34118839
3. Coolen RL, Groen J, Scheepe JR, et al. Transcutaneous Electrical Nerve Stimulation and Percutaneous Tibial Nerve Stimulation to Treat Idiopathic Nonobstructive Urinary Retention: A Systematic Review. *Eur Urol Focus*. Oct 22 2020. PMID 33268327
4. Ho FCS, He C, Yao HH, et al. Efficacy of sacral neuromodulation and percutaneous tibial nerve stimulation in the treatment of chronic nonobstructive urinary retention: A systematic review. *Neurourol Urodyn*. Jun 2021; 40(5): 1078-1088. PMID 33973670
5. Tutolo M, Ammirati E, Heesakkers J, et al. Efficacy and Safety of Sacral and Percutaneous Tibial Neuromodulation in Non-neurogenic Lower Urinary Tract Dysfunction and Chronic Pelvic Pain: A Systematic Review of the Literature. *Eur Urol*. Mar 2018; 73(3): 406-418. PMID 29336927
6. Tutolo M, Ammirati E, Van der Aa F. What Is New in Neuromodulation for Overactive Bladder?. *Eur Urol Focus*. Jan 2018; 4(1): 49-53. PMID 29773501
7. Stewart F, Gameiro LF, El Dib R, et al. Electrical stimulation with non-implanted electrodes for overactive bladder in adults. *Cochrane Database Syst Rev*. Dec 09 2016; 12: CD010098. PMID 27935011
8. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Percutaneous tibial nerve stimulation for the treatment of voiding dysfunction. *TEC Assessments*. 2013;Volume 28:Tab 10. PMID
9. Burton C, Sajja A, Latthe PM. Effectiveness of percutaneous posterior tibial nerve stimulation for overactive bladder: a systematic review and meta-analysis. *Neurourol Urodyn*. Nov 2012; 31(8): 1206-16. PMID 22581511
10. Levin PJ, Wu JM, Kawasaki A, et al. The efficacy of posterior tibial nerve stimulation for the treatment of overactive bladder in women: a systematic review. *Int Urogynecol J*. Nov 2012; 23(11): 1591-7. PMID 22411208
11. Moosdorff-Steinhauser HF, Berghmans B. Effects of percutaneous tibial nerve stimulation on adult patients with overactive bladder syndrome: a systematic review. *Neurourol Urodyn*. Mar 2013; 32(3): 206-14. PMID 22907807

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

12. Gaziev G, Topazio L, Iacovelli V, et al. Percutaneous Tibial Nerve Stimulation (PTNS) efficacy in the treatment of lower urinary tract dysfunctions: a systematic review. *BMC Urol.* Nov 25 2013; 13: 61. PMID 24274173
13. Shamliyan T, Wyman J, Kane RL. Nonsurgical Treatments for Urinary Incontinence in Adult Women: Diagnosis and Comparative Effectiveness (Comparative Effectiveness Review No. 36). Rockville, MD: Agency for Healthcare Research and Quality; 2012.
14. Finazzi-Agro E, Petta F, Sciobica F, et al. Percutaneous tibial nerve stimulation effects on detrusor overactivity incontinence are not due to a placebo effect: a randomized, double-blind, placebo controlled trial. *J Urol.* Nov 2010; 184(5): 2001-6. PMID 20850833
15. Peters KM, Carrico DJ, Perez-Marrero RA, et al. Randomized trial of percutaneous tibial nerve stimulation versus Sham efficacy in the treatment of overactive bladder syndrome: results from the SUmIT trial. *J Urol.* Apr 2010; 183(4): 1438-43. PMID 20171677
16. Peters K, Carrico D, Burks F. Validation of a sham for percutaneous tibial nerve stimulation (PTNS). *NeuroUrol Urodyn.* 2009; 28(1): 58-61. PMID 18671297
17. Peters KM, Carrico DJ, Wooldridge LS, et al. Percutaneous tibial nerve stimulation for the long-term treatment of overactive bladder: 3-year results of the STEP study. *J Urol.* Jun 2013; 189(6): 2194-201. PMID 23219541
18. Vecchioli-Scaldazza C, Morosetti C. Effectiveness and durability of solifenacin versus percutaneous tibial nerve stimulation versus their combination for the treatment of women with overactive bladder syndrome: a randomized controlled study with a follow-up of ten months. *Int Braz J Urol.* Jan-Feb 2018; 44(1): 102-108. PMID 29064651
19. Boudaoud N, Binet A, Line A, et al. Management of refractory overactive bladder in children by transcutaneous posterior tibial nerve stimulation: A controlled study. *J Pediatr Urol.* Jun 2015; 11(3): 138.e1-10. PMID 25979217
20. Gungor Ugurlucan F, Onal M, Aslan E, et al. Comparison of the effects of electrical stimulation and posterior tibial nerve stimulation in the treatment of overactive bladder syndrome. *Gynecol Obstet Invest.* 2013; 75(1): 46-52. PMID 23171636
21. Preyer O, Umek W, Laml T, et al. Percutaneous tibial nerve stimulation versus tolterodine for overactive bladder in women: a randomised controlled trial. *Eur J Obstet Gynecol Reprod Biol.* Aug 2015; 191: 51-6. PMID 26073262
22. Vecchioli-Scaldazza C, Morosetti C, Berouz A, et al. Solifenacin succinate versus percutaneous tibial nerve stimulation in women with overactive bladder syndrome: results of a randomized controlled crossover study. *Gynecol Obstet Invest.* 2013; 75(4): 230-4. PMID 23548260
23. Schreiner L, dos Santos TG, Knorst MR, et al. Randomized trial of transcutaneous tibial nerve stimulation to treat urge urinary incontinence in older women. *Int Urogynecol J.* Sep 2010; 21(9): 1065-70. PMID 20458465
24. Peters KM, Macdiarmid SA, Wooldridge LS, et al. Randomized trial of percutaneous tibial nerve stimulation versus extended-release tolterodine: results from the overactive bladder innovative therapy trial. *J Urol.* Sep 2009; 182(3): 1055-61. PMID 19616802
25. MacDiarmid SA, Peters KM, Shobeiri SA, et al. Long-term durability of percutaneous tibial nerve stimulation for the treatment of overactive bladder. *J Urol.* Jan 2010; 183(1): 234-40. PMID 19913821
26. Schneider MP, Gross T, Bachmann LM, et al. Tibial Nerve Stimulation for Treating Neurogenic Lower Urinary Tract Dysfunction: A Systematic Review. *Eur Urol.* Nov 2015; 68(5): 859-67. PMID 26194043
27. Monteiro ES, de Carvalho LB, Fukujima MM, et al. Electrical stimulation of the posterior tibialis nerve improves symptoms of poststroke neurogenic overactive bladder in men: a randomized controlled trial. *Urology.* Sep 2014; 84(3): 509-14. PMID 25168524
28. Perissinotto MC, D'Ancona CA, Lucio A, et al. Transcutaneous tibial nerve stimulation in the treatment of lower urinary tract symptoms and its impact on health-related quality of life in patients with Parkinson disease: a randomized controlled trial. *J Wound Ostomy Continence Nurs.* Jan-Feb 2015; 42(1): 94-9. PMID 25549314
29. Gaspard L, Tombal B, Opsomer RJ, et al. [Physiotherapy and neurogenic lower urinary tract dysfunction in multiple sclerosis patients: a randomized controlled trial]. *Prog Urol.* Sep 2014; 24(11): 697-707. PMID 25214451
30. Eftekhari T, Teimoori N, Miri E, et al. Posterior tibial nerve stimulation for treating neurologic bladder in women: a randomized clinical trial. *Acta Med Iran.* 2014; 52(11): 816-21. PMID 25415813
31. Zonic-Imamovic M, Imamovic S, Cickusic A, et al. Effects of Treating an Overactive Urinary Bladder in Patients with Multiple Sclerosis. *Acta Med Acad.* Dec 2019; 48(3): 271-277. PMID 32124625
32. Welk B, McKibbin M. A randomized, controlled trial of transcutaneous tibial nerve stimulation to treat overactive bladder and neurogenic bladder patients. *Can Urol Assoc J.* Jul 2020; 14(7): E297-E303. PMID 32017693
33. Sarveazad A, Babahajian A, Amini N, et al. Posterior Tibial Nerve Stimulation in Fecal Incontinence: A Systematic Review and Meta-Analysis. *Basic Clin Neurosci.* Sep-Oct 2019; 10(5): 419-431. PMID 32284831
34. Tan K, Wells CI, Dinning P, et al. Placebo Response Rates in Electrical Nerve Stimulation Trials for Fecal Incontinence and Constipation: A Systematic Review and Meta-Analysis. *Neuromodulation.* Dec 2020; 23(8): 1108-1116. PMID 31889364
35. Simillis C, Lal N, Qiu S, et al. Sacral nerve stimulation versus percutaneous tibial nerve stimulation for faecal incontinence: a systematic review and meta-analysis. *Int J Colorectal Dis.* May 2018; 33(5): 645-648. PMID 29470730
36. Edenfield AL, Amundsen CL, Wu JM, et al. Posterior tibial nerve stimulation for the treatment of fecal incontinence: a systematic evidence review. *Obstet Gynecol Surv.* May 2015; 70(5): 329-41. PMID 25974730
37. Horrocks EJ, Thin N, Thaha MA, et al. Systematic review of tibial nerve stimulation to treat faecal incontinence. *Br J Surg.* Apr 2014; 101(5): 457-68. PMID 24446127
38. George AT, Kalmar K, Sala S, et al. Randomized controlled trial of percutaneous versus transcutaneous posterior tibial nerve stimulation in faecal incontinence. *Br J Surg.* Feb 2013; 100(3): 330-8. PMID 23300071
39. Knowles CH, Horrocks EJ, Bremner SA, et al. Percutaneous tibial nerve stimulation versus sham electrical stimulation for the treatment of faecal incontinence in adults (CONFIDENT): a double-blind, multicentre, pragmatic, parallel-group, randomised controlled trial. *Lancet.* Oct 24 2015; 386(10004): 1640-8. PMID 26293315

40. Horrocks EJ, Chadi SA, Stevens NJ, et al. Factors Associated With Efficacy of Percutaneous Tibial Nerve Stimulation for Fecal Incontinence, Based on Post-Hoc Analysis of Data From a Randomized Trial. Clin Gastroenterol Hepatol. Dec 2017; 15(12): 1915-1921.e2. PMID 28647458
41. Thin NN, Taylor SJ, Bremner SA, et al. Randomized clinical trial of sacral versus percutaneous tibial nerve stimulation in patients with faecal incontinence. Br J Surg. Mar 2015; 102(4): 349-58. PMID 25644291
42. Leo CA, Thomas GP, Hodgkinson JD, et al. Randomized Pilot Study: Anal Inserts Versus Percutaneous Tibial Nerve Stimulation in Patients With Fecal Incontinence. Dis Colon Rectum. Apr 01 2021; 64(4): 466-474. PMID 33399411
43. Sanagapalli S, Neilan L, Lo JYT, et al. Efficacy of Percutaneous Posterior Tibial Nerve Stimulation for the Management of Fecal Incontinence in Multiple Sclerosis: A Pilot Study. Neuromodulation. Oct 2018; 21(7): 682-687. PMID 29575432
44. Lightner DJ, Gomelsky A, Souter L, et al. Diagnosis and Treatment of Overactive Bladder (Non-Neurogenic) in Adults: AUA/SUFU Guideline Amendment 2019. J Urol. Sep 2019; 202(3): 558-563. PMID 31039103
45. ACOG Practice Bulletin No. 155: Urinary Incontinence in Women. Obstet Gynecol. Nov 2015; 126(5): e66-e81. PMID 26488524
46. Bharucha AE, Rao SSC, Shin AS. Surgical Interventions and the Use of Device-Aided Therapy for the Treatment of Fecal Incontinence and Defecatory Disorders. Clin Gastroenterol Hepatol. Dec 2017; 15(12): 1844-1854. PMID 28838787

**POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:**

Date	Action	Description
December 2012	New policy	
September 2013	Replace policy	Policy updated with literature review. References 5, 6, 8-10, 12-15, 17 and 20 added; other references renumbered or removed. Policy statement unchanged.
September 2014	Replace policy	Policy updated with literature review; reference 1 added, 5 updated; policy statement unchanged.
September 2015	Replace policy	Policy updated with literature review; Title changed to "Percutaneous Tibial Nerve Stimulation." "Posterior" changed to "percutaneous" in existing policy statement. Policy statement edited to not medically necessary for all indications with bullet points for urinary and fecal incontinence. Reference 17-19 added.
June 2016	Replace policy	Policy updated with literature review through November 30, 2015; references 15, 17, 19-25, 27-28, and 30-31 added. Policy statements unchanged.
June 2018	Replace policy	Policy updated with literature review through September 15, 2017; reference 18 added; reference 31 updated. Revised policy statements for use of PTNS in OAB syndrome that has failed behavioral and pharmacologic therapy. In these patients, PTNS is considered medically necessary as an initial course of therapy and maintenance therapy for individuals who respond to initial course. Percutaneous tibial nerve stimulation changed from not medically necessary to investigational (due to FDA 510k approval status) for all other indications, including but not limited to the following: Neurogenic bladder dysfunction; Fecal incontinence.
December 2018		Policy updated with literature review through June 4, 2018; references 13-14, 27, 32, 34, and 37 added. Policy statements are unchanged.
December 2019	Replace policy	Policy updated with literature review through May 31, 2019; references added. Policy statements unchanged.
December 2020	Replace policy	Policy updated with literature review through May 28, 2020; references added. Policy statements unchanged.
December 2021	Replace policy	Policy updated with literature review through June 24, 2021; references added. Policy statements unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.