



## FEP Medical Policy Manual

### FEP 2.01.39 Quantitative Sensory Testing

**Effective Policy Date: October 1, 2022**

**Original Policy Date: September 2012**

**Related Policies:**

None

## Quantitative Sensory Testing

### Description

#### Description

Quantitative sensory testing (QST) systems are used for the noninvasive assessment and quantification of sensory nerve function in patients with symptoms of, or the potential for neurologic damage or disease. Types of sensory testing include current perception threshold testing, pressure-specified sensory testing, vibration perception testing (VPT), and thermal sensory testing. Information on sensory deficits identified using QST has been used in research settings to understand neuropathic pain better. It could be used to diagnose conditions linked to nerve damage and disease, and to improve patient outcomes by impacting management strategies.

Quantitative sensory test systems measure and quantify the amount of physical stimuli required for sensory perception to occur. As sensory deficits increase, the perception threshold of QST will increase, which may be informative in documenting the progression of neurologic damage or disease. Currently, QST has not been established for use as a sole tool for diagnosis and management but has been used with standard evaluative and management procedures (eg, physical and neurologic examination, monofilament testing, pinprick, grip and pinch strength, Tinel sign, and Phalen and Roos test) to enhance the diagnosis and treatment-planning process, and to confirm physical findings with quantifiable data. Stimuli used in QST include touch, pressure, pain, thermal (warm and cold), or vibratory stimuli.

The criterion standard for evaluation of myelinated, large fibers is the electromyography nerve conduction study. However, the function of smaller myelinated and unmyelinated sensory nerves, which may show pathologic changes before the involvement of the motor nerves, cannot be detected by

nerve conduction studies. Small fiber neuropathy has traditionally been a diagnosis of exclusion in patients who have symptoms of distal neuropathy and a negative nerve conduction study.

Depending on the type of stimuli used, QST can assess both small and large fiber dysfunction. Touch and vibration measure the function of large myelinated A alpha and A beta sensory fibers. Thermal stimulation devices are used to evaluate pathology of small myelinated and unmyelinated nerve fibers; they can be used to assess heat and cold sensation, as well as thermal pain thresholds. Pressure-specified sensory devices assess large myelinated sensory nerve function by quantifying the thresholds of pressure detected with light, static, and moving touch. Finally, current perception threshold testing involves the quantification of the sensory threshold to transcutaneous electrical stimulation. In current perception threshold testing, typically 3 frequencies are tested: 5 Hz, designed to assess C fibers; 250 Hz, designed to assess A delta fibers; and 2000 Hz, designed to assess A beta fibers. Results are compared with those of a reference population.

Because QST combines the objective physical, sensory stimuli with the subject patient response, it is psychophysical and requires patients who are alert, able to follow directions, and cooperative. Also, to get reliable results, examinations need to include standardized instructions to the patients, and stimuli must be applied consistently by trained staff. Psychophysical tests have greater inherent variability, making their results more difficult to reproduce.

Primarily, QST has been applied in patients with conditions associated with nerve damage and neuropathic pain. A retrospective analysis of a prospective database maintained by the German Research Network on Neuropathic Pain by Forstenpointner et al (2021) compared QST profiles between patients with painful neuropathic conditions (n=332), patients with neuropathic conditions who did not report pain (n=111), and healthy controls (n=112). After extensive QST testing, including thermal, mechanical/vibration, and pain sensitivity, the researchers found similar QST profiles between patients who reported pain and patients who did not report pain, which raises concern about the role of QST in general in decision-making for neuropathic conditions.<sup>3</sup> There have also been preliminary investigations to identify sensory deficits associated with conditions such as autism spectrum disorder, Tourette syndrome, restless legs syndrome, musculoskeletal pain, and response to opioid treatment.

## OBJECTIVE

The objective of this evidence review is to determine whether quantitative sensory testing improves the net health outcome in patients with conditions linked to nerve damage or disease.

## POLICY STATEMENT

Quantitative sensory testing, including but not limited to current perception threshold testing, pressure-specified sensory device testing, vibration perception threshold testing, and thermal threshold testing, is considered **investigational**.

## POLICY GUIDELINES

None

## BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

## FDA REGULATORY STATUS

A number of QST devices have been cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process. Examples are listed in Table 1.

**Table 1. FDA Approved Quantitative Sensory Testing Devices**

Device	Manufacturer	Date Cleared	510(k)	Indications
FDA product code: LLN				
Neurometer	Neurotron	Jun 1986	K853608	Current perception threshold testing
NK Pressure-Specified Sensory Device, Model PSSD	NK Biotechnical Engineering	Aug 1994	K934368	Pressure-specified sensory testing
AP-4000, Air Pulse Sensory Stimulator	Pentax Precision Instrument	Sep 1997	K964815	Pressure-specified sensory testing
Neural-Scan	Neuro-Diagnostic Assoc.	Dec 1997	K964622	Current perception threshold testing
Vibration Perception Threshold (VPT) METER	Xilas Medical	Dec 2003	K030829	Vibration perception testing
Pain Vision, Model PS-2100	Osachi Co., LTD	Jan 2009	K072882	Current perception threshold testing
FDA product code: NTU				
Contact Heat-Evoked Potential Stimulator (Cheps)	Medoc, Advanced Medical Systems	Feb 2005	K041908	Thermal sensory testing
Modified Contact-Heat Evoked Potential Stimulator (Cheps)	Medoc, Advanced Medical Systems	Jun 2005	K051448	Thermal sensory testing
Pathway - Ats/Cheps	Medoc, Advanced Medical Systems	Jan 2006	K052357	Thermal sensory testing

FDA: U.S. Food and Drug Administration.

## RATIONALE

### Summary of Evidence

For individuals who have conditions linked to nerve damage or disease (eg, diabetic neuropathy, carpal tunnel syndrome) who receive current perception threshold testing, the evidence includes several studies on technical performance and diagnostic accuracy. Relevant outcomes are test accuracy and validity, symptoms, and functional outcomes. The existing evidence does not support the accuracy of current perception threshold testing for diagnosing any condition linked to nerve damage or disease. Studies comparing current perception threshold testing with other testing methods have not reported on sensitivity or specificity. Also, there is a lack of direct evidence on the clinical utility of current perception testing and, because there is insufficient evidence on test performance, an indirect chain of evidence on clinical utility cannot be constructed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have conditions linked to nerve damage or disease (eg, diabetic neuropathy, carpal tunnel syndrome) who receive pressure-specified sensory testing, the evidence includes several studies on diagnostic accuracy. Relevant outcomes are test accuracy and validity, symptoms, and functional outcomes. Current evidence does not support the diagnostic accuracy of pressure-specified sensory testing for diagnosing any condition linked to nerve damage or disease. A systematic review found that pressure-specified sensory testing had low accuracy for diagnosing spinal conditions. Also, there is a lack of direct evidence on the clinical utility of pressure-specified sensory testing and, because there is insufficient evidence on test performance, an indirect chain of evidence on clinical utility cannot be constructed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have conditions linked to nerve damage or disease (eg, diabetic neuropathy, carpal tunnel syndrome) who receive vibration perception testing (VPT), the evidence includes several studies on diagnostic accuracy. Relevant outcomes are test accuracy and validity, symptoms, and functional outcomes. A few studies have assessed the diagnostic performance of vibration testing using devices not cleared by the U.S. Food and Drug Administration (FDA). Also, there is a lack of direct evidence on the clinical utility of VPT and, in the absence of sufficient evidence on test performance, an indirect chain of evidence on clinical utility cannot be constructed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have conditions linked to nerve damage or disease (eg, diabetic neuropathy, carpal tunnel syndrome) who receive thermal sensory testing, the evidence includes diagnostic accuracy studies. Relevant outcomes are test accuracy and validity, symptoms, and functional outcomes. Two studies identified evaluated the diagnostic accuracy of thermal quantitative sensory testing (QST) using the same FDA -cleared device. Neither found a high diagnostic accuracy for thermal QST but both studies found the test had potential when used with other tests. An additional study using a different device also supports the potential of thermal QST in combination with other tests. The optimal combination of tests is currently unclear. Also, there is a lack of direct evidence on the clinical utility of thermal sensory testing and, because there is insufficient evidence on test performance, an indirect chain of evidence on clinical utility cannot be constructed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

## SUPPLEMENTAL INFORMATION

### Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

#### American Academy of Neurology

The American Academy of Neurology (2003; reaffirmed 2022 ) concluded that quantitative sensory testing (QST) is probably (level B recommendation) an effective tool for documenting of sensory abnormalities and changes in sensory thresholds in longitudinal evaluation of patients with diabetic neuropathy.<sup>20,21</sup> Evidence was weak or insufficient to support the use of QST in patients with other conditions (small fiber sensory neuropathy, pain syndromes, toxic neuropathies, uremic neuropathy, acquired and inherited demyelinating neuropathies, or malingering).

#### American Association of Neuromuscular & Electrodiagnostic Medicine

In 2004, the American Association of Neuromuscular & Electrodiagnostic Medicine (AANEM) published a technology literature review on QST (light touch, vibration, thermal, pain).<sup>22</sup> The review concluded that QST is a reliable psychophysical test of large- and small-fiber sensory modalities but is highly dependent on the full patient cooperation. Abnormalities do not localize dysfunction to the central or peripheral nervous system, and no algorithm can reliably distinguish between psychogenic and organic abnormalities. The AANEM review also indicated that QST had been shown to be reasonably reproducible over a period of days or weeks in normal subjects, but, for individual patients, more studies are needed to determine the maximum allowable difference between 2 quantitative sensory tests that can be attributed to experimental error.

In 2005, the AANEM with American Academy of Neurology and American Academy of Physical Medicine & Rehabilitation developed a formal case definition of distal symmetrical polyneuropathy based on a systematic analysis of peer-reviewed literature supplemented by consensus from an expert panel.<sup>23</sup> QST was not included as part of the final case definition, given that the reproducibility of QST ranged from poor to excellent, and the sensitivities and specificities of QST varied widely among studies.

## American Diabetes Association

In 2021, the American Diabetes Association published an updated standard for microvascular complications and foot care.<sup>24</sup> Although temperature and vibration testing are recommended as part of the evaluation of small fiber and large fiber function, respectively, the specific screening tests for diabetic peripheral neuropathy that are described in the standard are manual/clinical rather than quantitative. Therefore, QST does not appear to have a role in the routine evaluation or diagnosis of diabetic peripheral neuropathy.

## U.S. Preventive Services Task Force Recommendations

Not applicable.

## Medicare National Coverage

Medicare (2002) announced a national noncoverage policy on sensory nerve conduction threshold testing. Medicare reconsidered its policy, but affirmed it, concluding that any use of sensory nerve conduction threshold testing to diagnose sensory neuropathies or radiculopathies is not reasonable and necessary. This decision was reaffirmed in 2004.<sup>25</sup> Medicare has not addressed coverage for other types of QST.

## REFERENCES

- Spanakis EK, Golden SH. Race/ethnic difference in diabetes and diabetic complications. *Curr Diab Rep*. Dec 2013; 13(6): 814-23. PMID 24037313
- Taylor YJ, Davis ME, Mahabaleshwarkar R et al. Racial/ethnic disparities in diabetes care and outcomes: A mixed methods study. *Journal of Health Disparities Research and Practice*. 2018; 11(2). Accessed April 27, 2022. <https://digitalscholarship.unlv.edu/jhdrp/vol11/iss2/9>
- Forstenpointner J, Ruscheweyh R, Attal N, et al. No pain, still gain (of function): the relation between sensory profiles and the presence or absence of self-reported pain in a large multicenter cohort of patients with neuropathy. *Pain*. Mar 01 2021; 162(3): 718-727. PMID 32868752
- Ziccardi VB, Dragoo J, Eliav E, et al. Comparison of current perception threshold electrical testing to clinical sensory testing for lingual nerve injuries. *J Oral Maxillofac Surg*. Feb 2012; 70(2): 289-94. PMID 22079068
- Park R, Wallace MS, Schulteis G. Relative sensitivity to alfentanil and reliability of current perception threshold vs von Frey tactile stimulation and thermal sensory testing. *J Peripher Nerv Syst*. Dec 2001; 6(4): 232-40. PMID 11800047
- Weber RA, Schuchmann JA, Albers JH, et al. A prospective blinded evaluation of nerve conduction velocity versus Pressure-Specified Sensory Testing in carpal tunnel syndrome. *Ann Plast Surg*. Sep 2000; 45(3): 252-7. PMID 10987525
- Nath RK, Bowen ME, Eichhorn MG. Pressure-specified sensory device versus electrodiagnostic testing in brachial plexus upper trunk injury. *J Reconstr Microsurg*. May 2010; 26(4): 235-42. PMID 20143301
- Hubscher M, Moloney N, Leaver A, et al. Relationship between quantitative sensory testing and pain or disability in people with spinal pain—a systematic review and meta-analysis. *Pain*. Sep 2013; 154(9): 1497-1504. PMID 23711482
- Suokas AK, Walsh DA, McWilliams DF, et al. Quantitative sensory testing in painful osteoarthritis: a systematic review and meta-analysis. *Osteoarthritis Cartilage*. Oct 2012; 20(10): 1075-85. PMID 22796624
- Mythili A, Kumar KD, Subrahmanyam KA, et al. A Comparative study of examination scores and quantitative sensory testing in diagnosis of diabetic polyneuropathy. *Int J Diabetes Dev Ctries*. Jan 2010; 30(1): 43-8. PMID 20431806
- Abraham A, Albulaihe H, Alabdali M, et al. Elevated Vibration Perception Thresholds in CIDP Patients Indicate More Severe Neuropathy and Lower Treatment Response Rates. *PLoS One*. 2015; 10(11): e0139689. PMID 26545096
- Goel A, Shivaprasad C, Kolly A, et al. Comparison of electrochemical skin conductance and vibration perception threshold measurement in the detection of early diabetic neuropathy. *PLoS One*. 2017; 12(9): e0183973. PMID 28880907
- Azzopardi K, Gatt A, Chockalingam N, et al. Hidden dangers revealed by misdiagnosed diabetic neuropathy: A comparison of simple clinical tests for the screening of vibration perception threshold at primary care level. *Prim Care Diabetes*. Apr 2018; 12(2): 111-115. PMID 29029862
- Papanas N, Pafili K, Demetriou M, et al. The Diagnostic Utility of VibraTip for Distal Symmetrical Polyneuropathy in Type 2 Diabetes Mellitus. *Diabetes Ther*. Jan 2020; 11(1): 341-346. PMID 31782049
- Ferdousi M, Kalteniece A, Azmi S, et al. Corneal confocal microscopy compared with quantitative sensory testing and nerve conduction for diagnosing and stratifying the severity of diabetic peripheral neuropathy. *BMJ Open Diabetes Res Care*. Dec 2020; 8(2). PMID 33355206
- Devigili G, Tugnoli V, Penza P, et al. The diagnostic criteria for small fibre neuropathy: from symptoms to neuropathology. *Brain*. Jul 2008; 131(Pt 7): 1912-25. PMID 18524793
- Lefaucheur JP, Wahab A, Plante-Bordeneuve V, et al. Diagnosis of small fiber neuropathy: A comparative study of five neurophysiological tests. *Neurophysiol Clin*. Dec 2015; 45(6): 445-55. PMID 26596193
- Anand P, Privitera R, Yiangou Y, et al. Trench Foot or Non-Freezing Cold Injury As a Painful Vaso-Neuropathy: Clinical and Skin Biopsy Assessments. *Front Neurol*. 2017; 8: 514. PMID 28993756
- Fabry V, Gerdelat A, Acket B, et al. Which Method for Diagnosing Small Fiber Neuropathy?. *Front Neurol*. 2020; 11: 342. PMID 32431663

20. Shy ME, Frohman EM, So YT, et al. Quantitative sensory testing: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. Mar 25 2003; 60(6): 898-904. PMID 12654951
21. American Academy of Neurology. Quantitative Sensory Testing. 2003 (reaffirmed 2022); <https://www.aan.com/Guidelines/home/GuidelineDetail/87>. Accessed April 27, 2022.
22. Chong PS, Cros DP. Technology literature review: quantitative sensory testing. *Muscle Nerve*. May 2004; 29(5): 734-47. PMID 15116380
23. England JD, Gronseth GS, Franklin G, et al. Distal symmetrical polyneuropathy: definition for clinical research. *Muscle Nerve*. Jan 2005; 31(1): 113-23. PMID 15536624
24. American Diabetes Association. 11. Microvascular Complications and Foot Care: Standards of Medical Care in Diabetes-2021. *Diabetes Care*. Jan 2021; 44(Suppl 1): S151-S167. PMID 33298422
25. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination (NCD) for sensory Nerve Conduction Threshold Tests (sNCTs) (160.23). 2004; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=270&ncdver=2>. Accessed April 27, 2022.

## POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
September 2012	New policy	Quantitative sensory testing, including but not limited to current perception threshold testing, pressure-specified sensory device testing, vibration perception threshold testing, and thermal threshold testing, is considered investigational.
December 2013	Replace policy	Policy updated with literature review, Policy statement unchanged. References added; other references reordered or removed.
December 2014	Replace policy	Policy updated with literature review. Policy statement unchanged. Reference 16 added.
March 2016	Replace policy	Policy updated with literature review Policy statement unchanged. Reference 14 added.
September 2018	Replace policy	Policy updated with literature review through May 1, 2018; references 9- 11, 13-14, and 16 added. Policy statement unchanged except "not medically necessary" corrected to "investigational" due to FDA 510k process.
September 2019	Replace policy	Policy updated with literature review through April 3, 2019, no references added. Policy statement unchanged.
December 2020	Replace policy	Policy updated with literature review through August 20, 2020; references added. Policy statement unchanged.
September 2021	Replace policy	Policy updated with literature review through April 19, 2021; references added. Policy statement unchanged.
September 2022	Replace policy	Policy updated with literature review through April 27, 2022; background references added. Policy statement unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.