



FEP Medical Policy Manual

FEP 2.01.56 Low Level Laser Therapy

Effective Policy Date: October 1, 2022

Original Policy Date: June 2012

Related Policies:

2.01.21 - Temporomandibular Joint Disorder

Low Level Laser Therapy

Description

Description

Low-level laser therapy (LLLT), also called photobiomodulation, is being evaluated to treat various conditions, including, among others, oral mucositis, myofascial pain, joint pain, lymphedema, and chronic wounds.

OBJECTIVE

The objective of this evidence review is to evaluate net health outcomes of low-level laser therapy for treating individuals at increased risk of mucositis and other conditions (eg, soft tissue injuries, myofascial pain, tendinopathies, nerve injuries, joint pain, lymphedema).

POLICY STATEMENT

Low-level laser therapy may be considered **medically necessary** for the prevention of oral mucositis in individuals undergoing cancer treatment associated with increased risk of oral mucositis, including chemotherapy and/or radiotherapy, and/or hematopoietic cell transplantation (see Policy Guidelines).

Low-level laser therapy is considered **investigational** for all other indications including but not limited to:

- Carpal tunnel syndrome
- Neck pain
- Subacromial impingement
- Adhesive capsulitis
- Temporomandibular joint pain
- Low back pain
- Osteoarthritic knee pain
- Heel pain (ie, Achilles tendinopathy, plantar fasciitis)
- Rheumatoid arthritis
- Bell palsy
- Fibromyalgia
- Wound healing
- Lymphedema.

POLICY GUIDELINES

In the meta-analysis of 18 trials comparing low-level laser therapy (LLL) to chemotherapy or chemoradiation for prevention of oral mucositis (Oberoi et al [2014]), the course of LLLT was generally from day 0 through treatment. In studies of hematopoietic cell transplant, the course of LLLT began between day -7 and day 0 and continued as long as day 14 or 15. In studies that began LLLT at day -7 or day -5 before hematopoietic cell transplant, the course of laser therapy ended at day -1 or day 0.

Other protocols have applied low-level laser energy to acupuncture points on the fingers and hand. This technique may be referred to as *laser acupuncture*. Laser acupuncture is not reviewed herein.

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

MicroLight Laser Corp. has published a list of providers offering low-level laser therapy; most of the providers are chiropractors. Because the therapy typically requires up to 15 treatments, contractual or benefit restrictions on chiropractic visits for an individual diagnosis may apply.

FDA REGULATORY STATUS

Table 1. Selected Low-Level Laser Therapy Devices Cleared by the U.S. Food and Drug Administration

Device	Manufacturer	Date Cleared	510(k) No.	Indication
FX-635	Erchonia Corporation	6/01/2019	K190572	For adjunctive use in whole body musculoskeletal pain therapy
Super Pulsed Laser Technology	Multi Radiance Medical	01/13/2018	K171354	Providing temporary relief of minor chronic neck and shoulder pain of musculoskeletal origin
Lightstream Low-Level Laser	SOLICA CORPORATION	04/03/2009	K081166	For adjunctive use in the temporary relief of pain associated with knee disorders with standard chiropractic practice
GRT LITE, MODEL 8-A	GRT SOLUTIONS, INC.	02/03/2006	K050668	Use in providing temporary relief of minor chronic neck and shoulder pain of musculoskeletal origin
MICROLIGHT 830 LASER SYSTEM	MICROLIGHT CORPORATION OF AMERICA	02/06/2002	K010175	Use in pain therapy or related indication

A number of low-level lasers have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process for the treatment of pain. Data submitted for the MicroLight 830 Laser consisted of the application of the laser over the carpal tunnel 3 times a week for 5 weeks. The labeling states that the "MicroLight 830 Laser is indicated for adjunctive use in the temporary relief of hand and wrist pain associated with Carpal Tunnel Syndrome." In 2006, GRT LITE™ was cleared for marketing, listing the TUCO Erchonia PL3000, the Excalibur System, the MicroLight 830 Laser, and the Acculaser Pro as predicate devices. Indications of the GRTLITE™ for CTS are similar to the predicate devices: "adjunctive use in providing temporary relief of minor chronic pain." In 2009, the LightStream™ LLL device was cleared for marketing by the FDA through the 510(k) process for adjunctive use in the temporary relief of pain associated with knee disorders treated in standard chiropractic practice. A number of clinical trials of LLLT are underway in the U.S., including studies of wound healing. Since 2009, many more similar LLLT devices have received 510(k) clearance from the FDA.

RATIONALE

Summary of Evidence

Oral Mucositis

For individuals who have an increased risk of oral mucositis due to some cancer treatments (eg, chemotherapy, radiotherapy) and/or hematopoietic cell transplantation (HCT) who receive low-level laser therapy (LLLT), the evidence includes systematic reviews. Relevant outcomes are symptoms, morbid events, quality of life (QOL), and treatment-related morbidity. Several systematic reviews of randomized controlled trials (RCTs) have found better outcomes with LLLT used to prevent oral mucositis than with control treatments. Results have consistently supported a reduction in severe oral mucositis in patients undergoing chemotherapy, HCT, radiotherapy, and chemoradiotherapy. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Musculoskeletal and Neurologic Disorders

For individuals who have carpal tunnel syndrome (CTS) who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. Both a 2016 systematic review and a TEC Assessment (2010) did not find sufficient evidence from RCTs that LLLT improves outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have neck pain who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. A 2013 systematic review identified 17 trials, most of which were considered low-quality. Only 2 trials were considered moderate quality, and they found that LLLT led to better outcomes than placebo for chronic neck pain. A TEC Assessment (2010) found conflicting evidence. Additionally, laser types, application dosages, and treatment schedules vary in the available evidence and require further study. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have subacromial impingement syndrome who receive LLLT, the evidence includes RCTs. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. Most trials did not show a significant benefit of LLLT compared with sham treatment or with an alternative intervention (eg, exercise). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have adhesive capsulitis who receive LLLT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. A Cochrane review evaluating treatments for adhesive capsulitis identified 2 RCTs assessing LLLT. Due to the small number of trials and study limitations, reviewers concluded that the evidence was insufficient to permit conclusions about the effectiveness of LLLT for adhesive capsulitis. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have temporomandibular joint (TMJ) pain who receive LLLT, the evidence includes RCTs and several systematic reviews. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. Meta-analyses of RCTs had mixed findings. A 2021 meta-analysis, which included 33 placebo-controlled randomized trials, found a statistically significant impact of LLLT on pain scores and improved functional outcomes (eg, mouth opening); however, heterogeneity was high among included trials. Furthermore, RCTs have not compared the impact of LLLT with physical therapy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have low back pain who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. Meta-analyses of RCTs found that LLLT resulted in a significantly greater reduction in pain scores and global assessment scores than a placebo control in the immediate posttreatment setting. Meta-analyses also found that other outcomes (eg, disability index, range of motion) were significantly better immediately after treatment with active rather than placebo LLLT but not at longer-term follow-up. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have osteoarthritis (OA) knee pain who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. A 2020 systematic review, which pooled study findings, did find that LLLT significantly reduced pain or improved functional outcomes compared with a sham intervention; however, the study was limited by high heterogeneity and inconsistency between regimens and follow-up duration. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heel pain (ie, Achilles tendinopathy, plantar fasciitis) who receive LLLT, the evidence includes RCTs and 2 systematic reviews. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. Findings of sham-controlled randomized trials were inconsistent, and RCTs lacked long term follow up. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have rheumatoid arthritis (RA) who receive LLLT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. A systematic review of RCTs found an inconsistent benefit of LLLT for a range of outcomes. A 2010 RCT, published after the systematic review, did not find that LLLT was significantly better than a placebo treatment on most outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have Bell palsy who receive LLLT, the evidence includes a systematic review of 4 RCTs, with 2 RCTs reported in English. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. One RCT found a significant short-term benefit of LLLT over exercise. Longer-term outcomes (>6 weeks) were not available. Because Bell palsy often improves within weeks and may completely resolve within months, it is difficult to isolate specific improvements from laser therapy over the natural resolution of the illness. Also, no sham-controlled trials are available. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have fibromyalgia who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. The RCTs evaluating LLLT for treatment of fibromyalgia are small (ie, <25 patients each). One RCT (N=20 patients) found significantly better outcomes with LLLT than with sham, while another (N=20 patients) did not find statistically

significant between-group differences for similar outcomes. Additional RCTs with sufficient numbers of patients are needed to establish the efficacy of LLLT for fibromyalgia. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Wound Care and Lymphedema

For individuals who have chronic nonhealing wounds who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. The few existing RCTs tend to have small sample sizes and potential risk of bias. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have lymphedema who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. Multiple systematic reviews detected methodologic flaws in the available studies and did not consistently find better outcomes for patients receiving LLLT than those receiving a control condition for the treatment of lymphedema. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

Multinational Association of Supportive Care in Cancer and International Society of Oral Oncology

In 2017, the Mucositis Prevention Guideline Development Group published guidelines on preventing oral and oropharyngeal mucositis in children undergoing hematopoietic cell transplantation.¹⁴⁷ The guidelines were based on an evidence review consisting of randomized controlled trials that evaluated interventions such as cryotherapy and low-level laser therapy (LLLT). The guidelines suggested that LLLT could be offered to children but classified this recommendation as weak.

In 2020, the Multinational Association of Supportive Care in Cancer and the International Society of Oral Oncology published joint guidelines on the management of mucositis secondary to cancer therapy.¹⁴⁸

For the prevention of oral mucositis, the 2 associations recommended the following treatments, based on level 1 evidence: LLLT in patients undergoing radiotherapy with chemotherapy for head and neck cancer; LLLT in patients receiving hematopoietic cell transplantation conditioned with high-dose chemotherapy with or without total body irradiation; recombinant human keratinocyte growth factor-1 in patients receiving high-dose chemotherapy and total body irradiation, followed by autologous cell transplantation for hematologic malignancy; and benzydamine mouthwash in patients with head and neck cancer receiving moderate-dose radiotherapy without concomitant chemotherapy.

Additionally, numerous treatments were recommended for the prevention of oral mucositis based on level II evidence, including LLLT in patients undergoing radiotherapy, without concomitant chemotherapy, for head and neck cancer. Several LLLT protocols are outlined by the guideline based on cancer treatment modality, ranging in wavelength from 632.9 to 660 nm.

American Physical Therapy Association

In 2018, the American Physical Therapy Association published an updated guideline on the diagnosis and treatment of Achilles tendinitis.¹⁴⁹ The use of LLLT was given a level D recommendation, meaning that no recommendation could be made due to contradictory evidence. This is a change from the previous version of the guideline published in 2010, which gave LLLT a level B recommendation.¹⁵⁰

National Institute for Health and Care Excellence

In 2009, NICE issued guidance on early management of persistent, nonspecific low back pain and did not recommend laser treatment, citing limited evidence.¹⁵¹ The 2016 and 2020 updated guidance does not mention laser therapy.¹⁵¹

North American Spine Society

In 2020, the North American Spine Society published a guideline on the diagnosis and treatment of low back pain.¹⁵² The guideline was based on a systematic review of the literature to address key clinical questions regarding the diagnosis and treatment of adults with nonspecific low back pain.

Table 2. North American Spine Society Guideline Recommendations for Laser Therapy

Guideline Recommendation	Grade of Recommendation
"It is suggested that the combination of laser therapy (low-level or high-level) with exercise provides better short-term relief of pain than either exercise or laser therapy alone."	B
"There is conflicting evidence that the combination of laser therapy with exercise provides better short-term improvement in function compared to exercise or laser therapy alone."	I
"It is suggested that there is no short-term benefit of laser therapy (low-level or high-level) when compared with exercise alone."	B

Grade of Recommendation (levels of evidence range from Level I [high quality randomized controlled trial] to Level V [expert consensus]): A=Good evidence (Level I studies with consistent findings) for or against recommending intervention; B=Fair evidence (Level II or III studies with consistent findings) for or against recommending intervention; C=Poor quality evidence (Level IV or V studies) for or against recommending intervention; I=Insufficient or conflicting evidence not allowing a recommendation for or against intervention.

American College of Physicians

In 2017, the American College of Physicians released guidelines relating to noninvasive treatments for chronic low back pain.¹⁵³ The guidelines strongly recommended that patients with chronic low back pain should first seek nonpharmacologic treatment such as exercise, multidisciplinary rehabilitation, acupuncture, and mindfulness-based stress reduction—all based on moderate quality evidence. The recommendation also stated that patients with chronic low back pain should seek treatments such as tai chi, yoga, motor control exercise, progressive relaxation, electromyography biofeedback, LLLT, operant therapy, cognitive behavioral therapy, or spinal manipulation—all based on low-quality evidence. While the College stated that LLLT has a small effect on pain and function, it found the evidence insufficient for the use of LLLT.

In 2020, the American College of Physicians published a joint guideline on management of acute pain from non-low back musculoskeletal injuries with the American Academy of Family Physicians.¹⁵⁴ No recommendations are made specific to LLLT, but the guideline notes that laser therapy did not significantly reduce pain in 1 to 7 days compared to placebo.

American Academy of Orthopaedic Surgeons

In 2016, the American Academy of Orthopaedic Surgeons' guidelines on the management of carpal tunnel syndrome indicated the: "limited evidence supports that laser therapy might be effective compared to placebo."¹⁵⁵

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

REFERENCES

1. Lalla RV, Bowen J, Barasch A, et al. MASCC/ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. *Cancer*. May 15 2014; 120(10): 1453-61. PMID 24615748
2. Schubert MM, Eduardo FP, Guthrie KA, et al. A phase III randomized double-blind placebo-controlled clinical trial to determine the efficacy of low level laser therapy for the prevention of oral mucositis in patients undergoing hematopoietic cell transplantation. *Support Care Cancer*. Oct 2007; 15(10): 1145-54. PMID 17393191
3. Figueiredo AL, Lins L, Cattony AC, et al. Laser therapy in the control of oral mucositis: a meta-analysis. *Rev Assoc Med Bras* (1992). Sep-Oct 2013; 59(5): 467-74. PMID 24119379
4. Doeuk C, Hersant B, Bosc R, et al. Current indications for low level laser treatment in maxillofacial surgery: a review. *Br J Oral Maxillofac Surg*. Apr 2015; 53(4): 309-15. PMID 25740083
5. Peng J, Shi Y, Wang J, et al. Low-level laser therapy in the prevention and treatment of oral mucositis: a systematic review and meta-analysis. *Oral Surg Oral Med Oral Pathol Oral Radiol*. Oct 2020; 130(4): 387-397.e9. PMID 32624448
6. Oberoi S, Zamperini-Netto G, Beyene J, et al. Effect of prophylactic low level laser therapy on oral mucositis: a systematic review and meta-analysis. *PLoS One*. 2014; 9(9): e107418. PMID 25198431
7. Rankin IA, Sargeant H, Rehman H, et al. Low-level laser therapy for carpal tunnel syndrome. *Cochrane*. <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012765/full> Published 22 August 2017. Accessed April 25, 2022.
8. Li ZJ, Wang Y, Zhang HF, et al. Effectiveness of low-level laser on carpal tunnel syndrome: A meta-analysis of previously reported randomized trials. *Medicine* (Baltimore). Aug 2016; 95(31): e4424. PMID 27495063
9. Fusakul Y, Aranyavalai T, Saensri P, et al. Low-level laser therapy with a wrist splint to treat carpal tunnel syndrome: a double-blinded randomized controlled trial. *Lasers Med Sci*. May 2014; 29(3): 1279-87. PMID 24477392
10. Low-level laser therapy for carpal tunnel syndrome and chronic neck pain. *Technol Eval Cent Assess Program Exec Summ*. Nov 2010; 25(4): 1-2. PMID 21638940
11. Chow RT, Heller GZ, Barnsley L. The effect of 300 mW, 830 nm laser on chronic neck pain: a double-blind, randomized, placebo-controlled study. *Pain*. Sep 2006; 124(1-2): 201-10. PMID 16806710
12. Gross AR, Dziengo S, Boers O, et al. Low Level Laser Therapy (LLLT) for Neck Pain: A Systematic Review and Meta-Regression. *Open Orthop J*. 2013; 7: 396-419. PMID 24155802
13. Yeldan I, Cetin E, Ozdincler AR. The effectiveness of low-level laser therapy on shoulder function in subacromial impingement syndrome. *Disabil Rehabil*. 2009; 31(11): 935-40. PMID 19031167
14. Dogan SK, Ay S, Evcik D. The effectiveness of low laser therapy in subacromial impingement syndrome: a randomized placebo controlled double-blind prospective study. *Clinics (Sao Paulo)*. 2010; 65(10): 1019-22. PMID 21120304
15. Abrisham SM, Kermani-Alghoraishi M, Ghahramani R, et al. Additive effects of low-level laser therapy with exercise on subacromial syndrome: a randomised, double-blind, controlled trial. *Clin Rheumatol*. Oct 2011; 30(10): 1341-6. PMID 21538218
16. Bal A, Eksioğlu E, Gurcay E, et al. Low-level laser therapy in subacromial impingement syndrome. *Photomed Laser Surg*. Feb 2009; 27(1): 31-6. PMID 19250050
17. Calis HT, Berberoglu N, Calis M. Are ultrasound, laser and exercise superior to each other in the treatment of subacromial impingement syndrome? A randomized clinical trial. *Eur J Phys Rehabil Med*. Mar 2011; 47(3): 375-380. PMID 21364511
18. Alfredo PP, Bjordal JM, Junior WS, et al. Efficacy of low-level laser therapy combined with exercise for subacromial impingement syndrome: A randomised controlled trial. *Clin Rehabil*. Jun 2021; 35(6): 851-860. PMID 33307783
19. Badil Guloglu S. Comparison of low-level laser treatment and extracorporeal shock wave therapy in subacromial impingement syndrome: a randomized, prospective clinical study. *Lasers Med Sci*. Jun 2021; 36(4): 773-781. PMID 32638239
20. Page MJ, Green S, Kramer S, et al. Electrotherapy modalities for adhesive capsulitis (frozen shoulder). *Cochrane Database Syst Rev*. Oct 01 2014; (10): CD011324. PMID 25271097
21. Stergioulas A, Stergioula M, Aarskog R, et al. Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic achilles tendinopathy. *Am J Sports Med*. May 2008; 36(5): 881-7. PMID 18272794
22. Chen J, Huang Z, Ge M, et al. Efficacy of low-level laser therapy in the treatment of TMDs: a meta-analysis of 14 randomised controlled trials. *J Oral Rehabil*. Apr 2015; 42(4): 291-9. PMID 25491183
23. Chang WD, Lee CL, Lin HY, et al. A Meta-analysis of Clinical Effects of Low-level Laser Therapy on Temporomandibular Joint Pain. *J Phys Ther Sci*. Aug 2014; 26(8): 1297-300. PMID 25202201
24. Hanna R, Dalvi S, Bensadoun RJ, et al. Role of Photobiomodulation Therapy in Modulating Oxidative Stress in Temporomandibular Disorders. A Systematic Review and Meta-Analysis of Human Randomised Controlled Trials. *Antioxidants* (Basel). Jun 25 2021; 10(7). PMID 34202292
25. Conti PC. Low level laser therapy in the treatment of temporomandibular disorders (TMD): a double-blind pilot study. *Cranio*. Apr 1997; 15(2): 144-9. PMID 9586517
26. Kulekcioglu S, Sivrioglu K, Ozcan O, et al. Effectiveness of low-level laser therapy in temporomandibular disorder. *Scand J Rheumatol*. 2003; 32(2): 114-8. PMID 12737331
27. Venancio Rde A, Camparis CM, Lizarelli Rde F. Low intensity laser therapy in the treatment of temporomandibular disorders: a double-blind study. *J Oral Rehabil*. Nov 2005; 32(11): 800-7. PMID 16202043
28. Cetiner S, Kahraman SA, Yucetas S. Evaluation of low-level laser therapy in the treatment of temporomandibular disorders. *Photomed Laser Surg*. Oct 2006; 24(5): 637-41. PMID 17069496
29. Fikackova H, Dostalova T, Navratil L, et al. Effectiveness of low-level laser therapy in temporomandibular joint disorders: a placebo-controlled study. *Photomed Laser Surg*. Aug 2007; 25(4): 297-303. PMID 17803388

30. Mazzetto MO, Carrasco TG, Bidinelo EF, et al. Low intensity laser application in temporomandibular disorders: a phase I double-blind study. *Cranio*. Jul 2007; 25(3): 186-92. PMID 17696035
31. Frare J.C., Nicolau R.A. Clinical analysis of the effect of laser photobiomodulation (GaAs904 nm) on temporomandibular joint dysfunction. *Rev. Bras. Fisioter.* 2008;12:3742. doi: 10.1590/S1413-35552008000100008.
32. da Cunha LA, Firoozmand LM, da Silva AP, et al. Efficacy of low-level laser therapy in the treatment of temporomandibular disorder. *Int Dent J*. Aug 2008; 58(4): 213-7. PMID 18783114
33. Lassemi E., Jafari S.M., Motamedi M.H.K., Navi F., Lasemi R. Low- level laser therapy in the management of temporomandibular joint disorder. *J. Oral Laser Appl.* 2008;8:8386
34. Carrasco TG, Mazzetto MO, Mazzetto RG, et al. Low intensity laser therapy in temporomandibular disorder: a phase II double-blind study. *Cranio*. Oct 2008; 26(4): 274-81. PMID 19004308
35. Emshoff R, Bosch R, Pempel E, et al. Low-level laser therapy for treatment of temporomandibular joint pain: a double-blind and placebo-controlled trial. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* Apr 2008; 105(4): 452-6. PMID 18329580
36. Carrasco TG, Guerisoli LD, Guerisoli DM, et al. Evaluation of low intensity laser therapy in myofascial pain syndrome. *Cranio*. Oct 2009; 27(4): 243-7. PMID 19891258
37. Shirani AM, Gutknecht N, Taghizadeh M, et al. Low-level laser therapy and myofascial pain dysfunction syndrome: a randomized controlled clinical trial. *Lasers Med Sci.* Sep 2009; 24(5): 715-20. PMID 19002646
38. Venezian GC, da Silva MA, Mazzetto RG, et al. Low level laser effects on pain to palpation and electromyographic activity in TMD patients: a double-blind, randomized, placebo-controlled study. *Cranio*. Apr 2010; 28(2): 84-91. PMID 20491229
39. Oz S, Gokcen-Rohlig B, Saruhanoglu A, et al. Management of myofascial pain: low-level laser therapy versus occlusal splints. *J Craniofac Surg.* Nov 2010; 21(6): 1722-8. PMID 21119408
40. Marini I, Gatto MR, Bonetti GA. Effects of superpulsed low-level laser therapy on temporomandibular joint pain. *Clin J Pain.* Sep 2010; 26(7): 611-6. PMID 20664343
41. Rohlig B.G., Kipirdi S., Meric U., Capan N., Keskin H. Masticatory muscle pain and low-level laser therapy: A double-blind and placebo-controlled study. *Turk. J. Phys. Med. Rehabil. Turk. Fiz. Tip Rehabil. Derg.* 2011;57:3137. doi: 10.4274/tftr.57.06.
42. Sattayut S, Bradley P. A study of the influence of low intensity laser therapy on painful temporomandibular disorder patients. *Laser Ther.* Sep 30 2012; 21(3): 183-92. PMID 24511188
43. de Carli ML, Guerra MB, Nunes TB, et al. Piroxicam and laser phototherapy in the treatment of TMJ arthralgia: a double-blind randomised controlled trial. *J Oral Rehabil.* Mar 2013; 40(3): 171-8. PMID 23252583
44. da Silva MA, Botelho AL, Turim CV, et al. Low level laser therapy as an adjunctive technique in the management of temporomandibular disorders. *Cranio*. Oct 2012; 30(4): 264-71. PMID 23156967
45. Panhoca VH, Lizarelli Rde F, Nunez SC, et al. Comparative clinical study of light analgesic effect on temporomandibular disorder (TMD) using red and infrared led therapy. *Lasers Med Sci.* Feb 2015; 30(2): 815-22. PMID 24197518
46. Uemoto L, Garcia MA, Gouvea CV, et al. Laser therapy and needling in myofascial trigger point deactivation. *J Oral Sci.* 2013; 55(2): 175-81. PMID 23748458
47. Ferreira LA, de Oliveira RG, Guimaraes JP, et al. Laser acupuncture in patients with temporomandibular dysfunction: a randomized controlled trial. *Lasers Med Sci.* Nov 2013; 28(6): 1549-58. PMID 23380907
48. Demirkol N, Sari F, Bulbul M, et al. Effectiveness of occlusal splints and low-level laser therapy on myofascial pain. *Lasers Med Sci.* Apr 2015; 30(3): 1007-12. PMID 24504660
49. Ahrari F, Madani AS, Ghafouri ZS, et al. The efficacy of low-level laser therapy for the treatment of myogenous temporomandibular joint disorder. *Lasers Med Sci.* Mar 2014; 29(2): 551-7. PMID 23318917
50. Pereira TS, Flecha OD, Guimaraes RC, et al. Efficacy of red and infrared lasers in treatment of temporomandibular disorders--a double-blind, randomized, parallel clinical trial. *Cranio*. Jan 2014; 32(1): 51-6. PMID 24660647
51. de Moraes Maia ML, Ribeiro MA, Maia LG, et al. Evaluation of low-level laser therapy effectiveness on the pain and masticatory performance of patients with myofascial pain. *Lasers Med Sci.* Jan 2014; 29(1): 29-35. PMID 23143142
52. Sancakli E, Gokcen-Rohlig B, Balik A, et al. Early results of low-level laser application for masticatory muscle pain: a double-blind randomized clinical study. *BMC Oral Health.* Oct 23 2015; 15(1): 131. PMID 26496720
53. Douglas De Oliveira DW, Lages FS, Guimaraes RC, et al. Do TMJ symptoms improve and last across time after treatment with red (660 nm) and infrared (790 nm) low level laser treatment (LLLT)? A survival analysis. *Cranio*. Nov 2017; 35(6): 372-378. PMID 28218006
54. Costa SAP, Florezi GP, Artes GE, et al. The analgesic effect of photobiomodulation therapy (830 nm) on the masticatory muscles: a randomized, double-blind study. *Braz Oral Res.* Dec 18 2017; 31: e107. PMID 29267668
55. Seifi M, Ebadifar A, Kabiri S, et al. Comparative effectiveness of Low Level Laser therapy and Transcutaneous Electric Nerve Stimulation on Temporomandibular Joint Disorders. *J Lasers Med Sci.* 2017; 8(Suppl 1): S27-S31. PMID 29071032
56. Shobha R, Narayanan VS, Jagadish Pai BS, et al. Low-level laser therapy: A novel therapeutic approach to temporomandibular disorder - A randomized, double-blinded, placebo-controlled trial. *Indian J Dent Res.* Jul-Aug 2017; 28(4): 380-387. PMID 28836528
57. Rezazadeh F, Hajian K, Shahidi S, et al. Comparison of the Effects of Transcutaneous Electrical Nerve Stimulation and Low-Level Laser Therapy on Drug-Resistant Temporomandibular Disorders. *J Dent (Shiraz).* Sep 2017; 18(3): 187-192. PMID 29034273
58. Varma S.R., al Shayeb M., el Kaseh A., Kuduruthullah S., Ashekhi A., al Khader E. Effectiveness of low-level laser therapy in the Management of the Temporomandibular Joint Disorders: A Placebo-controlled Trial. *World J. Dent.* 2018;9:316320. doi: 10.5005/jp-journals-10015-1555.
59. Borges RMM, Cardoso DS, Flores BC, et al. Effects of different photobiomodulation dosimetries on temporomandibular dysfunction: a randomized, double-blind, placebo-controlled clinical trial. *Lasers Med Sci.* Dec 2018; 33(9): 1859-1866. PMID 29850961
60. Brochado FT, Jesus LH, Carrard VC, et al. Comparative effectiveness of photobiomodulation and manual therapy alone or combined in TMD patients: a randomized clinical trial. *Braz Oral Res.* Jul 10 2018; 32: e50. PMID 29995062

61. Rodrigues CA, Melchior MO, Valencise Magri L, et al. Can the severity of orofacial myofunctional conditions interfere with the response of analgesia promoted by active or placebo low-level laser therapy?. *Cranio*. Jul 2020; 38(4): 240-247. PMID 30244669
62. Peimani A., Keshavarz S., Fathollahi M.S. Comparison of Low-Level Laser Therapy and Drug Therapy in Patients with Temporomandibular Disorders: A Randomized Clinical Trial. *J. Oral Health Dent*. 2020;38:240247. doi: 10.1080/08869634.2018.1520950.
63. Nadershah M, Abdel-Alim HM, Bayoumi AM, et al. Photobiomodulation Therapy for Myofascial Pain in Temporomandibular Joint Dysfunction: A Double-Blinded Randomized Clinical Trial. *J Maxillofac Oral Surg*. Mar 2020; 19(1): 93-97. PMID 31988570
64. Magri LV, Bataglion C, Leite-Panissi CRA. Follow-up results of a randomized clinical trial for low-level laser therapy in painful TMD of muscular origins. *Cranio*. Nov 2021; 39(6): 502-509. PMID 31585522
65. Al-Quisi AF, Al-Anee AM, Al-Jumaily HA, et al. Efficacy of the LED Red Light Therapy in the Treatment of Temporomandibular Disorders: Double Blind Randomized Controlled Trial. *Pain Res Treat*. 2019; 2019: 8578703. PMID 31205787
66. Herpich CM, Leal-Junior ECP, Politti F, et al. Intraoral photobiomodulation diminishes pain and improves functioning in women with temporomandibular disorder: a randomized, sham-controlled, double-blind clinical trial : Intraoral photobiomodulation diminishes pain in women with temporomandibular disorder. *Lasers Med Sci*. Mar 2020; 35(2): 439-445. PMID 31325122
67. Khairnar S, Bhate K, S N SK, et al. Comparative evaluation of low-level laser therapy and ultrasound heat therapy in reducing temporomandibular joint disorder pain. *J Dent Anesth Pain Med*. Oct 2019; 19(5): 289-294. PMID 31723669
68. Sobral APT, Godoy CLH, Fernandes KPS, et al. Photomodulation in the treatment of chronic pain in patients with temporomandibular disorder: protocol for cost-effectiveness analysis. *BMJ Open*. May 05 2018; 8(5): e018326. PMID 29730613
69. Maracci LM, Stasiak G, de Oliveira Chami V, et al. Treatment of myofascial pain with a rapid laser therapy protocol compared to occlusal splint: A double-blind, randomized clinical trial. *Cranio*. Jun 03 2020: 1-7. PMID 32491964
70. Chellappa D, Thirupathy M. Comparative efficacy of low-Level laser and TENS in the symptomatic relief of temporomandibular joint disorders: A randomized clinical trial. *Indian J Dent Res*. Jan-Feb 2020; 31(1): 42-47. PMID 32246680
71. Monteiro L, Ferreira R, Resende T, et al. Effectiveness of Photobiomodulation in Temporomandibular Disorder-Related Pain Using a 635 nm Diode Laser: A Randomized, Blinded, and Placebo-Controlled Clinical Trial. *Photobiomodul Photomed Laser Surg*. May 2020; 38(5): 280-288. PMID 32427553
72. Del Vecchio A, Floravanti M, Boccassini A, et al. Evaluation of the efficacy of a new low-level laser therapy home protocol in the treatment of temporomandibular joint disorder-related pain: A randomized, double-blind, placebo-controlled clinical trial. *Cranio*. Mar 2021; 39(2): 141-150. PMID 30999823
73. Aisaiti A, Zhou Y, Wen Y, et al. Effect of photobiomodulation therapy on painful temporomandibular disorders. *Sci Rep*. Apr 27 2021; 11(1): 9049. PMID 33907210
74. Glazov G, Yelland M, Emery J. Low-level laser therapy for chronic non-specific low back pain: a meta-analysis of randomised controlled trials. *Acupunct Med*. Oct 2016; 34(5): 328-341. PMID 27207675
75. Huang Z, Ma J, Chen J, et al. The effectiveness of low-level laser therapy for nonspecific chronic low back pain: a systematic review and meta-analysis. *Arthritis Res Ther*. Dec 15 2015; 17: 360. PMID 26667480
76. Chen YJ, Liao CD, Hong JP, et al. Effects of laser therapy on chronic low back pain: A systematic review and meta-analysis of randomized controlled trials. *Clin Rehabil*. Mar 2022; 36(3): 289-302. PMID 34757882
77. Alayat MS, Atya AM, Ali MM, et al. Long-term effect of high-intensity laser therapy in the treatment of patients with chronic low back pain: a randomized blinded placebo-controlled trial. *Lasers Med Sci*. May 2014; 29(3): 1065-73. PMID 24178907
78. Ay S, Dogan SK, Evcik D. Is low-level laser therapy effective in acute or chronic low back pain?. *Clin Rheumatol*. Aug 2010; 29(8): 905-10. PMID 20414695
79. Basford JR, Sheffield CG, Harmsen WS. Laser therapy: a randomized, controlled trial of the effects of low-intensity Nd:YAG laser irradiation on musculoskeletal back pain. *Arch Phys Med Rehabil*. Jun 1999; 80(6): 647-52. PMID 10378490
80. Djavid GE, Mehrdad R, Ghasemi M, et al. In chronic low back pain, low level laser therapy combined with exercise is more beneficial than exercise alone in the long term: a randomised trial. *Aust J Physiother*. 2007; 53(3): 155-60. PMID 17725472
81. Glazov G, Schattner P, Lopez D, et al. Laser acupuncture for chronic non-specific low back pain: a controlled clinical trial. *Acupunct Med*. Sep 2009; 27(3): 94-100. PMID 19734378
82. Glazov G, Yelland M, Emery J. Low-dose laser acupuncture for non-specific chronic low back pain: a double-blind randomised controlled trial. *Acupunct Med*. Apr 2014; 32(2): 116-23. PMID 24280948
83. Klein RG, Eek BC. Low-energy laser treatment and exercise for chronic low back pain: double-blind controlled trial. *Arch Phys Med Rehabil*. Jan 1990; 71(1): 34-7. PMID 2136991
84. Konstantinovic L, Lazovic M, Milovanovic N, et al.. Low level laser therapy in geriatric patients with low back pain. *Eur J Pain Suppl (Poster Sessions)* 2011;5:61 10.1016/S1754-3207(11)70205-X
85. Lin ML, Wu HC, Hsieh YH, et al. Evaluation of the effect of laser acupuncture and cupping with ryodoraku and visual analog scale on low back pain. *Evid Based Complement Alternat Med*. 2012; 2012: 521612. PMID 23118792
86. Okamoto H. Therapeutic effect of semiconductor laser irradiation on low-back pain. *J Jpn A Phys Med Balneology Climatology* 1989;52:13145.
87. Ruth M, Weber M, Zenz M. [Laser acupuncture for chronic back pain. A double-blind clinical study]. *Schmerz*. Sep 2010; 24(5): 485-93. PMID 20872127
88. Soriano F, Rios R. Gallium arsenide laser treatment of chronic low back pain: a prospective, randomized and double blind study. *Laser Ther* 1998;10:17580. 10.5978/islsm.10.175
89. Umegaki S. Effectiveness of low-power laser therapy on low-back pain: double blind comparative study to evaluate the analgesic effect of low-power laser therapy. *Kiso to Rinsho (The Clinical Report)* 1989;23:283946.
90. Vallone F, Benedicenti S, Sorrenti E, et al. Effect of diode laser in the treatment of patients with nonspecific chronic low back pain: a randomized controlled trial. *Photomed Laser Surg*. Sep 2014; 32(9): 490-4. PMID 25141218

91. Wallace G. The effects of laser acupuncture on chronic low back pain [Thesis]. Melbourne Monash University, 1996.
92. Gur A, Karakoc M, Cevik R, et al. Efficacy of low power laser therapy and exercise on pain and functions in chronic low back pain. *Lasers Surg Med.* 2003; 32(3): 233-8. PMID 12605431
93. Hsieh RL, Lee WC. Short-term therapeutic effects of 890-nanometer light therapy for chronic low back pain: a double-blind randomized placebo-controlled study. *Lasers Med Sci.* Mar 2014; 29(2): 671-9. PMID 23820974
94. de Carvalho ME, de Carvalho RM, Marques AP, et al. Low intensity laser and LED therapies associated with lateral decubitus position and flexion exercises of the lower limbs in patients with lumbar disk herniation: clinical randomized trial. *Lasers Med Sci.* Sep 2016; 31(7): 1455-63. PMID 27379776
95. Tantawy SA, Abdelbasset WK, Kamel DM, et al. Laser photobiomodulation is more effective than ultrasound therapy in patients with chronic nonspecific low back pain: a comparative study. *Lasers Med Sci.* Jun 2019; 34(4): 793-800. PMID 30334124
96. Nambi G, Kamal W, Es S, et al. Spinal manipulation plus laser therapy versus laser therapy alone in the treatment of chronic non-specific low back pain: a randomized controlled study. *Eur J Phys Rehabil Med.* Dec 2018; 54(6): 880-889. PMID 29687966
97. Shin JY, Ku B, Kim JU, et al. Short-Term Effect of Laser Acupuncture on Lower Back Pain: A Randomized, Placebo-Controlled, Double-Blind Trial. *Evid Based Complement Alternat Med.* 2015; 2015: 808425. PMID 26516333
98. Koldas Dogan S, Ay S, Evcik D. The effects of two different low level laser therapies in the treatment of patients with chronic low back pain: A double-blinded randomized clinical trial. *J Back Musculoskelet Rehabil.* 2017; 30(2): 235-240. PMID 27472858
99. Huang Z, Chen J, Ma J, et al. Effectiveness of low-level laser therapy in patients with knee osteoarthritis: a systematic review and meta-analysis. *Osteoarthritis Cartilage.* Sep 2015; 23(9): 1437-1444. PMID 25914044
100. Bjordal JM, Johnson MI, Lopes-Martins RA, et al. Short-term efficacy of physical interventions in osteoarthritic knee pain. A systematic review and meta-analysis of randomised placebo-controlled trials. *BMC Musculoskelet Disord.* Jun 22 2007; 8: 51. PMID 17587446
101. Stausholm MB, Naterstad IF, Joensen J, et al. Efficacy of low-level laser therapy on pain and disability in knee osteoarthritis: systematic review and meta-analysis of randomised placebo-controlled trials. *BMJ Open.* Oct 28 2019; 9(10): e031142. PMID 31662383
102. Al Rashoud AS, Abboud RJ, Wang W, et al. Efficacy of low-level laser therapy applied at acupuncture points in knee osteoarthritis: a randomised double-blind comparative trial. *Physiotherapy.* Sep 2014; 100(3): 242-8. PMID 24418801
103. Alfredo PP, Bjordal JM, Dreyer SH, et al. Efficacy of low level laser therapy associated with exercises in knee osteoarthritis: a randomized double-blind study. *Clin Rehabil.* Jun 2012; 26(6): 523-33. PMID 22169831
104. Alfredo PP, Bjordal JM, Junior WS, et al. Long-term results of a randomized, controlled, double-blind study of low-level laser therapy before exercises in knee osteoarthritis: laser and exercises in knee osteoarthritis. *Clin Rehabil.* Feb 2018; 32(2): 173-178. PMID 28776408
105. Alghadir A, Omar MT, Al-Askar AB, et al. Effect of low-level laser therapy in patients with chronic knee osteoarthritis: a single-blinded randomized clinical study. *Lasers Med Sci.* Mar 2014; 29(2): 749-55. PMID 23912778
106. Bagheri SR, Fatemi E, Fazeli SH, et al. Efficacy of low level laser on knee osteoarthritis treatment [Persian]. *Koomesh* 2011;12:28592.
107. Bulow PM, Jensen H, Danneskiold-Samsøe B. Low power Ga-Al-As laser treatment of painful osteoarthritis of the knee. A double-blind placebo-controlled study. *Scand J Rehabil Med.* Sep 1994; 26(3): 155-9. PMID 7801065
108. Delkosh CT, Fatemy E, Ghorbani R, et al. Comparing the immediate and long-term effects of low and high power laser on the symptoms of knee osteoarthritis [Persian]. *Journal of mazandaran university of medical sciences* 2018;28:6977.
109. Fukuda VO, Fukuda TY, Guimaraes M, et al. SHORT-TERM EFFICACY OF LOW-LEVEL LASER THERAPY IN PATIENTS WITH KNEE OSTEOARTHRITIS: A RANDOMIZED PLACEBO-CONTROLLED, DOUBLE-BLIND CLINICAL TRIAL. *Rev Bras Ortop.* Sep-Oct 2011; 46(5): 526-33. PMID 27027049
110. Gur A, Cosut A, Sarac AJ, et al. Efficacy of different therapy regimes of low-power laser in painful osteoarthritis of the knee: a double-blind and randomized-controlled trial. *Lasers Surg Med.* 2003; 33(5): 330-8. PMID 14677160
111. Gworys K, Gaszytch J, Puzder A, et al. Influence of various laser therapy methods on knee joint pain and function in patients with knee osteoarthritis. *Ortop Traumatol Rehabil.* May-Jun 2012; 14(3): 269-77. PMID 22764339
112. Hegedus B, Viharos L, Gervain M, et al. The effect of low-level laser in knee osteoarthritis: a double-blind, randomized, placebo-controlled trial. *Photomed Laser Surg.* Aug 2009; 27(4): 577-84. PMID 19530911
113. Helianthi DR, Simadibrata C, Srilestari A, et al. Pain Reduction After Laser Acupuncture Treatment in Geriatric Patients with Knee Osteoarthritis: a Randomized Controlled Trial. *Acta Med Indones.* Apr 2016; 48(2): 114-21. PMID 27550880
114. Hinman RS, McCrory P, Pirodda M, et al. Acupuncture for chronic knee pain: a randomized clinical trial. *JAMA.* Oct 01 2014; 312(13): 1313-22. PMID 25268438
115. Jensen H, Harreby M, Kjer J. [Infrared laser--effect in painful arthrosis of the knee?]. *Ugeskr Laeger.* Nov 09 1987; 149(46): 3104-6. PMID 3445368
116. Kheshie AR, Alayat MS, Ali MM. High-intensity versus low-level laser therapy in the treatment of patients with knee osteoarthritis: a randomized controlled trial. *Lasers Med Sci.* Jul 2014; 29(4): 1371-6. PMID 24487957
117. Koutenaei FR, Mosallanezhad Z, Naghikhani M, et al. The effect of low level laser therapy on pain and range of motion of patients with knee osteoarthritis. *Physical Treatments - Specific Physical Therapy* 2017;7:1318.
118. Mohammed N, Allam H, Elghoroury E, et al. Evaluation of serum beta-endorphin and substance P in knee osteoarthritis patients treated by laser acupuncture. *J Complement Integr Med.* Jan 05 2018; 15(2). PMID 29303777
119. S GN, Kamal W, George J, et al. Radiological and biochemical effects (CTX-II, MMP-3, 8, and 13) of low-level laser therapy (LLLT) in chronic osteoarthritis in Al-Kharij, Saudi Arabia. *Lasers Med Sci.* Feb 2017; 32(2): 297-303. PMID 27913970
120. Nivbrant B, Friberg S. [Laser treatment of knee joint arthrosis seems to be effective but scientific evidence is lacking]. *Lakartidningen.* Mar 11 1992; 89(11): 859-61. PMID 1545640
121. Rayegani SM, Bahrami MH, Elyaspour D, et al. Therapeutic effects of low level laser therapy (LLLT) in knee osteoarthritis, compared to therapeutic ultrasound. *J Lasers Med Sci* 2012;3:7174.

122. Tascioglu F, Armagan O, Tabak Y, et al. Low power laser treatment in patients with knee osteoarthritis. *Swiss Med Wkly*. May 01 2004; 134(17-18): 254-8. PMID 15243853
123. Youssef EF, Muaidi QI, Shanb AA. Effect of Laser Therapy on Chronic Osteoarthritis of the Knee in Older Subjects. *J Lasers Med Sci*. 2016; 7(2): 112-9. PMID 27330707
124. Tumilty S, McDonough S, Hurley DA, et al. Clinical effectiveness of low-level laser therapy as an adjunct to eccentric exercise for the treatment of Achilles' tendinopathy: a randomized controlled trial. *Arch Phys Med Rehabil*. May 2012; 93(5): 733-9. PMID 22541305
125. Wang W, Jiang W, Tang C, et al. Clinical efficacy of low-level laser therapy in plantar fasciitis: A systematic review and meta-analysis. *Medicine (Baltimore)*. Jan 2019; 98(3): e14088. PMID 30653125
126. Guimaraes JS, Arcanjo FL, Leporace G, et al. Effect of low-level laser therapy on pain and disability in patients with plantar fasciitis: A systematic review and meta-analysis. *Musculoskelet Sci Pract*. Feb 2022; 57: 102478. PMID 34847470
127. Macias DM, Coughlin MJ, Zang K, et al. Low-Level Laser Therapy at 635 nm for Treatment of Chronic Plantar Fasciitis: A Placebo-Controlled, Randomized Study. *J Foot Ankle Surg*. Sep-Oct 2015; 54(5): 768-72. PMID 25769363
128. Kiritsi O, Tsitias K, Malliaropoulos N, et al. Ultrasonographic evaluation of plantar fasciitis after low-level laser therapy: results of a double-blind, randomized, placebo-controlled trial. *Lasers Med Sci*. Mar 2010; 25(2): 275-81. PMID 19841862
129. Cinar E, Saxena S, Uygur F. Low-level laser therapy in the management of plantar fasciitis: a randomized controlled trial. *Lasers Med Sci*. Jul 2018; 33(5): 949-958. PMID 29273892
130. Brosseau L, Robinson V, Wells G, et al. Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis. *Cochrane Database Syst Rev*. Oct 19 2005; (4): CD002049. PMID 16235295
131. Meireles SM, Jones A, Jennings F, et al. Assessment of the effectiveness of low-level laser therapy on the hands of patients with rheumatoid arthritis: a randomized double-blind controlled trial. *Clin Rheumatol*. May 2010; 29(5): 501-9. PMID 20082104
132. Alayat MS, Elsodany AM, El Fiky AA. Efficacy of high and low level laser therapy in the treatment of Bell's palsy: a randomized double blind placebo-controlled trial. *Lasers Med Sci*. Jan 2014; 29(1): 335-42. PMID 23709010
133. Ordahan B, Karahan AY. Role of low-level laser therapy added to facial expression exercises in patients with idiopathic facial (Bell's) palsy. *Lasers Med Sci*. May 2017; 32(4): 931-936. PMID 28337563
134. Honda Y, Sakamoto J, Hamaue Y, et al. Effects of Physical-Agent Pain Relief Modalities for Fibromyalgia Patients: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Pain Res Manag*. 2018; 2018: 2930632. PMID 30402199
135. Ruaro JA, Frez AR, Ruaro MB, et al. Low-level laser therapy to treat fibromyalgia. *Lasers Med Sci*. Nov 2014; 29(6): 1815-9. PMID 24801056
136. Matsutani LA, Marques AP, Ferreira EA, et al. Effectiveness of muscle stretching exercises with and without laser therapy at tender points for patients with fibromyalgia. *Clin Exp Rheumatol*. May-Jun 2007; 25(3): 410-5. PMID 17631737
137. Samson D, Lefevre F, Aronson N. Wound-healing technologies: low-level laser and vacuum-assisted closure. *Evid Rep Technol Assess (Summ)*. Dec 2004; (111): 1-6. PMID 15663354
138. Chen C, Hou WH, Chan ES, et al. Phototherapy for treating pressure ulcers. *Cochrane Database Syst Rev*. Jul 11 2014; (7): CD009224. PMID 25019295
139. Machado RS, Viana S, Sbruzzi G. Low-level laser therapy in the treatment of pressure ulcers: systematic review. *Lasers Med Sci*. May 2017; 32(4): 937-944. PMID 28116536
140. Taradaj J, Halski T, Kucharzewski M, et al. Effect of laser irradiation at different wavelengths (940, 808, and 658 nm) on pressure ulcer healing: results from a clinical study. *Evid Based Complement Alternat Med* 2013;2013:960240. PMID
141. Lucas C, van Gemert MJ, de Haan RJ. Efficacy of low-level laser therapy in the management of stage III decubitus ulcers: a prospective, observer-blinded multicentre randomised clinical trial. *Lasers Med Sci*. 2003; 18(2): 72-7. PMID 12928815
142. Nussbaum EL, Biemann I, Mustard B. Comparison of ultrasound/ultraviolet-C and laser for treatment of pressure ulcers in patients with spinal cord injury. *Phys Ther*. Sep 1994; 74(9): 812-23; discussion 824-5. PMID 8066108
143. Taly AB, Sivaraman Nair KP, Murali T, et al. Efficacy of multiwavelength light therapy in the treatment of pressure ulcers in subjects with disorders of the spinal cord: A randomized double-blind controlled trial. *Arch Phys Med Rehabil*. Oct 2004; 85(10): 1657-61. PMID 15468027
144. Li S, Wang C, Wang B, et al. Efficacy of low-level light therapy for treatment of diabetic foot ulcer: A systematic review and meta-analysis of randomized controlled trials. *Diabetes Res Clin Pract*. Sep 2018; 143: 215-224. PMID 30009935
145. Smoot B, Chiavola-Larson L, Lee J, et al. Effect of low-level laser therapy on pain and swelling in women with breast cancer-related lymphedema: a systematic review and meta-analysis. *J Cancer Surviv*. Jun 2015; 9(2): 287-304. PMID 25432632
146. Omar MT, Shaheen AA, Zafar H. A systematic review of the effect of low-level laser therapy in the management of breast cancer-related lymphedema. *Support Care Cancer*. Nov 2012; 20(11): 2977-84. PMID 22875413
147. Sung L, Robinson P, Treister N, et al. Guideline for the prevention of oral and oropharyngeal mucositis in children receiving treatment for cancer or undergoing haematopoietic stem cell transplantation. *BMJ Support Palliat Care*. Mar 2017; 7(1): 7-16. PMID 25818385
148. Elad S, Cheng KKF, Lalla RV, et al. MASCC/ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. *Cancer*. Oct 01 2020; 126(19): 4423-4431. PMID 32786044
149. Martin RL, Chimenti R, Cuddeford T, et al. Achilles Pain, Stiffness, and Muscle Power Deficits: Midportion Achilles Tendinopathy Revision 2018. *J Orthop Sports Phys Ther*. May 2018; 48(5): A1-A38. PMID 29712543
150. Carcia CR, Martin RL, Houck J, et al. Achilles pain, stiffness, and muscle power deficits: achilles tendinitis. *J Orthop Sports Phys Ther*. Sep 2010; 40(9): A1-26. PMID 20805627
151. National Institute for Health and Care Excellence (NICE). Low back pain in adults: early management [CG88]. 2009; <https://www.nice.org.uk/guidance/CG88>. Accessed April 26, 2022.
152. North American Spine Society. Evidence-based clinical guidelines for multidisciplinary spine care. 2020. <https://www.spine.org/Portals/0/assets/downloads/ResearchClinicalCare/Guidelines/LowBackPain.pdf>. Accessed April 25, 2022.

153. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med.* Apr 04 2017; 166(7): 514-530. PMID 28192789
154. Qaseem A, McLean RM, O'Gurek D, et al. Nonpharmacologic and Pharmacologic Management of Acute Pain From Non-Low Back, Musculoskeletal Injuries in Adults: A Clinical Guideline From the American College of Physicians and American Academy of Family Physicians. *Ann Intern Med.* Nov 03 2020; 173(9): 739-748. PMID 32805126
155. American Academy of Orthopaedic Surgeons. Management of Carpal Tunnel Syndrome: Evidence-Based Clinical Guideline. 2016; <https://www.aaos.org/quality/quality-programs/upper-extremity-programs/carpal-tunnel-syndrome/>. Accessed April 25, 2022.

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
June 2012	New policy	
March 2013	Replace policy	Policy updated with literature review through September 2012, references added and reordered, policy statement unchanged.
March 2014	Replace policy	Policy updated with literature review. References 14, 37, 46-47, 56, 59- 60 added. Policy statement unchanged.
March 2015	Replace policy	Policy updated with literature review. References 29, 31, 58-59 and 68 added. Policy statement unchanged.
December 2016	Replace policy	Policy updated with literature review; references 5, 9-11, 15, 17, 25, 35, 37, 48, and 51 added. Statement added that low-level laser therapy may be considered medically necessary for prevention of oral mucositis in selected patients. Not medically necessary statement changed to investigational with added bullet points.
September 2018	Replace policy	Policy updated with literature review through April 26, 2018; references 13, 26, 29, 36, 40, 45-49, 53, 55, and 60 added; reference 45 updated. Policy statements unchanged.
September 2019	Replace policy	Policy updated with literature review through April 3, 2019, references added. Policy statements unchanged.
December 2020	Replace policy	Policy updated with literature review through August 20, 2020, references added. Policy statements unchanged.
September 2021	Replace policy	Policy updated with literature review through April 23, 2021; references added. Policy statements unchanged.
September 2022	Replace policy	Policy updated with literature review through April 25, 2022; references added. Minor editorial changes to policy statements; intent unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.