



FEP Medical Policy Manual

FEP 2.02.15 Wearable Cardioverter Defibrillators

Effective Policy Date: October 1, 2021

Original Policy Date: December 2011

Related Policies:

None

Wearable Cardioverter Defibrillators

Description

Description

A wearable cardioverter defibrillator (WCD) is a temporary, external device that is an alternative to an implantable cardioverter defibrillator (ICD). It is primarily intended for temporary conditions for which an implantable device is contraindicated, or for the period during which the need for a permanent implantable device is uncertain.

OBJECTIVE

The objective of this evidence review is to assess whether the use of a wearable cardioverter defibrillator improves net health outcome in patients with a temporary contraindication to implantable cardioverter defibrillator or as a bridge to implantable cardioverter defibrillator placement or heart transplantation, or recovery.

POLICY STATEMENT

Use of wearable cardioverter defibrillators (WCDs) for the prevention of sudden cardiac death is considered **medically necessary** as interim treatment for those who:

- meet the criteria for an implantable cardioverter defibrillator (ICD; see indications in evidence review 7.01.44); and
- have a temporary contraindication to receiving an ICD, such as a systemic infection, at the current time; and
- have been scheduled for an ICD placement or who had an ICD removed and have been rescheduled for placement of another ICD once the contraindication is treated.

Use of WCDs for the prevention of sudden cardiac death is considered **not medically necessary** for the following indications when they are the sole indication for a WCD:

- Patients in the immediate (ie, <40 days) period following an acute myocardial infarction
- Patients post coronary artery bypass graft surgery
- High-risk patients awaiting heart transplant
- Patients with newly diagnosed nonischemic cardiomyopathy
- Women with peripartum cardiomyopathy.

Use of WCDs is considered **not medically necessary** for all other indications.

POLICY GUIDELINES

It is uncommon for patients to have a temporary contraindication to implantable cardioverter defibrillator placement. The most common reason will be a systemic infection that requires treatment before the implantable cardioverter defibrillator can be implanted. The wearable cardioverter defibrillator should only be used short-term while the temporary contraindication (eg, systemic infection) is being clinically managed. Once treatment is completed, the permanent implantable cardioverter defibrillator should be implanted.

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

FDA REGULATORY STATUS

In 2001, the Lifecor WCD 2000 system was approved by the FDA through the premarket approval process for "adult patients who are at risk for cardiac arrest and are either not candidates for or refuse an implantable defibrillator." The vest was renamed the Zoll LifeVest.

In 2015, the FDA approved the LifeVest for certain children who are at risk for sudden cardiac arrest, but are not candidates for an implantable defibrillator due to certain medical conditions or lack of parental consent."

FDA product code: MVK.

RATIONALE

Summary of Evidence

Overview of Wearable Cardioverter Defibrillator Versus Implantable Cardioverter Defibrillator

One randomized controlled trial (RCT) has compared wearable cardioverter defibrillator (WCD) with usual guideline-based care and found no significant benefit to WCD over usual care. No studies have directly compared the performance of a WCD with a permanent implantable cardiac defibrillator (ICD). One small study in an electrophysiology lab demonstrated that the WCD can correctly identify and terminate most induced ventricular arrhythmias. A cohort study of WCD use estimated that the percentage of successful resuscitations was approximately 70%. Multiple studies have demonstrated suboptimal adherence. Device failures were largely attributed to incorrect device use and/or nonadherence. A more recent registry study has reported a high compliance rate, although these results may be biased by self-selection. Collectively, this evidence indicates that the WCD can successfully detect and terminate arrhythmias in at least some patients but that overall performance in clinical practice might be inferior to a permanent ICD.

Temporary Contraindications

For individuals who have a temporary contraindication to an ICD who receive a WCD, the evidence includes prospective cohort studies and a technology assessment that assessed ICD devices, given the absence of evidence on WCD devices. Relevant outcomes are overall survival (OS), morbid events, functional outcomes, and treatment-related morbidity. A small number of patients meet established criteria for an ICD but have a transient contraindication for an implantable device, most commonly an infectious process. The available data have established that the WCD device can detect lethal arrhythmias and can successfully deliver a countershock in most cases. In patients scheduled for ICD placement, the WCD will improve outcomes as an interim treatment. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Immediate Post Myocardial Infarction

For individuals who are in the immediate post myocardial infarction period who receive a WCD, the evidence includes an RCT comparing WCD with guideline-based therapy, 2 cohort studies, and a systematic review. Relevant outcomes are OS, morbid events, functional outcomes, and treatment-related morbidity. The RCT reported no benefit of WCD over guideline-based therapy. The cohort study of 8453 patients showed that 252 shocks successfully terminated ventricular fibrillation or ventricular tachycardia (82% success rate), but without a control group, interpretation is difficult. Similarly, a retrospective cohort of Medicare data found that WCD use was associated with lower 1-year mortality than no WCD use, but potential biases were noted. Evidence from the systematic review was deemed of low to very low quality, and the reviewers had weak confidence in the reported estimates. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Post - Coronary Artery Bypass Graft Surgery at High Risk for Lethal Arrhythmias

For individuals who are post - coronary artery bypass graft surgery and are at high risk for lethal arrhythmias, the evidence includes an RCT for ICD and a registry study. Relevant outcomes are OS, morbid events, functional outcomes, and treatment-related morbidity. For high-risk post coronary artery bypass graft patients, an RCT reported no difference in OS associated with early ICD placement. The registry study found survival benefits with WCD but had limited interpretation of data. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Awaiting Heart Transplantation at High Risk for Lethal Arrhythmias

For individuals who are awaiting heart transplantation and are at high risk for lethal arrhythmias, the evidence includes analyses of subsets of patients from the manufacturer registry, a subset from a prospective cohort study, and a case series. Relevant outcomes are OS, morbid events, functional outcomes, and treatment-related morbidity. These studies do not provide sufficient evidence to determine whether a WCD is of benefit compared with usual care. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Newly Diagnosed Nonischemic Cardiomyopathy

For individuals who have newly diagnosed nonischemic cardiomyopathy, the evidence includes an RCT for ICD and several retrospective analyses of WCD registry data. Relevant outcomes are OS, morbid events, functional outcomes, and treatment-related morbidity. The RCT found that prophylactic ICD placement for nonischemic cardiomyopathy did not improve mortality compared with usual care. Evidence from the retrospective analysis was not sufficient to determine whether WCD improves outcomes compared with usual care. Given the lack of evidence that ICD improves outcomes, WCD is not expected to improve outcomes under the conditions studied in these trials. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Peripartum Cardiomyopathy

For individuals who have peripartum cardiomyopathy, the evidence includes a retrospective registry data analysis and a small cohort study. Relevant outcomes are OS, morbid events, functional outcomes, and treatment-related morbidity. The registry study revealed that no shocks were delivered during use over an average of 124 days. The cohort study identified 4 episodes of appropriate electric shock over 133 days. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in "Supplemental Information" if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Heart Association et al

In 2018, the American Heart Association (AHA), the American College of Cardiology and the Heart Rhythm Society published a guideline on the management of patients with ventricular arrhythmias and prevention of sudden cardiac death.²⁹ The guidelines note that "the patients listed in this recommendation are represented in clinical series and registries that demonstrate the safety and effectiveness of the wearable cardioverter-defibrillator. Patients with recent MI, newly diagnosed NICM, recent revascularization, myocarditis, and secondary cardiomyopathy are at increased risk of VT/SCA. However, the wearable cardioverter-defibrillator is of unproven benefit in these settings, in part because the clinical situation may improve with therapy and time." The specific recommendations are summarized in Table 1. Class IIa is moderate recommendation, and class IIb is a weak recommendation.

Table 1. Guidelines for Wearable Cardioverter Defibrillator Therapy

Recommendation	COR	LOE
"In patients with an ICD and a history of SCA or sustained VA in whom removal of the ICD is required (as with infection), the wearable cardioverter-defibrillator is reasonable for the prevention of SCD."	IIa	B-NR
"In patients at an increased risk of SCD but who are not ineligible for an ICD, such as awaiting cardiac transplant, having an LVEF of 35% or less and are within 40 days from an MI, or have newly diagnosed NICM, revascularization within the past 90 days, myocarditis or secondary cardiomyopathy or a systemic infection, the wearable cardioverter-defibrillator may be reasonable."	IIb	B-NR

B-NR: Level B - nonrandomized; COR: class of recommendation; ICD: implantable cardioverter defibrillator; LOE: *level of evidence*; LVEF: left ventricular ejection fraction; MI: myocardial infarction; NICM: non-ischemic cardiomyopathy; SCA: sudden cardiac arrest; SCD: sudden cardiac death; WCD: wearable cardioverter defibrillator.

In 2016, the AHA published a scientific advisory on the WCD.³⁰ The AHA stated that "because there is a paucity of prospective data supporting the use of the WCD, particularly in the absence of any published, randomized, clinical trials, the recommendations provided in this advisory are not

intended to be prescriptive or to suggest an evidence-based approach to the management of patients with FDA-approved indications for use." The specific recommendations are summarized in Table 2.

Table 2. Guidelines for Wearable Cardioverter Defibrillator Therapy

Recommendation	COR	LOE
"Use of WCDs is reasonable when there is a clear indication for an implanted/permanent device accompanied by a transient contraindication or interruption in ICD care such as infection."	IIa	C
"Use of WCDs is reasonable as a bridge to more definitive therapy such as cardiac transplantation"	IIa	C
"Use of WCDs may be reasonable when there is concern about a heightened risk of SCD that may resolve over time or with treatment of left ventricular dysfunction/ for example, in ischemic heart disease with recent revascularization, newly diagnosed nonischemic dilated cardiomyopathy in patients starting guideline-directed medical therapy, or secondary cardiomyopathy (tachycardia mediated, thyroid mediated, etc) in which the underlying cause is potentially treatable."	IIb	C
"WCDs may be appropriate as bridging therapy in situations associated with increased risk of death in which ICDs have been shown to reduce SCD but not overall survival such as within 40 D of MI."	IIb	C
"WCDs should not be used when nonarrhythmic risk is expected to significantly exceed arrhythmic risk, particularly in patients who are not expected to survive >6 mo."	III	C

COR: class of recommendation; ICD: implantable cardioverter defibrillator; LOE: *level of evidence*; MI: myocardial infarction; SCD: sudden cardiac death; WCD: wearable cardioverter defibrillator.

American College of Cardiology et al

In 2018, the American College of Cardiology, AHA, and the Heart Rhythm Society jointly published guidelines on the management of adults who have ventricular arrhythmias or who are at risk for sudden cardiac death, including diseases and syndromes associated with a risk of sudden cardiac death from ventricular arrhythmias.²⁹ Recommendations related to the use of WCDs are provided in Table 3.

Table 3. Guidelines for Wearable Cardioverter Defibrillator Therapy

Recommendations	COR	LOE
In patients with an implantable cardioverter defibrillator and a history of sudden cardiac arrest or sustained ventricular arrhythmia in whom removal of the implantable cardioverter defibrillator is required (as with infection), the wearable cardioverter defibrillator is reasonable for the prevention of sudden cardiac death. ^a	IIa	B-NR
In patients at an increased risk of sudden cardiac death but who are not ineligible for an implantable cardioverter defibrillator, such as awaiting cardiac transplant, having an LVEF of 35% or less and are within 40 days from an MI, or have newly diagnosed nonischemic cardiomyopathy, revascularization within the past 90 days, myocarditis or secondary cardiomyopathy or a systemic infection, wearable cardioverter defibrillator may be reasonable. ^b	IIb	B-NR

B-NR: data derived from ≥ 1 nonrandomized trials or meta-analysis of such studies; COR: class of recommendation; ICD: implantable cardioverter defibrillator; LOE: level of evidence; LVEF: left ventricular ejection fraction; MI: myocardial infarction.

^a Removal of an ICD for a period of time, most commonly due to infection, exposes the patient to risk of untreated ventricular tachycardia/sudden cardiac death unless monitoring and access to emergency external defibrillation is maintained. In 1 series of 354 patients who received the WCD, the indication was infection in 10%.³¹ For patients with a history of sudden cardiac arrest or sustained ventricular arrhythmia, the WCD may allow the patient to be discharged from the hospital with protection from ventricular tachycardia/sudden cardiac death until the clinical situation allows reimplantation of an ICD.

^b The patients listed in this recommendation are represented in clinical series and registries that demonstrate the safety and effectiveness of the WCD. Patients with recent MI, newly

diagnosed nonischemic cardiomyopathy, recent revascularization, myocarditis, and secondary cardiomyopathy are at increased risk of ventricular tachycardia or sudden cardiac death. However, the WCD is of unproven benefit in these settings, in part because the clinical situation may improve with therapy and time. In patients awaiting transplant, even with anticipated survival <1 year without transplant, and depending on clinical factors such as use of intravenous inotropes and ambient ventricular arrhythmia, a WCD may be an alternative to an ICD.

International Society for Heart and Lung Transplantation

In 2006, the International Society for Heart and Lung Transplantation issued guidelines on the care of cardiac transplant candidates that addressed use of ICDs or WCDs.³² Recommendations on the use of WCDs are provided in Table 4.

Table 4. Guidelines on Management of Cardiac Transplant Candidates With ICDs

Recommendation	COR	LOE
"An implanted or wearable ICD should be provided for Status 1B patients [ie, dependent on intravenous medications or a mechanical assist device] who are discharged home given that the wait for transplantation remains significant."	I	C
"It is reasonable to consider placement of a defibrillator in patients with Stage D failure who are candidates for transplantation or LVAD destination therapy (see subsequent considerations for MCSD referral: bridge or destination)."	Ila	C

COR: class of recommendation; ICD: implantable cardioverter defibrillator; LOE: *level of evidence*; LVAD: left ventricular assist device; MCSD: mechanical circulatory support device.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

REFERENCES

1. Food and Drug Administration. Summary of Safety and Effectiveness Data, P010030, Lifecor, Inc. WCD 2000 System. 2001; https://www.accessdata.fda.gov/cdrh_docs/pdf/p010030b.pdf. Accessed April 12, 2021.
2. Beauregard LA. Personal security: Clinical applications of the wearable defibrillator. *Pacing Clin Electrophysiol*. Jan 2004; 27(1): 1-3. PMID 14720147
3. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Wearable cardioverter-defibrillator as a bridge to implantable cardioverter-defibrillator treatment. *TEC Assessments*. 2010;Volume 25:Tab 2.
4. Auricchio A, Klein H, Geller CJ, et al. Clinical efficacy of the wearable cardioverter-defibrillator in acutely terminating episodes of ventricular fibrillation. *Am J Cardiol*. May 15 1998; 81(10): 1253-6. PMID 9604964
5. Chung MK, Szymkiewicz SJ, Shao M, et al. Aggregate national experience with the wearable cardioverter-defibrillator: event rates, compliance, and survival. *J Am Coll Cardiol*. Jul 13 2010; 56(3): 194-203. PMID 20620738
6. Tanawuttiwat T, Garisto JD, Salow A, et al. Protection from outpatient sudden cardiac death following ICD removal using a wearable cardioverter defibrillator. *Pacing Clin Electrophysiol*. May 2014; 37(5): 562-8. PMID 24762055
7. Mitrani RD, McArdle A, Slane M, et al. Wearable defibrillators in uninsured patients with newly diagnosed cardiomyopathy or recent revascularization in a community medical center. *Am Heart J*. Mar 2013; 165(3): 386-92. PMID 23453108
8. Kao AC, Krause SW, Handa R, et al. Wearable defibrillator use in heart failure (WIF): results of a prospective registry. *BMC Cardiovasc Disord*. Dec 12 2012; 12: 123. PMID 23234574
9. Feldman AM, Klein H, Tchou P, et al. Use of a wearable defibrillator in terminating tachyarrhythmias in patients at high risk for sudden death: results of the WEARIT/BIROAD. *Pacing Clin Electrophysiol*. Jan 2004; 27(1): 4-9. PMID 14720148
10. Kutyifa V, Moss AJ, Klein H, et al. Use of the wearable cardioverter defibrillator in high-risk cardiac patients: data from the Prospective Registry of Patients Using the Wearable Cardioverter Defibrillator (WEARIT-II Registry). *Circulation*. Oct 27 2015; 132(17): 1613-9. PMID 26316618

11. Gregoratos G, Cheitlin MD, Conill A, et al. ACC/AHA guidelines for implantation of cardiac pacemakers and antiarrhythmia devices: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Pacemaker Implantation). *J Am Coll Cardiol*. Apr 1998; 31(5): 1175-209. PMID 9562026
12. Hohnloser SH, Kuck KH, Dorian P, et al. Prophylactic use of an implantable cardioverter-defibrillator after acute myocardial infarction. *N Engl J Med*. Dec 09 2004; 351(24): 2481-8. PMID 15590950
13. Steinbeck G, Andresen D, Seidl K, et al. Defibrillator implantation early after myocardial infarction. *N Engl J Med*. Oct 08 2009; 361(15): 1427-36. PMID 19812399
14. Olgin JE, Pletcher MJ, Vittinghoff E, et al. Wearable Cardioverter-Defibrillator after Myocardial Infarction. *N Engl J Med*. Sep 27 2018; 379(13): 1205-1215. PMID 30280654
15. Epstein AE, Abraham WT, Bianco NR, et al. Wearable cardioverter-defibrillator use in patients perceived to be at high risk early post-myocardial infarction. *J Am Coll Cardiol*. Nov 19 2013; 62(21): 2000-2007. PMID 23916930
16. Clark MA, Szymkiewicz SJ, Volosin K. Mortality and Costs Associated with Wearable Cardioverter-defibrillators after Acute Myocardial Infarction: A Retrospective Cohort Analysis of Medicare Claims Data. *J Innov Card Rhythm Manag*. Oct 2019; 10(10): 3866-3873. PMID 32477706
17. Uyei J, Braithwaite RS. Effectiveness of wearable defibrillators: systematic review and quality of evidence. *Int J Technol Assess Health Care*. Apr 2014; 30(2): 194-202. PMID 24893969
18. Bigger JT. Prophylactic use of implanted cardiac defibrillators in patients at high risk for ventricular arrhythmias after coronary-artery bypass graft surgery. *Coronary Artery Bypass Graft (CABG) Patch Trial Investigators*. *N Engl J Med*. Nov 27 1997; 337(22): 1569-75. PMID 9371853
19. Zishiri ET, Williams S, Cronin EM, et al. Early risk of mortality after coronary artery revascularization in patients with left ventricular dysfunction and potential role of the wearable cardioverter defibrillator. *Circ Arrhythm Electrophysiol*. Feb 2013; 6(1): 117-28. PMID 23275233
20. Opreanu M, Wan C, Singh V, et al. Wearable cardioverter-defibrillator as a bridge to cardiac transplantation: A national database analysis. *J Heart Lung Transplant*. Oct 2015; 34(10): 1305-9. PMID 26094085
21. Wassnig NK, Gunther M, Quick S, et al. Experience With the Wearable Cardioverter-Defibrillator in Patients at High Risk for Sudden Cardiac Death. *Circulation*. Aug 30 2016; 134(9): 635-43. PMID 27458236
22. Rao M, Goldenberg I, Moss AJ, et al. Wearable defibrillator in congenital structural heart disease and inherited arrhythmias. *Am J Cardiol*. Dec 01 2011; 108(11): 1632-8. PMID 21890075
23. Kadish A, Schaechter A, Subacius H, et al. Patients with recently diagnosed nonischemic cardiomyopathy benefit from implantable cardioverter-defibrillators. *J Am Coll Cardiol*. Jun 20 2006; 47(12): 2477-82. PMID 16781376
24. Salehi N, Nasiri M, Bianco NR, et al. The Wearable Cardioverter Defibrillator in Nonischemic Cardiomyopathy: A US National Database Analysis. *Can J Cardiol*. Oct 2016; 32(10): 1247.e1-1247.e6. PMID 26975224
25. Duncker D, Konig T, Hohmann S, et al. Ventricular arrhythmias in patients with newly diagnosed nonischemic cardiomyopathy: Insights from the PROLONG study. *Clin Cardiol*. Aug 2017; 40(8): 586-590. PMID 28333373
26. Duncker D, Konig T, Hohmann S, et al. Avoiding Untimely Implantable Cardioverter/Defibrillator Implantation by Intensified Heart Failure Therapy Optimization Supported by the Wearable Cardioverter/Defibrillator-The PROLONG Study. *J Am Heart Assoc*. Jan 17 2017; 6(1). PMID 28096098
27. Saltzberg MT, Szymkiewicz S, Bianco NR. Characteristics and outcomes of peripartum versus nonperipartum cardiomyopathy in women using a wearable cardiac defibrillator. *J Card Fail*. Jan 2012; 18(1): 21-7. PMID 22196837
28. Duncker D, Haghikia A, Konig T, et al. Risk for ventricular fibrillation in peripartum cardiomyopathy with severely reduced left ventricular function-value of the wearable cardioverter/defibrillator. *Eur J Heart Fail*. Dec 2014; 16(12): 1331-6. PMID 25371320
29. Al-Khatib SM, Stevenson WG, Ackerman MJ, et al. 2017 AHA/ACC/HRS Guideline for Management of Patients With Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Circulation*. Sep 25 2018; 138(13): e272-e391. PMID 29084731
30. Piccini JP, Allen LA, Kudenchuk PJ, et al. Wearable Cardioverter-Defibrillator Therapy for the Prevention of Sudden Cardiac Death: A Science Advisory From the American Heart Association. *Circulation*. Apr 26 2016; 133(17): 1715-27. PMID 27022063
31. Klein HU, Meltendorf U, Reek S, et al. Bridging a temporary high risk of sudden arrhythmic death. Experience with the wearable cardioverter defibrillator (WCD). *Pacing Clin Electrophysiol*. Mar 2010; 33(3): 353-67. PMID 19889186
32. Gronda E, Bourge RC, Costanzo MR, et al. Heart rhythm considerations in heart transplant candidates and considerations for ventricular assist devices: International Society for Heart and Lung Transplantation guidelines for the care of cardiac transplant candidates--2006. *J Heart Lung Transplant*. Sep 2006; 25(9): 1043-56. PMID 16962465

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
December 2011	New policy	
December 2012	Replace policy	Policy updated with literature review, reference 6 updated, reference 14 added. Wording "have all of the following" stricken from medically necessary policy statement. No other changes to policy statement.
March 2014	Replace policy	Policy updated with literature review through August 2013, references 6-7, 13 and 15 added. No change to policy statement. Removed "as a Bridge to Implantable Cardioverter-Defibrillator Placement" from the title.
March 2015	Replace policy	Policy updated with literature review through November 30, 2014. References 17, 23, and 26-27 added. Investigational policy statements changed to not medically necessary.
September 2016	Replace policy	Policy updated with literature review, references 13, 20, 30-32 added. FDA regulatory status updated. Policy statements and guideline revised.
September 2018	Replace policy	Policy updated with literature review through March 14, 2018; reference 27-28 added; reference 1 updated. "High-risk patients awaiting heart transplant" was added to the not medically necessary policy statement; and an additional policy statement that use of wearable cardioverter-defibrillators is considered not medically necessary for all other indications was added.
May 2019	Replace policy	Policy updated with literature review through March 4, 2019.; reference 31 added. Policy statements unchanged.
September 2020	Replace policy	Policy updated with literature review through March 9, 2020; reference added. Policy statements unchanged.
September 2021	Replace policy	Policy updated with literature review through April 10, 2021; reference added. Policy statements unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.