



FEP Medical Policy Manual

FEP 1.01.24 Interferential Current Stimulation

Effective Policy Date: October 1, 2020

Related Policies:

Original Policy Date: September 2011

1.01.09 - Transcutaneous Electrical Nerve Stimulation
7.01.29 - Percutaneous Electrical Nerve Stimulation and Percutaneous Neuromodulation Therapy

Interferential Current Stimulation

Description

Description

Interferential current stimulation (IFS) is a type of electrical stimulation used to reduce pain. The technique has been proposed to decrease pain and increase function in patients with osteoarthritis and to treat other conditions such as constipation, irritable bowel syndrome, dyspepsia, and spasticity.

Interferential current stimulation (IFS) is a type of electrical stimulation that has been investigated as a technique to reduce pain, improve function and range of motion, and treat gastrointestinal disorders.

IFS uses paired electrodes of 2 independent circuits carrying high-frequency and medium-frequency alternating currents. The superficial electrodes are aligned on the skin around the affected area. It is believed that IFS permeates the tissues more effectively and with less unwanted stimulation of cutaneous nerves, is more comfortable than transcutaneous electrical nerve stimulation. There are no standardized protocols for the use of IFS; IFS may vary by the frequency of stimulation, the pulse duration, treatment time, and electrode-placement technique.

OBJECTIVE

The objective of this evidence review is to determine whether inferential current stimulation improves the net health outcome in patients with musculoskeletal conditions, gastrointestinal disorders, or post-stroke spasticity.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

POLICY STATEMENT

Interferential current stimulation is considered **investigational**.

POLICY GUIDELINES

None

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

FDA REGULATORY STATUS

A number of IFS devices have been cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process, including the Medstar™ 100 (MedNet Services) and the RS-4i (RS Medical). IFS may be included in multimodal electrotherapy devices such as transcutaneous electrical nerve stimulation and functional electrostimulation.

RATIONALE

Summary of Evidence

For individuals who have musculoskeletal conditions who receive Interferential current stimulation (IFS), the evidence includes randomized controlled trials (RCTs) and meta-analyses. Relevant outcomes are symptoms, functional outcomes, quality of life, medication use, and treatment-related morbidity. Placebo-controlled randomized trial(s) have found that IFS when used to treat musculoskeletal pain and impaired function(s), does not significantly improve outcomes; additionally, a meta-analysis of placebo-controlled trials did not find a significant benefit of IFS for decreasing pain or improving function. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have gastrointestinal disorders who receive IFS, the evidence includes RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, medication use, and treatment-related morbidity. IFS has been tested for a variety of gastrointestinal conditions, with a small number of trials completed for each condition. The results of the trials are mixed, with some reporting benefit and others not. This body of evidence is inconclusive on whether IFS is an efficacious treatment for gastrointestinal conditions. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have post-stroke spasticity who receive IFS, the evidence includes RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The RCTs had small sample sizes and very short follow-up (immediately post-treatment to 5 weeks). The evidence is insufficient to determine the effects of the technology on health outcomes.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

American College of Physicians and the American Pain Society

In 2009, the clinical practice guidelines from the American College of Physicians and the American Pain Society concluded that there was insufficient evidence to recommend interferential current stimulation (IFS) for the treatment of low back pain.²⁴ An update of these guidelines by the American College of Physicians (2017) confirmed the 2009 findings that there was insufficient evidence to determine the effectiveness of interferential current stimulation (IFS) for the treatment of low back pain.²⁵

American College of Occupational and Environmental Medicine

The American College of Occupational and Environmental Medicine published several relevant guidelines. For shoulder disorders, guidelines found the evidence on IFS to be insufficient and, depending on the specific disorder, either did not recommend IFS or were neutral on whether to recommend it.²⁶ For low back disorders, guidelines found the evidence on IFS to be insufficient and did not recommend it. The sole exception was that IFS could be considered as an option on a limited basis for acute low back pain with or without radicular pain.²⁷ For knee disorders, guidelines recommended IFS for postoperative anterior cruciate ligament reconstruction, meniscectomy, and knee chondroplasty immediately postoperatively in the elderly.²⁸ This was a level C recommendation.

National Institute for Health and Care Excellence

In 2016, the National Institute for Health and Care Excellence had a guideline (NG59) on assessment and management of low back pain and sciatica in people aged 16 and over.³ The guideline states "Do not offer interferential therapy for managing low back pain with or without sciatica".

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

REFERENCES

1. Zeng C, Li H, Yang T, et al. Electrical stimulation for pain relief in knee osteoarthritis: systematic review and network meta-analysis. *Osteoarthritis Cartilage*. Feb 2015;23(2):189-202. PMID 25497083
2. Ferreira RM, Torres RT, Duarte JA, et al. Non-Pharmacological and Non-Surgical Interventions for Knee Osteoarthritis: A Systematic Review and Meta-Analysis. *Acta Reumatol Port*. Jul 29 2019; 44(3): 173-217. PMID 31356585
3. National Institute for Health and Care Excellence (NICE). Low back pain and sciatica in over 16s: assessment and management [NG59]. 2016; <https://www.nice.org.uk/guidance/ng59>. Accessed April 21, 2020.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

4. Fuentes JP, Armijo Olivo S, Magee DJ, et al. Effectiveness of interferential current therapy in the management of musculoskeletal pain: a systematic review and meta-analysis. *Phys Ther*. Sep 2010;90(9):1219-1238. PMID 20651012
5. Defrin R, Ariel E, Peretz C. Segmental noxious versus innocuous electrical stimulation for chronic pain relief and the effect of fading sensation during treatment. *Pain*. May 2005;115(1-2):152-160. PMID 15836978
6. Taylor K, Newton RA, Personius WJ, et al. Effects of interferential current stimulation for treatment of subjects with recurrent jaw pain. *Phys Ther*. Mar 1987;67(3):346-350. PMID 3493493
7. Atamaz FC, Durmaz B, Baydar M, et al. Comparison of the efficacy of transcutaneous electrical nerve stimulation, interferential currents, and shortwave diathermy in knee osteoarthritis: a double-blind, randomized, controlled, multicenter study. *Arch Phys Med Rehabil*. May 2012;93(5):748-756. PMID 22459699
8. Gundog M, Atamaz F, Kanyilmaz S, et al. Interferential current therapy in patients with knee osteoarthritis: comparison of the effectiveness of different amplitude-modulated frequencies. *Am J Phys Med Rehabil*. Feb 2012;91(2):107-113. PMID 22019968
9. Kadi MR, Hepguler S, Atamaz FC, et al. Is interferential current effective in the management of pain, range of motion, and edema following total knee arthroplasty surgery? A randomized double-blind controlled trial. *Clin Rehabil*. Jun 2019; 33(6): 1027-1034. PMID 30764635
10. Koca I, Boyaci A, Tutoglu A, et al. Assessment of the effectiveness of interferential current therapy and TENS in the management of carpal tunnel syndrome: a randomized controlled study. *Rheumatol Int*. Dec 2014;34(12):1639-1645. PMID 24728028
11. Lara-Palomo IC, Aguilar-Ferrandiz ME, Mataran-Penarrocha GA, et al. Short-term effects of interferential current electro-massage in adults with chronic non-specific low back pain: a randomized controlled trial. *Clin Rehabil*. May 2013;27(5):439-449. PMID 23035006
12. Facci LM, Nowotny JP, Tormem F, et al. Effects of transcutaneous electrical nerve stimulation (TENS) and interferential currents (IFC) in patients with nonspecific chronic low back pain: randomized clinical trial. *Sao Paulo Med J*. 2011;129(4):206-216. PMID 21971895
13. Albornoz-Cabello M, Maya-Martin J, Dominguez-Maldonado G, et al. Effect of interferential current therapy on pain perception and disability level in subjects with chronic low back pain: a randomized controlled trial. *Clin Rehabil*. Feb 2017;31(2):242-249. PMID 26975312
14. Dissanayaka TD, Pallegama RW, Suraweera HJ, et al. Comparison of the effectiveness of transcutaneous electrical nerve stimulation and interferential therapy on the upper trapezius in myofascial pain syndrome: a randomized controlled study. *Am J Phys Med Rehabil*. Sep 2016;95(9):663-672. PMID 26945216
15. Albornoz-Cabello M, Perez-Marmol JM, Barrios Quinta CJ, et al. Effect of adding interferential current stimulation to exercise on outcomes in primary care patients with chronic neck pain: a randomized controlled trial. *Clin Rehabil*. Sep 2019; 33(9): 1458-1467. PMID 31007047
16. Iacona R, Ramage L, Malakounides G. Current State of Neuromodulation for Constipation and Fecal Incontinence in Children: A Systematic Review. *Eur J Pediatr Surg*. Dec 2019; 29(6): 495-503. PMID 30650450
17. Kajbafzadeh AM, Sharifi-Rad L, Nejat F, et al. Transcutaneous interferential electrical stimulation for management of neurogenic bowel dysfunction in children with myelomeningocele. *Int J Colorectal Dis*. Apr 2012;27(4):453-458. PMID 22065105
18. Clarke MC, Chase JW, Gibb S, et al. Improvement of quality of life in children with slow transit constipation after treatment with transcutaneous electrical stimulation. *J Pediatr Surg*. Jun 2009;44(6):1268-1272; discussion 1272. PMID 19524752
19. Moore JS, Gibson PR, Burgell RE. Randomised clinical trial: transabdominal interferential electrical stimulation vs sham stimulation in women with functional constipation. *Aliment Pharmacol Ther*. Apr 2020; 51(8): 760-769. PMID 32128859
20. Coban S, Akbal E, Koklu S, et al. Clinical trial: transcutaneous interferential electrical stimulation in individuals with irritable bowel syndrome - a prospective double-blind randomized study. *Digestion*. Aug 2012;86(2):86-93. PMID 22846190
21. Koklu S, Koklu G, Ozguclu E, et al. Clinical trial: interferential electric stimulation in functional dyspepsia patients - a prospective randomized study. *Aliment Pharmacol Ther*. May 2010;31(9):961-968. PMID 20136803
22. Suh HR, Han HC, Cho HY. Immediate therapeutic effect of interferential current therapy on spasticity, balance, and gait function in chronic stroke patients: a randomized control trial. *Clin Rehabil*. Sep 2014;28(9):885-891. PMID 24607801
23. Eslamian F, Farhoudi M, Jahanjoo F, et al. Electrical interferential current stimulation versus electrical acupuncture in management of hemiplegic shoulder pain and disability following ischemic stroke-a randomized clinical trial. *Arch Physiother*. 2020; 10: 2. PMID 31938571
24. Chou R, Atlas SJ, Stanos SP, et al. Nonsurgical interventional therapies for low back pain: a review of the evidence for an American Pain Society clinical practice guideline. *Spine (Phila Pa 1976)*. May 1 2009;34(10):1078-1093. PMID 19363456
25. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med*. Apr 04 2017; 166(7): 514-530. PMID 28192789
26. American College of Occupational and Environmental Medicine (ACOEM). Shoulder disorders. In: Hegmann KT, ed. *Occupational medicine practice guidelines. Evaluation and management of common health problems and functional recovery in workers*. 3rd ed. Elk Grove Village, IL: ACOEM; 2011:1-297.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

27. American College of Occupational and Environmental Medicine (ACOEM). Low Back Disorders. In: Hegmann KT, ed. Occupational medicine practice guidelines: evaluation and management of common health problems and functional recovery in workers. Westminster, CO: Reed Group; 2016:1-844.
28. American College of Occupational and Environmental Medicine (ACOEM). Knee Disorders. In: Hegmann KT, ed. Occupational medicine practice guidelines. Evaluation and management of common health problems and functional recovery in workers. 3rd ed. Elk Grove Village, IL: ACOEM; 2011:1-503.

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
September 2011	New policy	
June 2012	Replace policy	Policy updated with literature search; references 6 and 7 added; other references re-numbered or removed. Policy statement changed to not medically necessary to use IFS for the treatment of pain
March 2013	Replace policy	Policy updated with literature search. References 5, 9-12 added; other references renumbered or removed. Policy changed to include not medically necessary for treatment of other conditions. Title changed to "Interferential Current Stimulation."
March 2014	Replace policy	Policy updated with literature review; references 4 and 12 added; other references renumbered or removed. No change in policy statement.
March 2015	Replace policy	Policy updated with literature review. References 7, 12, and 14-16 added. No change in policy statement.
December 2017	Replace policy	Policy updated with literature review through July 21, 2017; no references added; references 18-20 updated. Policy statement corrected from "not medically necessary" to "investigational" due to FDA 510(k) clearance.
September 2018	Replace policy	Policy updated with literature review through April 9, 2018; reference 17 added. Policy statement unchanged.
September 2019	Replace policy	Policy updated with literature review through April 30, 2019, no references added. Policy statement unchanged.
September 2020	Replace policy	Policy updated with literature review through April 17, 2020; references added. Policy statement unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.