



# FEP Medical Policy Manual

---

## FEP 2.04.56 Immune Cell Function Assay

---

**Effective Policy Date: April 1, 2021**

**Related Policies:**

**Original Policy Date: December 2011**

None

---

## Immune Cell Function Assay

### Description

#### Description

Careful monitoring of lifelong immunosuppression is required to ensure the long-term viability of solid organ allografts without incurring an increased risk of infection. The monitoring of immunosuppression parameters attempts to balance the dual risks of rejection and infection. It is proposed that individual immune profiles, such as an immune cell function assay, will help assess the immune function of the transplant recipient and individualize immunosuppressive therapy.

#### OBJECTIVE

The objective of this evidence review is to determine whether the use of commercially available assays to assess immune cell function in patients with solid organ transplants and hematopoietic cell transplantation improves the net health outcome.

---

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

## POLICY STATEMENT

Use of the immune cell function assay to monitor and predict immune function after solid organ transplantation is considered **investigational**.

Use of the immune cell function assay to monitor and predict immune function after hematopoietic cell transplantation is considered **investigational**.

Use of the immune cell function assay for all other indications is considered **investigational**.

## POLICY GUIDELINES

None

## BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

## FDA REGULATORY STATUS

In April 2002, ImmuKnow (Cylex, acquired by ViraCor-IBT Laboratories), an immune cell function assay, was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process (K013169). The FDA indicated use of ImmuKnow is for the detection of a cell-mediated immune response in populations undergoing immunosuppressive therapy for an organ transplant.

In April 2002, Immune Cell Function Assay (Cylex) was cleared for marketing by the FDA through the 510(k) process. The FDA indicated use of the Immune Cell Function Assay is for the detection of a cell-mediated immune response in an immunosuppressed population. In 2010, a device modification for this assay was cleared for marketing by FDA through the 510(k) (K101911). There were no changes to the indications or intended use.<sup>1</sup>

In August 2014, Pleximmune™ (Plexision) was approved by the FDA through the humanitarian device exemption process.<sup>2</sup> The test is intended for use in the pretransplantation and early and late posttransplantation period in pediatric liver and small bowel transplant patients for the purpose of predicting the risk of transplant rejection within 60 days after transplantation or 60 days after sampling.

## RATIONALE

### Summary of Evidence

For individuals who have a solid organ transplant or hematopoietic cell transplant who receive immune cell function assay testing with ImmuKnow, the evidence includes numerous studies on the association between assay test values and subsequent rejection or infection, and a randomized controlled trial in liver transplant patients. Relevant outcomes are overall survival, other test performance measures, and morbid events. The ImmuKnow test has shown variable associations with infection and rejection, depending on the type of transplant and context of the study. Across all the studies among various types of patients, ImmuKnow levels are associated with the risk of rejection when levels are high and risk of infection when levels are low. However, the absolute risk and increments of risk are uncertain because of the heterogeneity of the studies. The predictive characteristics of the test are still uncertain and do not allow a strong chain of evidence for clinical utility. The trial of the ImmuKnow test in liver transplant patients showed improvement in overall survival; however, the trial had several limitations. The evidence is insufficient to determine that the technology results in an improvement in the net health outcomes.

---

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

For individuals who have a solid organ transplant or hematopoietic cell transplant who receive immune cell function assay testing with Pleximmune, the evidence includes the U.S. Food and Drug Administration (FDA) documentation and a report on the test's development and validation. Relevant outcomes are overall survival, other measures of test performance, and morbid events. Small studies have shown that Pleximmune values correlate with long-term survival. Pleximmune test results correlated with rejection, but conclusions are uncertain because of extremely limited evidence deriving from a small number of patients described briefly in the FDA approval documents and a second study, in which the confidence interval bounds for sensitivity and specificity estimates were wide. No direct studies of clinical utility were identified. An argument for clinical utility using a chain of evidence would rely on both a demonstration of clinical validity and a rationale that specific clinical interventions based on the results of the test decrease the risk of a poor health outcome. At present, the clinical interventions that would occur as a result of the test result are uncertain, and so the clinical validity is uncertain. The evidence is insufficient to determine that the technology results in an improvement in the net health outcomes.

## SUPPLEMENTAL INFORMATION

### Practice Guidelines and Position Statements

#### Transplantation Society

In 2018, the International Cytomegalovirus Consensus Group of the Transplantation Society updated its consensus statement on the management of cytomegalovirus in solid organ transplant.<sup>39</sup> The statement indicated that "there are no clinical studies demonstrating that management decisions based on immunologic monitoring affect patient outcomes." Routine immunologic monitoring was not recommended.

#### U.S. Preventive Services Task Force Recommendations

Not applicable.

#### Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

## REFERENCES

1. Food and Drug Administration (FDA). Special 510(k): Device Modification 2010 (K101911). n.d.; [http://www.accessdata.fda.gov/cdrh\\_docs/reviews/K101911.pdf](http://www.accessdata.fda.gov/cdrh_docs/reviews/K101911.pdf). Accessed October 27, 2020.
2. Food and Drug Administration (FDA). Summary of Safety and Probable Benefit: Pleximmune. 2014; [http://www.accessdata.fda.gov/cdrh\\_docs/pdf13/H130004b.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf13/H130004b.pdf). Accessed October 27, 2020.
3. Ling X, Xiong J, Liang W, et al. Can immune cell function assay identify patients at risk of infection or rejection? A meta-analysis. *Transplantation*. Apr 15 2012;93(7):737-743. PMID 22357178
4. Rodrigo E, Lopez-Hoyos M, Corral M, et al. ImmuKnow((R)) as a diagnostic tool for predicting infection and acute rejection in adult liver transplant recipients: Systematic review and meta-analysis. *Liver Transpl*. Jun 27 2012;18(10):1245-1253. PMID 22740321
5. Rossano JW, Denfield SW, Kim JJ, et al. Assessment of the Cylex ImmuKnow cell function assay in pediatric heart transplant patients. *J Heart Lung Transplant*. Jan 2009;28(1):26-31. PMID 19134527
6. Wong MS, Boucek R, Kemna M, et al. Immune cell function assay in pediatric heart transplant recipients. *Pediatr Transplant*. Aug 2014;18(5):485-490. PMID 24930882

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

7. Ryan CM, Chaudhuri A, Concepcion W, et al. Immune cell function assay does not identify biopsy-proven pediatric renal allograft rejection or infection. *Pediatr Transplant*. Aug 2014;18(5):446-452. PMID 24930482
8. Wozniak LJ, Venick RS, Gordon Burroughs S, et al. Utility of an immune cell function assay to differentiate rejection from infectious enteritis in pediatric intestinal transplant recipients. *Clin Transplant*. Feb 2014;28(2):229-235. PMID 24433466
9. Liu W, Wang K, Zhao YH, et al. Clinical relevance of a CD4 + T cell immune function assay in the diagnosis of infection in pediatric living-donor liver transplantation. *Exp Ther Med*. Nov 2019; 18(5): 3823-3828. PMID 31602249
10. Nishikawa K, Mizuno S, Masui S, et al. Usefulness of monitoring cell-mediated immunity for predicting post- kidney transplantation viral infection. *Transplant Proc*. Mar 2014;46(2):552-555. PMID 24656010
11. Sageshima J, Ciancio G, Chen L, et al. Lack of clinical association and effect of peripheral WBC counts on immune cell function test in kidney transplant recipients with T-cell depleting induction and steroid-sparing maintenance therapy. *Transpl Immunol*. Mar 2014;30(2-3):88-92. PMID 24518158
12. Torio A, Fernandez EJ, Montes-Ares O, et al. Lack of association of immune cell function test with rejection in kidney transplantation. *Transplant Proc*. Jul-Aug 2011;43(6):2168-2170. PMID 21839223
13. Zhou H, Wu Z, Ma L, et al. Assessing immunologic function through CD4 T-lymphocyte adenosine triphosphate levels by ImmuKnow assay in Chinese patients following renal transplantation. *Transplant Proc*. Sep 2011;43(7):2574-2578. PMID 21911125
14. Huskey J, Gralla J, Wiseman AC. Single time point immune function assay (ImmuKnow) testing does not aid in the prediction of future opportunistic infections or acute rejection. *Clin J Am Soc Nephrol*. Feb 2011;6(2):423-429. PMID 21088287
15. Reinsmoen NL, Cornett KM, Kloehn R, et al. Pretransplant donor-specific and non-specific immune parameters associated with early acute rejection. *Transplantation*. Feb 15 2008;85(3):462-470. PMID 18301338
16. Serban G, Whittaker V, Fan J, et al. Significance of immune cell function monitoring in renal transplantation after Thymoglobulin induction therapy. *Hum Immunol*. Nov 2009;70(11):882-890. PMID 19664673
17. Libri I, Gnappi E, Zanelli P, et al. Trends in immune cell function assay and donor-specific HLA antibodies in kidney transplantation: A 3-year prospective study. *Am J Transplant*. Dec 2013;13(12):3215-3222. PMID 24266972
18. Myslik F, House AA, Yanko D, et al. Preoperative Cylex assay predicts rejection risk in patients with kidney transplant. *Clin Transplant*. May 2014;28(5):606-610. PMID 24628326
19. Quaglia M, Cena T, Fenoglio R, et al. Immune function assay (immunknow) drop over first 6 months after renal transplant: a predictor of opportunistic viral infections? *Transplant Proc*. Sep 2014;46(7):2220-2223. PMID 25242755
20. Wang XZ, Jin ZK, Tian XH, et al. Increased intracellular adenosine triphosphate level as an index to predict acute rejection in kidney transplant recipients. *Transpl Immunol*. Jan 2014;30(1):18-23. PMID 24211610
21. Weston MW, Rinde-Hoffman D, Lopez-Cepero M. Monitoring cell-mediated immunity during immunosuppression reduction in heart transplant recipients with severe systemic infections. *Clin Transplant*. Mar 2020; 34(3): e13809. PMID 32003048
22. Israeli M, Ben-Gal T, Yaari V, et al. Individualized immune monitoring of cardiac transplant recipients by noninvasive longitudinal cellular immunity tests. *Transplantation*. Apr 27 2010;89(8):968-976. PMID 20075792
23. Kobashigawa JA, Kiyosaki KK, Patel JK, et al. Benefit of immune monitoring in heart transplant patients using ATP production in activated lymphocytes. *J Heart Lung Transplant*. May 2010;29(5):504-508. PMID 20133166
24. Gupta S, Mitchell JD, Markham DW, et al. Utility of the Cylex assay in cardiac transplant recipients. *J Heart Lung Transplant*. Aug 2008;27(8):817-822. PMID 18656792
25. Shearer GM, Clerici M. In vitro analysis of cell-mediated immunity: clinical relevance. *Clin Chem*. Nov 1994;40(11 Pt 2):2162-2165. PMID 7955403
26. Cheng JW, Shi YH, Fan J, et al. An immune function assay predicts post-transplant recurrence in patients with hepatocellular carcinoma. *J Cancer Res Clin Oncol*. Oct 2011;137(10):1445-1453. PMID 21809031
27. Dong JY, Yin H, Li RD, et al. The relationship between adenosine triphosphate within CD4(+) T lymphocytes and acute rejection after liver transplantation. *Clin Transplant*. May-Jun 2011;25(3):E292-296. PMID 21470308
28. Hashimoto K, Miller C, Hirose K, et al. Measurement of CD4+ T-cell function in predicting allograft rejection and recurrent hepatitis C after liver transplantation. *Clin Transplant*. Sep-Oct 2010;24(5):701-708. PMID 20047619
29. Cabrera R, Ararat M, Soldevila-Pico C, et al. Using an immune functional assay to differentiate acute cellular rejection from recurrent hepatitis C in liver transplant patients. *Liver Transpl*. Feb 2009;15(2):216-222. PMID 19177434
30. Piloni D, Magni S, Oggionni T, et al. Clinical utility of CD4+ function assessment (ViraCor-IBT ImmuKnow test) in lung recipients. *Transpl Immunol*. Jul 2016;37:35-39. PMID 27095000
31. Husain S, Raza K, Pilewski JM, et al. Experience with immune monitoring in lung transplant recipients: correlation of low immune function with infection. *Transplantation*. Jun 27 2009;87(12):1852-1857. PMID 19543064
32. Bhorade SM, Janata K, Vigneswaran WT, et al. Cylex ImmuKnow assay levels are lower in lung transplant recipients with infection. *J Heart Lung Transplant*. Sep 2008;27(9):990-994. PMID 18765191
33. Shino MY, Weigt SS, Saggat R, et al. Usefulness of immune monitoring in lung transplantation using adenosine triphosphate production in activated lymphocytes. *J Heart Lung Transplant*. Sep 2012;31(9):996-1002. PMID 22884386

---

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

34. Ravaioli M, Neri F, Lazzarotto T, et al. Immunosuppression modifications based on an immune response assay: results of a randomized, controlled trial. *Transplantation*. Aug 2015;99(8):1625-1632. PMID 25757214
35. Manga K, Serban G, Schwartz J, et al. Increased adenosine triphosphate production by peripheral blood CD4+ cells in patients with hematologic malignancies treated with stem cell mobilization agents. *Hum Immunol*. Jul 2010;71(7):652-658. PMID 20381567
36. Gesundheit B, Budowski E, Israeli M, et al. Assessment of CD4 T-lymphocyte reactivity by the Cylex ImmuKnow assay in patients following allogeneic hematopoietic SCT. *Bone Marrow Transplant*. Mar 2010;45(3):527-533. PMID 19718067
37. Ashokkumar C, Talukdar A, Sun Q, et al. Allospecific CD154+ T cells associate with rejection risk after pediatric liver transplantation. *Am J Transplant*. Jan 2009;9(1):179-191. PMID 18976293
38. Ashokkumar C, Soltys K, Mazariegos G, et al. Predicting cellular rejection with a cell-based assay: preclinical evaluation in children. *Transplantation*. Jan 2017;101(1):131-140. PMID 26950712
39. Kotton CN, Kumar D, Caliendo AM, et al. International consensus guidelines on the management of cytomegalovirus in solid organ transplantation. *Transplantation*. Apr 15 2010;89(7):779-795. PMID 20224515

## POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
December 2011	New policy	
March 2012	Replace policy	Policy statement updated to read not medically necessary. References 10, 15-25 added. Previous references renumbered.
March 2013	Replace policy	Policy updated with literature review, two systematic reviews added and summary revised; references reordered; no change in policy statement.
March 2014	Replace policy	Policy updated with literature review, references 14-17 and 19 added; no change in policy statements.
March 2015	Replace policy	Policy updated with literature review, adding references 9-11, 13-14, 27-30 and 39-40; references 1 and 42 were updated. There are no changes to the policy statements.
June 2016	Replace policy	Policy updated with literature review through November 10, 2015; references 2 and 33-34 added. References on HIV, lupus nephritis deleted. Policy statements unchanged.
March 2018	Replace policy	lupus nephritis deleted. Policy statements unchanged. March 2018 Update Policy Policy updated with literature review through October 25, 2017; references 28, 29, 37 and 41 added. Policy statements corrected from "not medically necessary" to "investigational" based on FDA 510(k) and HDE approvals of assay tests.
March 2019	Replace policy	Policy updated with literature review through October 1, 2018; reference 38 added. Policy statements unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

---

<b>Date</b>	<b>Action</b>	<b>Description</b>
March 2020	Replace policy	Policy updated with literature review through October 14, 2019; no references added. Policy statements unchanged.
March 2021	Replace policy	Policy updated with literature review through October 27, 2020; references added. Policy statements unchanged.

---

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.