

Federal Employee Program® 1310 G Street, N.W. Washington, D.C. 20005 202.942.1000 Fax 202.942.1125

5.21.125

Section Prescription Drugs Effective Date: April 1, 2021

Subsection: Antineoplastic Agents Original Policy Date: January 11, 2019

Subject: Asparlas Page: 1 of 4

Last Review Date: March 12, 2021

# Asparlas

### **Description**

Asparlas (calaspargase pegol-mknl)

#### **Background**

Asparlas (calaspargase pegol-mknl) is an asparagine specific enzyme. L-asparaginase is an enzyme that catalyzes the conversion of the amino acid L-asparagine into aspartic acid and ammonia. The pharmacological effect of Asparlas is thought to be based on selective killing of leukemic cells due to depletion of plasma L-asparagine. Leukemic cells with low expression of asparagine synthetase have a reduced ability to synthesize L-asparagine, and therefore depend on an exogenous source of L-asparagine for survival (1).

The toxicity and efficacy of asparaginase is being studied in adult patients with acute lymphoblastic leukemia (ALL) (2).

#### **Regulatory Status**

FDA-approved indication: Asparlas is indicated as a component of a multi-agent chemotherapeutic regimen for the treatment of acute lymphoblastic leukemia (ALL) in pediatric and young adult patients age 1 month to 21 years (1).

Asparlas is contraindicated in patients with a history of pancreatitis, thrombosis, hemorrhagic events or anaphylaxis reaction with prior L-asparaginase therapy. Discontinue Asparlas in the event of serious hypersensitivity reactions, including anaphylaxis, and severe or hemorrhagic pancreatitis. Glucose intolerance can occur. Bilirubin, transaminases, and glucose should be monitored at least weekly until recovery from the cycle of therapy (1).

# 5.21.125

Section: Prescription Drugs Effective Date: April 1, 2021

Subsection: Antineoplastic Agents Original Policy Date: January 11, 2019

Subject: Asparlas Page: 2 of 4

Due to the risk of serious allergic reactions (such as life-threatening anaphylaxis), Asparlas should be administered in a clinical setting with resuscitation equipment and other agents necessary to treat anaphylaxis and patients should be observed for 1 hour after administration (1).

The safety and effectiveness of Asparlas in patients 1 month to 21 years of age have been established (1).

### Related policies

Erwinaze, Oncaspar

## Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Asparlas may be considered **medically necessary** in patients age 1 month to 21 years of age with acute lymphoblastic leukemia (ALL) and if the conditions indicated below are met.

Asparlas may be considered **investigational** in patients not between the ages of 1 month and 21 years and for all other indications.

## **Prior-Approval Requirements**

Age 1 month to 21 years of age

### **Diagnosis**

Patient must have the following:

- 1. Acute lymphoblastic leukemia (ALL)
  - a. Prescriber agrees to monitor bilirubin, liver function tests (LFTs), and glucose

# Prior – Approval Renewal Requirements

Same as above

# 5.21.125

Section: Prescription Drugs Effective Date: April 1, 2021

Subsection: Antineoplastic Agents Original Policy Date: January 11, 2019

Subject: Asparlas Page: 3 of 4

## **Policy Guidelines**

### **Pre - PA Allowance**

None

## **Prior - Approval Limits**

**Duration** 12 months

## Prior - Approval Renewal Limits

Same as above

### Rationale

### **Summary**

Asparlas (calaspargase pegol-mknl) is an asparagine specific enzyme. L-asparaginase is an enzyme that catalyzes the conversion of the amino acid L-asparagine into aspartic acid and ammonia. The pharmacological effect of Asparlas is thought to be based on selective killing of leukemic cells due to depletion of plasma L-asparagine. Leukemic cells with low expression of asparagine synthetase have a reduced ability to synthesize L-asparagine, and therefore depend on an exogenous source of L-asparagine for survival. The safety and effectiveness of Asparlas in patients 1 month to 21 years of age have been established (1).

Prior approval is required to ensure the safe, clinically appropriate and cost effective use of Asparlas while maintaining optimal therapeutic outcomes.

#### References

- 1. Asparlas [package insert]. Boston, MA: Servier Pharmaceuticals LLC.; June 2020.
- Koprivnikar J, McCloskey J, Faderl S. Safety, efficacy, and clinical utility of asparaginase in the treatment of adult patients with acute lymphoblastic leukemia. Onco Targets Ther. 2017; 10:1413-1422.

Policy History	
Date	Action
January 2019	Addition to PA
March 2019	Annual review
September 2019	Annual review, Added Onco Targets and Therapy reference per SME

# 5.21.125

Section: Prescription Drugs Effective Date: April 1, 2021

Subsection: Antineoplastic Agents Original Policy Date: January 11, 2019

Subject: Asparlas Page: 4 of 4

June 2020 Annual review and reference update

March 2021 Annual editorial review and reference update

Keywords

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on March 12, 2021 and is effective on April 1, 2021.