



5.30.65

Section:	Prescription Drugs	Effective Date:	April 1, 2021
Subsection:	Endocrine and Metabolic Drugs	Original Policy Date:	March 6, 2020
Subject:	Bynfezia	Page:	1 of 4

Last Review Date: March 12, 2021

Bynfezia

Description

Bynfezia (octreotide acetate) pen

Background

Bynfezia (octreotide acetate) is a subcutaneous injection for the treatment of acromegaly, diarrhea or flushing episodes that are associated with metastatic carcinoid tumors, and diarrhea that is associated with vasoactive intestinal peptide (VIP)-secreting tumors. Acromegaly is a rare and debilitating endocrine disorder caused by excess production of growth hormone (GH) and insulin-like growth factor-1 (IGF-1). Metastatic carcinoid tumors are found along the gastrointestinal (GI) tract and release too much serotonin into the body, while VIP-secreting tumors cause increased secretions from the intestines. Bynfezia mimics natural somatostatin by inhibiting the secretion of growth hormone, glucagon, insulin, serotonin, gastrin, VIP, secretin, motilin, and pancreatic polypeptide (1).

Regulatory Status

FDA-approved indication: Bynfezia is a somatostatin analogue indicated for: (1)

1. Reduction of growth hormone (GH) and insulin-like growth factor 1 (IGF-1) [somatomedin C] in adult patients with acromegaly who have had inadequate response to or cannot be treated with surgical resection, pituitary irradiation, and bromocriptine mesylate at maximally tolerated doses
2. Treatment of severe diarrhea/flushing episodes associated with metastatic carcinoid tumors in adult patients
3. Treatment of profuse watery diarrhea associated with vasoactive intestinal peptide tumors (VIPomas) in adult patients

Section:	Prescription Drugs	Effective Date:	April 1, 2021
Subsection:	Endocrine and Metabolic Drugs	Original Policy Date:	March 6, 2020
Subject:	Bynfezia	Page:	2 of 4

Limitations of Use: (1)

- In patients with acromegaly, the effect of Bynfezia on improvement in clinical signs and symptoms, reduction in tumor size and rate of growth, has not been determined.
- In patients with carcinoid syndrome and VIPomas, the effect of Bynfezia on tumor size, rate of growth, and development of metastases has not been determined.

The safety and effectiveness of Bynfezia in pediatric patients less than 18 years of age have not been established (1).

Related policies

Mycapssa, Sandostatin LAR, Signifor LAR, Somatuline Depot

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Bynfezia may be considered **medically necessary** in patients 18 years of age or older with acromegaly; severe diarrhea or flushing episodes associated with metastatic carcinoid tumors; severe diarrhea associated with VIP-secreting tumors and if the conditions indicated below are met.

Bynfezia is considered **investigational** in patients less than 18 years of age and for all other indications.

Prior-Approval Requirements

Age 18 years of age and older

Diagnoses

Patient must have **ONE** of the following:

1. Acromegaly
 - a. Inadequate response or patient is **NOT** a candidate for **ALL** of the following:
 - i. Surgery resection
 - ii. Pituitary irradiation
 - iii. Bromocriptine mesylate

Section:	Prescription Drugs	Effective Date:	April 1, 2021
Subsection:	Endocrine and Metabolic Drugs	Original Policy Date:	March 6, 2020
Subject:	Bynfezia	Page:	3 of 4

2. Severe diarrhea or flushing episodes associated with metastatic carcinoid tumor(s)
3. Profuse watery diarrhea associated with VIP-secreting tumor(s)

AND the following:

- a. Must have documented reason for requiring special injection device such as: lack of dexterity, visual acuity issues

Prior – Approval *Renewal* Requirements

Age 18 years of age or older

Diagnoses

Patient must have **ONE** of the following:

1. Acromegaly
2. Severe diarrhea or flushing episodes associated with metastatic carcinoid tumor(s)
3. Profuse watery diarrhea associated with VIP-secreting tumor(s)

Policy Guidelines

Pre - PA Allowance

None

Prior - Approval Limits

Duration 12 months

Prior – Approval *Renewal* Limits

Same as above

Rationale

Summary

Section:	Prescription Drugs	Effective Date:	April 1, 2021
Subsection:	Endocrine and Metabolic Drugs	Original Policy Date:	March 6, 2020
Subject:	Bynfezia	Page:	4 of 4

Bynfezia (octreotide acetate) is a subcutaneous injection for the treatment of acromegaly, diarrhea or flushing episodes that are associated with metastatic carcinoid tumors, and diarrhea that is associated with vasoactive intestinal peptide (VIP)-secreting tumors. Acromegaly is a rare and debilitating endocrine disorder caused by excess production of growth hormone (GH) and insulin-like growth factor-1 (IGF-1). Metastatic carcinoid tumors are found along the gastrointestinal (GI) tract and release too much serotonin into the body, while VIP-secreting tumors cause increased secretions from the intestines. Bynfezia mimics natural somatostatin by inhibiting the secretion of growth hormone, glucagon, insulin, serotonin, gastrin, VIP, secretin, motilin, and pancreatic polypeptide. The safety and effectiveness of Bynfezia in pediatric patients less than 18 years of age have not been established (1).

Prior approval is required to ensure the safe, clinically appropriate and cost-effective use of Bynfezia while maintaining optimal therapeutic outcomes.

References

1. Bynfezia [package insert]. Cranbury, NJ: Sun Pharmaceutical Industries, Inc.; January 2020.

Policy History

Date	Action
March 2020	Addition to PA
June 2020	Annual review
September 2020	Annual review
December 2020	Annual review
March 2021	Annual review

Keywords

This policy was approved by the FEP® Pharmacy Medical Policy Committee on March 12, 2021 and is effective on April 1, 2021.