

5.30.030

Section:	Prescription Drugs	Effective Date:	January 1, 2023
Subsection:	Endocrine and Metabolic Drugs	Original Policy Date:	January 1, 2014
Subject:	Buphenyl Pheburane	Page:	1 of 4

Last Review Date: December 2, 2022

Buphenyl Pheburane

Description

Buphenyl tablet, powder for solution (sodium phenylbutyrate)

Pheburane oral pellets (sodium phenylbutyrate)

Background

Urea cycle disorders include a group of diseases, each having a specific liver enzyme deficiency. Because they are inherited, other family members may be affected. These disorders vary in severity and may be first detected at various ages, from newborn infants to adults. They lead to increased amounts of ammonia in the blood, which may cause disturbed brain function and severe brain damage (1-2).

Buphenyl is available both as tablets and a powder for oral use (via mouth, gastrostomy, or nasogastric tube) with meals or feedings and helps dispose of ammonia in the body. Pheburane is available as a taste-masked oral pellet. Sodium phenylbutyrate is intended for patients who have UCD that cannot be managed by a protein-restricted diet or amino acid supplements alone. Buphenyl and Pheburane must be used with a protein-restricted diet and, in some cases, dietary supplements (1).

Regulatory Status

FDA-approved indication: Buphenyl and Pheburane are indicated as adjunctive therapy in the chronic management of patients with urea cycle disorders involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS) (1-2).

Section:	Prescription Drugs	Effective Date:	January 1, 2023
Subsection:	Endocrine and Metabolic Drugs	Original Policy Date:	January 1, 2014
Subject:	Buphenyl Pheburane	Page:	2 of 4

Limitations of Use:

Buphenyl and Pheburane are not indicated for treatment of acute hyperammonemia in patients with UCDs (1-2).

Caution should be used when using haloperidol and valproic acid. Buphenyl and Pheburane should be used with great care, if at all, in patients with congestive heart failure or severe renal insufficiency and in clinical states in which there is sodium retention with edema. Probenecid may inhibit renal transport of Buphenyl and Pheburane. Use of corticosteroids may cause the breakdown of body protein and increase plasma ammonia levels. The use of tablets for neonates, infants, and children under the weight of 20kg is not recommended (1-2).

Related policies

Ravicti

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Buphenyl and Pheburane may be considered **medically necessary** in patients with the diagnosis of urea cycle disorders (UCDs) involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS) and if the conditions indicated below are met.

Buphenyl and Pheburane may be considered **investigational** for all other indications.

Prior-Approval Requirements

Diagnosis

Patient must have the following:

Urea cycle disorders (UCDs) involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS)

AND ALL of the following:

- a. Failure to control ammonia level with dietary restrictions and/or amino acid supplementation

Section:	Prescription Drugs	Effective Date:	January 1, 2023
Subsection:	Endocrine and Metabolic Drugs	Original Policy Date:	January 1, 2014
Subject:	Buphenyl Pheburane	Page:	3 of 4

- b. Prescribing physician should be experienced in the management of UCDs
- c. Prescriber agrees to monitor electrolytes at baseline and as clinically indicated
- d. Must be used with dietary protein restrictions
- e. **NO** acute hyperammonemic encephalopathy

Prior – Approval *Renewal* Requirements

Diagnosis

Patient must have the following:

Urea cycle disorders (UCDs) involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS)

AND ALL of the following:

- a. Prescriber agrees to monitor electrolytes at baseline and as clinically indicated
- b. Must be used with dietary protein restrictions

Policy Guidelines

Pre - PA Allowance

None

Prior - Approval Limits

Duration 2 years

Prior – Approval *Renewal* Limits

Same as above

Rationale

Summary

Buphenyl and Pheburane are indicated as adjunctive therapy in the chronic management of patients with urea cycle disorders involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS). Patients with urea cycle disorders should not take valproic acid, haloperidol, or steroids as these drugs

Section:	Prescription Drugs	Effective Date:	January 1, 2023
Subsection:	Endocrine and Metabolic Drugs	Original Policy Date:	January 1, 2014
Subject:	Buphenyl Pheburane	Page:	4 of 4

have been reported to increase blood ammonia levels, and probenecid may affect the kidneys' excretion. Use with great care, if at all, in patients with congestive heart failure or severe renal insufficiency, and in clinical states where there is sodium retention with edema. Use caution when administering to patients with hepatic or renal insufficiency or inborn errors of beta oxidation. The safety or efficacy of doses in excess of 20 grams (40 tablets) per day has not been established (1).

Prior approval is required to ensure the safe, clinically appropriate, and cost-effective use of Buphenyl and Pheburane while maintaining optimal therapeutic outcomes.

References

1. Buphenyl [package insert]. Lake Forest, IL: Horizon Therapeutics, Inc. May 2021.
2. Pheburane [package insert]. Bryn Mawr, PA: Medunik USA, Inc. June 2022.

Policy History

Date	Action
January 2014	New addition to PA
March 2014	Annual review
March 2015	Annual criteria review and reference update
September 2016	Annual editorial review and reference update Policy code changed from 5.08.30 to 5.30.30
December 2017	Annual editorial review
November 2018	Annual review
December 2019	Annual editorial review and reference update. Changed approval duration from lifetime to 2 years
December 2020	Annual review and reference update
March 2021	Annual review
March 2022	Annual review and reference update
September 2022	Addition of Pheburane to policy
December 2022	Annual review. Per SME, added requirement for prescribers to monitor electrolytes at baseline and as clinically indicated

Keywords

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on December 2, 2022 and is effective on January 1, 2023.