



FEP Medical Policy Manual

FEP 7.01.151 Prostatic Urethral Lift

Annual Effective Policy Date: January 1, 2024

Original Policy Date: January 2015

Related Policies:

7.01.175 - Temporarily Implanted Nitinol Device (iTind) for Benign Prostatic Hyperplasia

Prostatic Urethral Lift

Description

Description

Benign prostatic hyperplasia (BPH) is a common condition in older individuals that can lead to increased urinary frequency, an urgency to urinate, a hesitancy to urinate, nocturia, and a weak stream when urinating. The prostatic urethral lift (PUL) procedure involves the insertion of one or more permanent implants into the prostate, which retracts prostatic tissue and maintains an expanded urethral lumen.

OBJECTIVE

The objective of this evidence review is to determine whether prostatic urethral lift improves the net health outcome in individuals with benign prostatic hyperplasia.

POLICY STATEMENT

Use of prostatic urethral lift in individuals with moderate-to-severe lower urinary tract obstruction due to benign prostatic hyperplasia may be considered **medically necessary** when all of the following criteria are met:

- The individual has persistent or progressive lower urinary tract symptoms despite medical therapy (α_1 -adrenergic antagonists maximally titrated, 5 α -reductase inhibitors, or combination medication therapy maximally titrated) over a trial period of no less than 6 months, or is unable to tolerate medical therapy; AND,
- Prostate gland volume is \leq 80 mL; AND,
- Prostate anatomy demonstrates normal bladder neck without an obstructive or protruding median lobe; AND,
- Individual does not have urinary retention related to conditions other than benign prostatic hyperplasia, urinary tract infection, or recent prostatitis (within past year); AND,
- Individual has had appropriate testing to exclude diagnosis of prostate cancer; AND,
- Individual does not have a known allergy to nickel, titanium or stainless steel.

Use of prostatic urethral lift in other situations, including repeat procedures, is considered **investigational**.

POLICY GUIDELINES

Use of temporarily implanted nitinol devices for benign prostatic hyperplasia is addressed separately in evidence review 7.01.175.

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

FDA REGULATORY STATUS

One implantable transprostatic tissue retractor system has been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. In 2013, the NeoTract UroLift System UL400 (NeoTract) was cleared (after receiving clearance through the FDA's de novo classification process in March 2013; K130651/DEN130023). In 2016, the FDA determined that the UL500 was substantially equivalent to existing devices (UL400) for the treatment of symptoms of urinary flow obstruction secondary to BPH in individuals ages 50 years and older. In 2017, the FDA expanded the indication for the UL400 and UL500 to include *lateral and median* lobe hyperplasia in men 45 years or older. An additional clearance in 2019 (K193269) modified an existing contraindication for use from men with a prostate volume of $>$ 80 cc to men with a prostate volume of $>$ 100 cc. FDA product code: PEW.

RATIONALE

Summary of Evidence

For individuals who have lower urinary tract obstruction symptoms due to benign prostatic hyperplasia (BPH) who do not have sufficient response to medical therapy or are experiencing significant side effects with medical therapy and receive a prostatic urethral lift (PUL), the evidence includes systematic reviews, randomized controlled trials (RCTs), and nonrandomized studies. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. One RCT, the BPH6 study, compared the PUL procedure with transurethral resection of the prostate (TURP) and reported that the PUL procedure was noninferior for the study's composite endpoint, which required concurrent fulfillment of 6 independently validated measures of symptoms, safety, and sexual health. While TURP was superior to PUL in managing lower urinary tract symptoms, PUL did provide significant symptom improvement over 2 years. Prostatic urethral lift was further superior to TURP in preserving ejaculatory function. These findings were corroborated by another RCT (the LIFT study), which compared PUL with sham control. Patients underwent washout of BPH medications before enrollment. LIFT reported that patients with the PUL procedure, compared with patients who had sham surgery and no BPH medication, had greater improvements in lower urinary tract symptoms without worsened sexual function at 3 months. After 3 months, patients were given the option to have PUL surgery; 80% of the patients with sham procedures chose that option. Publications from this trial reported these findings were preserved in a subset of patients over 3 to 5 years; however, a high number of patients were either excluded or lost to follow-up during this time. The BPH6 and LIFT RCTs included men with a prostate volume up to 80 cm³ and excluded men with median lobe obstruction. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have lower urinary tract obstruction symptoms due to BPH who have had a prior PUL procedure who are treated with a repeat PUL, the evidence includes long-term follow-up data from the LIFT study, a systematic review, and reports on care setting real world experience. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. Clinical data on the occurrence of repeat PUL, and consensus on clinically relevant definitions of retreatment/reintervention and subsequent outcomes are lacking. The 5 year surgical reintervention rate in the LIFT study was reported as 13.6%, while a meta-analysis concluded that the surgical reintervention rate following PUL is 6% per year. An analysis of clinical care setting real world experience reported the overall retreatment rate at 1 and 2 years to be 5.2% (95% confidence interval [CI], 4.2 to 6.1) and 11.9% (95% CI, 10.1 to 13.6), respectively, following an initial PUL. A retrospective healthcare system database analysis of endoscopic procedures for BPH found that patients treated with PUL were almost twice as likely to be retreated at 2-year follow-up compared to those receiving TURP (odds ratio [OR], 1.78; p<.01). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in "Supplemental Information" if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

National Institute for Health and Care Excellence

In 2014, the National Institute for Health and Care Excellence published guidance on urethral lift implants to treat lower urinary tract symptoms (LUTS) secondary to benign prostatic hyperplasia (BPH).⁴⁹ The guidance stated:

"Current evidence on the efficacy and safety of insertion of prostatic urethral lift implants to treat lower urinary tract symptoms secondary to benign prostatic hyperplasia is adequate to support the use of this procedure."

In 2021, the National Institute for Health and Care Excellence published updated guidance on the use of UroLift for treating LUTS of BPH.⁵⁰ The guidance stated: "the UroLift system relieves lower urinary tract symptoms, avoids risk to sexual function, and improves quality of life " and "the UroLift system should be considered as an alternative to transurethral resection of the prostate (TURP) and holmium laser enucleation of the prostate (HoLEP). It can be done as a day-case or outpatient procedure for people aged 50 and older with a prostate volume between 30 and 80 mL."

American Urological Association

In 2018, the American Urological Association published guidelines on the surgical management of LUTS attributed to BPH; the 2018 guidelines were most recently amended in 2021.⁷ The guidelines made the following recommendations and statements regarding prostatic urethral lift (PUL).

- "PUL may be offered as an option for patients with LUTS [lower urinary tract symptoms] /BPH [benign prostatic hyperplasia] provided prostate volume 30-80cc and verified absence of an obstructive middle lobe "
 - "Moderate Recommendation; Evidence Level: Grade C indicating "Benefits > Risks/Burdens (or vice versa); Net benefit (or net harm) appears moderate. Applies to most patients in most circumstances but better evidence is likely to change confidence"
- "PUL may be offered as a treatment option to eligible patients who desire preservation of erectile and ejaculatory function."
 - "Conditional Recommendation; Evidence Level: Grade C indicating "Risks/Burdens unclear; Alternative strategies may be equally reasonable. Better evidence likely to change confidence"
- "Clinicians should inform patients of the possibility of treatment failure and the need for additional or secondary treatments when considering surgical and minimally-invasive treatments for LUTS/BPH."
- "Surgery is recommended for patients who have renal insufficiency secondary to BPH, refractory urinary retention secondary to BPH, recurrent urinary tract infections (UTIs), recurrent bladder stones or gross hematuria due to BPH, and/or with LUTS/BPH refractory to or unwilling to use other therapies."

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

REFERENCES

1. Sarma AV, Wei JT. Clinical practice. Benign prostatic hyperplasia and lower urinary tract symptoms. *N Engl J Med*. Jul 19 2012; 367(3): 248-57. PMID 22808960
2. Barry MJ, Fowler FJ, O'Leary MP, et al. Measuring disease-specific health status in men with benign prostatic hyperplasia. Measurement Committee of The American Urological Association. *Med Care*. Apr 1995; 33(4 Suppl): AS145-55. PMID 7536866
3. O'leary MP. Validity of the "bother score" in the evaluation and treatment of symptomatic benign prostatic hyperplasia. *Rev Urol*. 2005; 7(1): 1-10. PMID 16985801
4. Djavan B, Marberger M. A meta-analysis on the efficacy and tolerability of alpha1-adrenoceptor antagonists in patients with lower urinary tract symptoms suggestive of benign prostatic obstruction. *Eur Urol*. 1999; 36(1): 1-13. PMID 10364649
5. Foster HE, Barry MJ, Dahm P, et al. Surgical Management of Lower Urinary Tract Symptoms Attributed to Benign Prostatic Hyperplasia: AUA Guideline. *J Urol*. Sep 2018; 200(3): 612-619. PMID 29775639
6. Reich O, Gratzke C, Bachmann A, et al. Morbidity, mortality and early outcome of transurethral resection of the prostate: a prospective multicenter evaluation of 10,654 patients. *J Urol*. Jul 2008; 180(1): 246-9. PMID 18499179
7. Lerner LB, McVary KT, Barry MJ, et al. Management of Lower Urinary Tract Symptoms Attributed to Benign Prostatic Hyperplasia: AUA GUIDELINE PART II-Surgical Evaluation and Treatment. *J Urol*. Oct 2021; 206(4): 818-826. PMID 34384236
8. Sundaram D, Sankaran PK, Raghunath G, et al. Correlation of Prostate Gland Size and Uroflowmetry in Patients with Lower Urinary Tract Symptoms. *J Clin Diagn Res*. May 2017; 11(5): AC01-AC04. PMID 28658743
9. Rosen RC, Catania JA, Althof SE, et al. Development and validation of four-item version of Male Sexual Health Questionnaire to assess ejaculatory dysfunction. *Urology*. May 2007; 69(5): 805-9. PMID 17482908
10. Cappelleri JC, Rosen RC. The Sexual Health Inventory for Men (SHIM): a 5-year review of research and clinical experience. *Int J Impot Res*. 2005; 17(4): 307-19. PMID 15875061
11. Snksen J, Barber NJ, Speakman MJ, et al. Prospective, randomized, multinational study of prostatic urethral lift versus transurethral resection of the prostate: 12-month results from the BPH6 study. *Eur Urol*. Oct 2015; 68(4): 643-52. PMID 25937539
12. Barry MJ, Williford WO, Chang Y, et al. Benign prostatic hyperplasia specific health status measures in clinical research: how much change in the American Urological Association symptom index and the benign prostatic hyperplasia impact index is perceptible to patients?. *J Urol*. Nov

- 1995; 154(5): 1770-4. PMID 7563343
13. Roehrborn CG, Wilson TH, Black LK. Quantifying the contribution of symptom improvement to satisfaction of men with moderate to severe benign prostatic hyperplasia: 4-year data from the CombAT trial. *J Urol.* May 2012; 187(5): 1732-8. PMID 22425127
 14. McVary KT, Roehrborn CG, Avins AL, et al. American Urological Association Guideline: Management of Benign Prostatic Hyperplasia (BPH). 2010 (affirmed 2014); [http://www.auanet.org/guidelines/benign-prostatic-hyperplasia-\(2010-reviewed-and-validity-confirmed-2014\)](http://www.auanet.org/guidelines/benign-prostatic-hyperplasia-(2010-reviewed-and-validity-confirmed-2014)). Accessed July 7, 2023.
 15. Perera M, Roberts MJ, Doi SA, et al. Prostatic urethral lift improves urinary symptoms and flow while preserving sexual function for men with benign prostatic hyperplasia: a systematic review and meta-analysis. *Eur Urol.* Apr 2015; 67(4): 704-13. PMID 25466940
 16. Cantwell AL, Bogache WK, Richardson SF, et al. Multicentre prospective crossover study of the 'prostatic urethral lift' for the treatment of lower urinary tract symptoms secondary to benign prostatic hyperplasia. *BJU Int.* Apr 2014; 113(4): 615-22. PMID 24765680
 17. Shore N, Freedman S, Gange S, et al. Prospective multi-center study elucidating patient experience after prostatic urethral lift. *Can J Urol.* Feb 2014; 21(1): 7094-101. PMID 24529008
 18. McNicholas TA, Woo HH, Chin PT, et al. Minimally invasive prostatic urethral lift: surgical technique and multinational experience. *Eur Urol.* Aug 2013; 64(2): 292-9. PMID 23357348
 19. Chin PT, Bolton DM, Jack G, et al. Prostatic urethral lift: two-year results after treatment for lower urinary tract symptoms secondary to benign prostatic hyperplasia. *Urology.* Jan 2012; 79(1): 5-11. PMID 22202539
 20. Woo HH, Bolton DM, Laborde E, et al. Preservation of sexual function with the prostatic urethral lift: a novel treatment for lower urinary tract symptoms secondary to benign prostatic hyperplasia. *J Sex Med.* Feb 2012; 9(2): 568-75. PMID 22172161
 21. Woo HH, Chin PT, McNicholas TA, et al. Safety and feasibility of the prostatic urethral lift: a novel, minimally invasive treatment for lower urinary tract symptoms (LUTS) secondary to benign prostatic hyperplasia (BPH). *BJU Int.* Jul 2011; 108(1): 82-8. PMID 21554526
 22. Hoffman RM, Monga M, Elliott SP, et al. Microwave thermotherapy for benign prostatic hyperplasia. *Cochrane Database Syst Rev.* Sep 12 2012; (9): CD004135. PMID 22972068
 23. Roehrborn CG, Gange SN, Shore ND, et al. The prostatic urethral lift for the treatment of lower urinary tract symptoms associated with prostate enlargement due to benign prostatic hyperplasia: the L.I.F.T. Study. *J Urol.* Dec 2013; 190(6): 2161-7. PMID 23764081
 24. Shore N. A Review of the Prostatic Urethral Lift for Lower Urinary Tract Symptoms: Symptom Relief, Flow Improvement, and Preservation of Sexual Function in Men With Benign Prostatic Hyperplasia. *Curr Bladder Dysfunct Rep.* 2015; 10(2): 186-192. PMID 25984251
 25. Roehrborn CG, Rukstalis DB, Barkin J, et al. Three year results of the prostatic urethral L.I.F.T. study. *Can J Urol.* Jun 2015; 22(3): 7772-82. PMID 26068624
 26. McVary KT, Gange SN, Shore ND, et al. Treatment of LUTS secondary to BPH while preserving sexual function: randomized controlled study of prostatic urethral lift. *J Sex Med.* Jan 2014; 11(1): 279-87. PMID 24119101
 27. Garrido Abad P, Coloma Del Peso A, Sinues Ojas B, et al. [Urolift, a new minimally invasive treatment for patients with low urinary tract symptoms secondary to BPH. Preliminary results]. *Arch Esp Urol.* 2013; 66(6): 584-91. PMID 23985459
 28. Jones P, Rajkumar GN, Rai BP, et al. Medium-term Outcomes of Urolift (Minimum 12 Months Follow-up): Evidence From a Systematic Review. *Urology.* Nov 2016; 97: 20-24. PMID 27208817
 29. Bozkurt A, Karabakan M, Keskin E, et al. Prostatic Urethral Lift: A New Minimally Invasive Treatment for Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia. *Urol Int.* 2016; 96(2): 202-6. PMID 26613256
 30. Ray A, Morgan H, Wilkes A, et al. The Urolift System for the Treatment of Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia: A NICE Medical Technology Guidance. *Appl Health Econ Health Policy.* Oct 2016; 14(5): 515-26. PMID 26832146
 31. Tanneru K, Gautam S, Norez D, et al. Meta-analysis and systematic review of intermediate-term follow-up of prostatic urethral lift for benign prostatic hyperplasia. *Int Urol Nephrol.* Jun 2020; 52(6): 999-1008. PMID 32065331
 32. Rukstalis D, Rashid P, Bogache WK, et al. 24-month durability after crossover to the prostatic urethral lift from randomised, blinded sham. *BJU Int.* Oct 2016; 118 Suppl 3: 14-22. PMID 27684483
 33. Sievert KD, Schonhaler M, Berges R, et al. Minimally invasive prostatic urethral lift (PUL) efficacious in TURP candidates: a multicenter German evaluation after 2 years. *World J Urol.* Jul 2019; 37(7): 1353-1360. PMID 30283994
 34. Jung JH, Reddy B, McCutcheon KA, et al. Prostatic urethral lift for the treatment of lower urinary tract symptoms in men with benign prostatic hyperplasia. *Cochrane Database Syst Rev.* May 25 2019; 5(5): CD012832. PMID 31128077
 35. Gratzke C, Barber N, Speakman MJ, et al. Prostatic urethral lift vs transurethral resection of the prostate: 2-year results of the BPH6 prospective, multicentre, randomized study. *BJU Int.* May 2017; 119(5): 767-775. PMID 27862831
 36. Franco JVA, Jung JH, Imamura M, et al. Minimally invasive treatments for benign prostatic hyperplasia: a Cochrane network meta-analysis. *BJU Int.* Aug 2022; 130(2): 142-156. PMID 34820997
 37. Roehrborn CG. Prostatic Urethral Lift: A Unique Minimally Invasive Surgical Treatment of Male Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia. *Urol Clin North Am.* Aug 2016; 43(3): 357-69. PMID 27476128
 38. Roehrborn CG, Barkin J, Gange SN, et al. Five year results of the prospective randomized controlled prostatic urethral L.I.F.T. study. *Can J Urol.* Jun 2017; 24(3): 8802-8813. PMID 28646935
 39. Rukstalis D, Grier D, Stroup SP, et al. Prostatic Urethral Lift (PUL) for obstructive median lobes: 12 month results of the MedLift Study. *Prostate Cancer Prostatic Dis.* Sep 2019; 22(3): 411-419. PMID 30542055
 40. Eure G, Rukstalis D, Roehrborn C. Prostatic Urethral Lift for Obstructive Median Lobes: Consistent Results Across Controlled Trial and Real-World Settings. *J Endourol.* Jan 2023; 37(1): 50-59. PMID 35876440
 41. Shah BB, Tayon K, Madiraju S, et al. Prostatic Urethral Lift: Does Size Matter?. *J Endourol.* Jul 2018; 32(7): 635-638. PMID 29631445
 42. Eure G, Gange S, Walter P, et al. Real-World Evidence of Prostatic Urethral Lift Confirms Pivotal Clinical Study Results: 2-Year Outcomes of a Retrospective Multicenter Study. *J Endourol.* Jul 2019; 33(7): 576-584. PMID 31115257

43. Kaplan SA. Surgical Reintervention Rate after Prostatic Urethral Lift: Systematic Review and Meta-Analysis Involving over 2,000 Patients. Letter. J Urol. Mar 2021; 205(3): 939-940. PMID 33393811
44. Miller LE, Chughtai B, Dornbier RA, et al. Surgical Reintervention Rate after Prostatic Urethral Lift: Systematic Review and Meta-Analysis Involving over 2,000 Patients. Reply. J Urol. Mar 2021; 205(3): 940-941. PMID 33393812
45. McVary KT, Kaplan SA. A Tower of Babel in Today's Urology: Disagreement in Concepts and Definitions of Lower Urinary Tract Symptoms/Benign Prostatic Hyperplasia Re-Treatment. J Urol. Aug 2020; 204(2): 213-214. PMID 32469261
46. Miller LE, Chughtai B, Dornbier RA, et al. Surgical Reintervention Rate after Prostatic Urethral Lift: Systematic Review and Meta-Analysis Involving over 2,000 Patients. J Urol. Nov 2020; 204(5): 1019-1026. PMID 32396049
47. Gaffney CD, Basourakos SP, Al Hussein Al Awamih B, et al. Adoption, Safety, and Retreatment Rates of Prostatic Urethral Lift for Benign Prostatic Enlargement. J Urol. Aug 2021; 206(2): 409-415. PMID 33793296
48. Page T, Veeratterapillay R, Keltie K, et al. Prostatic urethral lift (UroLift): a real-world analysis of outcomes using hospital episodes statistics. BMC Urol. Apr 07 2021; 21(1): 55. PMID 33827525
49. National Institute for Health and Care Excellence (NICE). Insertion of prostatic urethral lift implants to treat lower urinary tract symptoms secondary to benign prostatic hyperplasia [IPG475]. 2014; <https://www.nice.org.uk/guidance/ipg475/chapter/1-recommendations>. Accessed July 7, 2023.
50. National Institute for Health and Care Excellence (NICE). UroLift for treating lower urinary tract symptoms of benign prostatic hyperplasia [MTG58]. 2021; <https://www.nice.org.uk/guidance/MTG58>. Accessed July 7, 2023.

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
December 2015	New policy	
December 2016	Replace policy	Policy updated with literature review through July 10, 2016; references 11, 21-22, 24, 26, and 28 added. Policy statement unchanged.
March 2018	Replace policy	Policy updated with literature review through October 9, 2017; references 4-5, 24, 28-29, and 31 added. Use of prostatic urethral lift in individuals with moderate to severe lower urinary tract obstruction due to benign prostatic hyperplasia may be considered medically necessary when all of the specified criteria are met.
December 2018	Replace policy	Policy updated with literature review through June 4, 2018; references 6 and 37-38 added. The medically necessary statement related to not being a surgical candidate for TURP was removed.
December 2019	Replace policy	Policy updated with literature review through June 19, 2019; references added. The medically necessary (MN) statement was updated to remove the clause 'Patient does not have prostate-specific antigen level - 3 ng/mL' from the fifth criterion. The MN criterion regarding nickle allergy was expanded to include titanium and stainless steel.
December 2020	Replace policy	Policy updated with literature review through July 2, 2020; references added. Repeat procedures explicitly added to the investigational policy statement; statements otherwise unchanged.
December 2021	Replace policy	Policy updated with literature review through July 7, 2021; references added. Policy statements unchanged.
December 2022	Replace policy	Policy updated with literature review through July 7, 2022; references added. Minor editorial refinements to policy statements; intent unchanged
December 2023	Replace policy	Policy updated with literature review through July 7, 2023; reference added. Minor editorial refinements to policy statements; intent unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.